

The Generic Names of Moths of the World

Edited by I. W. B. Nye

Volume 2

Noctuoidea (part):

Arctiidae Lymantriidae

Cocytiidae Notodontidae

Ctenuchidae Strepsimanidae

Dilobidae Thaumetopoeidae

Dioptidae Thyretidae

By Allan Watson, D. S. Fletcher and I. W. B. Nye

British Museum (Natural History)

Trustees of the British Museum (Natural History)
London 1980

© Trustees of the British Museum (Natural History), 1980 Publication Number 811 ISBN 0-565-00811-0

Computer composed and phototypeset by Unwin Brothers Limited The Gresham Press, Old Woking, Surrey, England

This volume should preferably be cited as: Watson, A., Fletcher, D. S. & Nye, I. W. B., 1980, *in* Nye, I. W. B. [Ed.], *The Generic Names of Moths of the World*, volume 2

British Library Cataloguing in Publication Data

The generic names of moths of the world Vol. 2: Noctuoidea (part).

1. Moths—Nomenclature

I. Watson, Allan II. Fletcher, David Stephen

III. Nye, Ian William Beresford

595.7'81'012 QL545

ISBN 0-565-00811-0

Editor's preface

In the whole of the Animal Kingdom about 290 000 genus-group names have been used and of these nearly 10% have been applied to moths and butterflies. It is intended to catalogue all of the 24 000 moth names in the following volumes which have either already been published or are in preparation.

The Generic Names of Moths of the World

Edited by I. W. B. Nye

VOLUME 1 by I. W. B. Nye (published 11th December 1975)

Superfamily NOCTUOIDEA (part)

AGARISTIDAE NOCTUIDAE **NOLIDAE**

VOLUME 2 by A. Watson, D. S. Fletcher and I. W. B. Nye

Superfamily NOCTUOIDEA (part)

ARCTIIDAE COCYTIIDAE CTENUCHIDAE DILOBIDAE DIOPTIDAE LYMANTRIIDAE NOTODONTIDAE STREPSIMANIDAE THAUMETOPOEIDAE THYRETIDAE

VOLUME 3 by D. S. Fletcher (published 3rd December 1979)

Superfamily GEOMETROIDEA

APOPROGONIDAE AXIIDAE CALLIDULIDAE CYCLIDIIDAE DREPANIDAE EPICOPEIIDAE EPIPLEMIDAE GEOMETRIDAE PTEROTHYSANIDAE SEMATURIDAE THYATIRIDAE URANIIDAE

VOLUME 4 by D. S. Fletcher and I. W. B. Nye (in press)

Superfamily BOMBYCOIDEA

ANTHELIDAE
APATELODIDAE
BOMBYCIDAE
BRAHMAEIDAE
CARTHAEIDAE
CERCOPHANIDAE
ENDROMIDAE

EUPTEROTIDAE LASIOCAMPIDAE LEMONIIDAE MIMALLONIDAE OXYTENIDAE SATURNIIDAE SPHINGIDAE

Superfamily CASTNIOIDEA

CASTNIIDAE

Superfamily COSSOIDEA

CHRYSOPOLOMIDAE COSSIDAE DALCERIDAE LIMACODIDAE MEGALOPYGIDAE METARBELIDAE PYROMORPHIDAE RATARDIDAE

Superfamily ZYGAENOIDEA

ANOMOEOTIDAE CYCLOTORNIDAE EPIPYROPIDAE HETEROGYNIDAE HIMANTOPTERIDAE ZYGAENIDAE

Superfamily SESIOIDEA

CHOREUTIDAE DUDGEONEIDAE SESIIDAE

Suborder ZEUGLOPTERA

VOLUME 5 (in preparation by D. S. Fletcher, I. W. B. Nye and S. H. Halsey) Suborder DITRYSIA (part) Superfamilies PYRALOIDEA ALUCITOIDEA VOLUME 6 (in preparation by I. W. B. Nye, D. S. Fletcher and S. H. Halsey) Suborder DITRYSIA (part) Superfamilies GELECHIOIDEA YPONOMEUTOIDEA TINEOIDEA **TORTRICOIDEA** Suborder INCURVARIINA Superfamily **INCURVARIOIDEA** Suborder NANNOLEPIDOPTERA Superfamily **NEPTICULOIDEA** Suborder EXOPORIA Superfamilies **HEPIALOIDEA** MNESARCHAEOIDEA Suborder DACNONYPHA Superfamilies **ERIOCRANIOIDEA NEOPSEUSTOIDEA** Suborder AGLOSSATA Superfamily **AGATHIPHAGOIDEA**

Superfamily

MICROPTERIGOIDEA

Contents

												Page
EDITOR'S PREFAC	Ε.											iii
Synopsis .												vii
AIM, SCOPE AND	STYLE											vii
FORM OF ENTRIES												vii
SEQUENCE												vii
References												vii
DATES .												vii
Type-localiti	ES											vii
Unavailable	NAMES											viii
HOMONYMY.												viii
MISIDENTIFIED T	YPE-SPE	CIES										ix
FAMILY-GROUP N	NAMES											ix
ACKNOWLEDGEM												xi
Additions to T	he Gene	eric N	ames	of M	<i>loths</i>	of the	World	d, vol	UME 1			xi
SUMMARY OF INT	ER-FAM	IILY T	RANSI	EREN	CES							xiii
SUMMARY OF NE	W NAMI	ES ANI) NEV	V TAX	A EST	ABLISI	HED					xiii
ALPHABETICAL C	CATALO	GUE O	F GEN	US-G	ROUP	NAME	s.					1
SYMBOLS USED				•		•	٠		•			1
ABBREVIATION	S USED	FOR F	AMIL	Y-GR	OUP N	AMES	•		•	٠	•	I
CATALOGUE			•	٠			•		•			200
INDEX TO SPECIE	s-grou	P NAM	IES		•			•	•			206



Synopsis

The genus-group names of the families Cocytiidae, Arctiidae, Ctenuchidae, Lymantriidae, Notodontidae, Strepsimanidae, Thaumetopoeidae, Dioptidae, Thyretidae and Dilobidae are listed with their type-species and original bibliographical references. The means of fixation of each type-species together with its type-locality is given. The catalogue includes among its 3650 entries, 93 noctuid and agaristid names supplementary to volume 1 of this series. New genus-group names are established for 46 junior homonyms for which no replacement name could be found and two new nominal genera are established. One new replacement name is established for a type-species. Type-species are designated for 24 genera for which no previous type-species fixation could be found, and 27 genera have been transferred to other families.

Aim, Scope and Style

The aim, scope and style of this catalogue are closely similar to those defined in the first volume of this series. However, it has now been possible to include all relevant names contained in the *Zoological Record* up to and including volume 112 published in 1979, together with names from recent works that have come to our notice up to the end of 1979.

Each nomenclaturally available generic name has been objectively defined by linking it with its valid type-species designation, determined according to the *International Code of Zoological Nomenclature* (Edn 2), 1964, with amendments, 1974, *Bull. zool Nom.* 31: 77–89. Invalid and incorrect designations of type-species which antedate the valid designation are listed and the reasons for their rejection are given.

Subjective synonymy of genus-group names has not been included, except in the case of: (a) a junior homonym having no objective replacement name, or (b) two or more genera having type-species which are subjectively synonymous; such genus-group names have been cross-referenced by the phrase 'See also *Genus* Author, date'. Where the type-species of a genus is a junior subjective synonym, its senior subjective synonym and reference are also included.

Several cases of generic homonymy and objective synonymy have been revealed during the course of this study, such as the use of identical generic names for different taxa in the Diptera and Lepidoptera (Anthomyza Fallén, 1810, and Anthomyza Swainson, 1833). Hitherto wrongly accepted type-species have resulted in, for example, the association of the generic name Antarctia Hübner, [1820], with species of Arctiidae although the originally included species lacked a single arctiid. It is hoped that disentanglement of problems such as these will contribute towards stability in the nomenclature of the Lepidoptera.

Form of Entries

SEQUENCE

The catalogue is arranged alphabetically. Junior homonyms, junior objective synonyms and names not nomenclaturally available are listed chronologically under their potentially valid name, and are also cross-referenced in the main alphabetic sequence.

REFERENCES

The bibliographical source of every generic and specific name has been checked.

Abbreviations of journals follow those of the World List of Scientific Periodicals (Edn 4), 1963–1965. The titles of books and other works are abbreviated as though they were journals, using the 'Select List of Abbreviations Used' in volume 3 of the World List.

The term Commission refers to the International Commission on Zoological Nomenclature and the term *Code* refers to the *International Code of Zoological Nomenclature* (Edn 2), 1964, with amendments 1974, *Bull. zool. Nom.* 31: 77–89.

DATES

Following Recommendation 22A of the *Code*, dates of publication have been cited in square brackets if they have been determined from external evidence, and in parenthesis if determined from evidence in the volume itself. These dates have usually been followed by those printed on the title-page to assist in finding the work in library catalogues.

Dates of publication given in Sherborn's *Index Animalium* and in the *Catalogue of Books*, *Manuscripts*, *Maps and Drawings in the British Museum* (*Natural History*) have been followed unless bibliographical research has shown that changes should be made.

Type-localities

The stated place of origin of the type-material is that given in the original description of a type-species. Minor misspellings have been corrected, and where foreign spellings differ little from the English

COMPARATIVE TABLE OF FAMILY-GROUP NAME USAGES:

Hampson 1898–1920 Cat. Lepid. Phalaenae Br. Mus. 1–3; Suppls 1, 2	Seitz/Draudt/ Rothschild/Gaede 1909–1934 in Seitz, Gross-Schmett. Erde 2, 6, 10, 14	Zerny/Strand/ Bryk/Gaede 1912–1937 Lepid. Cat. 7, 22, 24, 26, 45, 52, 82	Forbes 1939 Bull. Mus. comp. Zool. Harv. 85
ARCTIADAE (2, 3, Suppl. 1) (Included Nolidae) LITHOSIADAE (Suppl. 2) (Included Nolidae)	ARCTIIDAE	ARCTIIDAE	ARCTIIDAE
Arctianae	Arctiinae Callimorphinae Micrarctiinae Nyctemerinae Spilosominae (Included Nolidae and Cocytiidae)	Arctiinae Callimorphinae Nyctemerinae (Included Nolidae)	Arctiinae
Account	Pericopinae	Pericopinae	PERICOPIDAE
_	Hypsinae	AGANAIDAE	HYPSIDAE
Lithosianae	Lithosiinae	Lithosiinae	Lithosiinae
	_	_	EUCHROMIIDAE (Included Amatinae Ctenuchinae Euchromiinae)

spelling, e.g., Brasil (Portuguese spelling) and Brazil (English spelling), these also have been cited in their English form. In all other instances square brackets have been used to enclose the present correct name for a locality, which is then followed by the originally cited name of the locality, for example, '[Ambon I.] Amboina' or '[Sulawesi] Celebes', but modern interpretations have not been given for originally stated localities such as 'Congo' which comprise more than one present-day African state. Square brackets are used also to add information not stated in the original description, for example in the entry '[Malawi]: Fwambo' where we have located the originally cited Fwambo as a village or town in the African state of Malawi. When localities such as 'Rio Grande' have been given in the original description and there is no indication in the paper or book as a whole as to the country of origin of the type-series, inferences have been drawn from the place of capture of specimens judged to be conspecific with the type-series or from the known distribution of species congeneric with the type-species. Such localities are cited, for example, as '[SOUTH AMERICA]: tropical' or other appropriate entry. A type-locality originally given as 'E. Indiis' presents a particular problem because the type-species in question may occur both in eastern India and in the islands of South East Asia (the East Indies). The spellings of modern names follow those used in *The Times Atlas of the World* (Edn 5), 1975.

UNAVAILABLE NAMES

Unavailable names (names that are not nomenclaturally available under the rules of the *Code*) are preceded by a double dagger (‡) and reasons for their rejection are given.

Names recorded in Neave, *Nomenclator zoologicus* as '(pro *A-us* Author, date)' usually lack positive indication as to whether the names are misspellings without nomenclatural availability or whether they are emendations with nomenclatural availability. All such entries have been checked and the correct status given.

Homonymy

Names that have been proposed expressly to replace junior homonyms, and junior objective synonyms that have been used for the same purpose, are referred to in this catalogue as objective replacement

ARCTIIDAE AND CTENUCHIDAE

Forbes 1960 Lepid. New York & neighboring States 4	Inoue 1961 Check List Lepid. Japan 6	Kimball 1965 Arthropods of Florida 1	Family and subfamily names adopted in this catalogue
ARCTIIDAE	ARCTIIDAE	ARCTIIDAE	ARCTIIDAE
Arctiini Callimorphini Phegopterini Utetheisini	Arctiinae Nyctemerinae	Arctiinae	Arctiinae
Pericopinae		PERICOPIDAE	Pericopinae
Hypsinae	AGANAIDAE	_	Aganainae
LITHOSIIDAE	Lithosiinae	-	Lithosiinae
EUCHROMIIDAE	AMATIDAE	AMATIDAE	CTENUCHIDAE (Including Syntominae Ctenuchinae Euchromiinae)

names. Junior subjective synonyms used to replace preoccupied senior synonyms are referred to as subjective replacement names.

Generic names that are junior homonyms have been placed, wherever possible, under their objective replacement names. Some generic homonyms were found to have no objective synonyms but it has been possible to place these as subjective synonyms of other genera by finding, in each case, a genus whose type-species we consider to be congeneric with that of the homonymous genus. There remained, however, 46 for which there was neither an objective nor a subjective replacement name; new names have therefore been proposed for these homonyms. A list of the new names have been included in a 'Summary of new names and new taxa established'.

Misidentified Type-species

Under Article 70 of the *Code* it is to be assumed than an author has correctly identified the nominal species that he

(1) referred to a new genus when he established it, or

(2) designated as type-species of a new or established genus.

In this catalogue there are several genera having type-species now known to be based on misidentified specimens. Under Article 70 such cases should be referred to the Commission to designate whichever species would promote nomenclatural stability. The authors of this catalogue have provisionally chosen type-species that they consider will fulfil this objective.

Family-group Names

A conservative approach has been taken towards family and subfamily classification. When discussing the subfamily classification of the Arctiidae, Watson (1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25) commented that a reclassification that reflects the degree of similarity of only the type-genera and a few of their allies is not a great improvement on the existing classification. The same comments apply at the family level. Areas where comparative work is needed include parts of the Ctenuchidae, for example those which resemble the arctiid genera Eucereon Hübner, Nelphe Herrich-Schäffer and Galethalea

Butler. In the Arctiidae, Forbes' (1960, Lepid. N. Y. & neighboring States 4) subdivision of the family into subfamilies (but termed 'tribes' by Forbes) reflects the phylogenetic structure fairly well but is an oversimplification. Most Old World genera were not dealt with either in 1960 or in an earlier paper by Forbes (1939, Bull. Mus. comp. Zool. Harv. 85) and several genera do not appear to fit into any of his seven subfamilies. The Old World Aganainae, the New World Pericopinae and the more cosmopolitan Lithosiinae each seem to be fairly homogeneous, but the genera now lumped under the present conception of the Arctiinae comprise a much less homogeneous unit and will probably prove to include not only Forbes' Arctiinae, Callimorphinae, Phaegopterinae and Utetheisinae but other currently unrecognized taxa to accommodate genera not discussed by Forbes.

The names used in this catalogue are the oldest available family-group names applied to a particular taxon, except in the case of Lymantriidae discussed below.

Arctiidae Leach, [1815], in Brewster, Edinburgh Encycl. 9: 133 (as Arctides) has had almost universal usage during this century, as also has Lithosiinae Stephens, 1829, Illust. Br. Ent. (Haustellata) 2: 88, and Pericopinae Walker, 1869, Characters undescr. Lepid. Heterocera: 1 (as Pericopidae).

Aganainae Boisduval, 1833, *Nouv. Annls Mus. Paris* 2: 244 (as Aganaides) has had some usage this century but it is unfortunate that its junior synonym Hypsidae Moore, 1878, *Proc. zool. Soc. Lond.* 1878: 3 was used by Hampson [1893], *Fauna Br. India* (Moths) 1) and throughout the volumes edited by Seitz (see table).

Cocytiidae Rothschild, 1915, Lepid. Br. Ornith. Union and Wollaston Expedns Snow Mts S. Dutch New Guinea: 57 (as Cocytianae) was established as a subfamily of the Noctuidae. Rothschild stated 'Sir George Hampson has raised it to the rank of a family, Cocytiadae, and places it after the Lymantriidae just before the Hypsidae'. This must have been an unpublished personal communication, as Hampson, 1918, Novit. zool. 25: 366, cited Rothschild, 1915, as the origin of this family name, and at the same time replaced it by a new name Eucocytiadae Hampson, 1918, as Hampson considered that Cocytia Boisduval was a junior homonym of Cocytius Hübner.

Ctenuchidae Kirby, 1837, in Richardson, Fauna Boreali-Am. 4: 305 is earlier than the family-group names Syntomidae or Amatidae and has had general acceptance in North and South America except by Forbes, who used the name Euchromiidae while admitting that he should have used the earlier name Ctenuchidae. The subfamily classification of the Ctenuchidae is accepted (see table) but its genus-group names have not been placed into subfamilies in this catalogue.

Dilobidae Aurivillius, 1889, *Nordens Fjärilar*: 79, 95 was established as a subfamily of the Arctiidae and was placed, as Dilobinae, on the *Official List of Family-group Names in Zoology* with Name Number 198. Kiriakoff, 1970, *NachrBl. bayer. Ent.* 19: 101, considered that on the basis of the tympanal characters of its type-species, *Diloba* Boisduval, [1837], should be placed in a separate family and raised the Dilobinae to family rank.

Dioptidae Walker, [1865], List Specimens lepid. Insects Colln Br. Mus. 31: 146 has had almost universal use during this century. Josiidae Piepers & Snellen, 1900, Tijdschr. Ent. 43: 26 has not been used since its proposal. A useful summary of the Dioptidae was published by Seitz & Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 499–501.

Lymantriidae Hampson, [1893], Fauna Br. India (Moths) 1: 432. This family was first separated under the name Lariidae Newman, 1832, Sphinx vespiformis, an essay: 40, 44 (as Lariae), based on the nominal genus Laria Schrank, 1802, a junior homonym of Laria Scopoli, 1763. The next name established for this family was Liparidae Boisduval, 1834, Icon. hist. Lépid. nouv. ou peu connus 2: 134 (as Liparides), based on the nominal genus Liparis Ochsenheimer, 1810, a junior homonym of Liparis Scopoli, 1777. Neither of these family-group names may be used as a valid name, the type-genus in each case being a junior homonym. The following names have also been established for the family: (a) Orgyiidae Wallengren, 1861, K. svenska Fregatten Eugenies Resa... C.A. Virgin aren 1851–1853 (Zool.) 1 (10, Lepidoptera): 369 (as Orgyides), based on the nominal genus Orgyia Ochsenheimer, 1810; (b) Dasychiridae Packard, 1864, Proc. ent. Soc. Philad. 3: 331 (as Dasychirae), based on the nominal genus Dasychira Hübner, [1809]; (c) Lymantriidae Hampson, [1893], Fauna Br. India (Moths) 1: 432, based on the nominal genus Lymantria Hübner, [1819]; (d) Leucomidae Grote, 1895, Mitt. Roemermus. Hildesh. 1: 3, based on the nominal genus Leucoma Hübner, 1822; (e) Ocneriidae Meyrick, 1895, Handbk Br. Lepid.: 169 (as Ocneriadae), based on the nominal genus Ocneria Hübner, [1819]; (f) Hypogymnidae Grote, 1896, Mitt Roemermus. Hildesh. 7: 3, based on the nominal genus Hypogymna Billberg, 1820.

Of the family-group names listed in the last paragraph Liparidae was the most widely used during the nineteenth century; Orgyiidae and Dasychiridae had minor usage, but neither name became widely adopted. During the present century Orgyiidae has been used occasionally in contrast with Lymantriidae, which has been used many hundreds of times throughout the world. In North America the use of Liparidae has continued until, in the most recent revision of the family by Ferguson, 1978, in Dominick et al., Moths Am. N. of Mexico 22 (2), the family name Lymantriidae has been adopted. Because of the overwhelming worldwide use of the name Lymantriidae an application has been submitted by D. S. Fletcher, I. W. B. Nye and D. C. Ferguson to the International Commission on Zoological Nomenclature requesting them to rule that the family-group name Lymantriidae Hampson, [1893] is to be given nomenclatural precedence over the family-group names Orgyiidae Wallengren, 1861, and Dasychiridae Packard, 1864, when applied to the same taxon.

Notodontidae Stephens, 1829, *The Nomenclature of British Insects*: 39 has had almost universal use during this century.

Strepsimanidae Meyrick, 1930, Exot. Microlepid. 4: 10 was established for a small oriental family originally placed in the Microlepidoptera. It has been transferred to the Noctuoidea by Hodges, 1978, in Dominick et al., Moths Am. N. of Mexico 6: (1): 11.

Thaumetopoeidae Aurivillius, 1889, Nordens Fjärilar: 68, 75 was established as a subfamily of the Notodontidae and was raised to family rank by Staudinger, 1901, in Staudinger & Rebel, Cat. Lepid. palaearct. Faunengeb. (1): 113. Cnethocampinae Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 362, a family-group name not used outside Australia, is a junior objective synonym of Thaumetopoeidae, both family-group names being based on the same type-species, Phalaena processionea Linnaeus, 1758. Thaumetopoeidae has had wide use and was adopted by Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E).

Thyretidae Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 99 (as Thyretinae) was established as a subfamily of the Zygaenidae. The genus *Thyretes* Boisduval, 1847, was included in the Syntomidae, now Ctenuchidae, by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 137. Thyretidae was unnecessarily proposed as a new family name by Kiriakoff, 1948, *Bull. Annls Soc. ent. Belg.* 84: 271.

Acknowledgements

We are grateful to the many colleagues in Britain and abroad who have helped us during the preparation of this catalogue. They include Dr Pastor Alayo D., Havana; Dr E. Berio, Genoa; Mr F. M. Brown, Colorado Springs; Dr I. F. B. Common, Canberra; Dr W. Dierl, Munich; Mr R. E. Dietz, Berkeley; Mr J. P. Donahue, Los Angeles; Dr D. C. Ferguson, Washington; Prof. J. G. Franclemont, Ithaca; Dr L. A. Gozmány, Budapest; Dr H. J. Hannemann, Berlin; Dr H. Inoue, Fujisawa; Prof. S. G. Kiriakoff, Ghent; Mr I. Lansbury, Oxford; Dr C. Lemaire, Paris; Prof. O. Mielke, Curitiba; Dr I. Persson, Stockholm; Dr E. L. Todd, Washington; Dr L. Vári, Pretoria and Dr P. Viette, Paris.

Our colleagues in the BMNH, Dr G. S. Robinson, Dr K. S. O. Sattler and Mr W. G. Tremewan, have frequently come to our aid with helpful discussions. Thanks are extended also to Mrs S. D. M. Knapp for typing into the computer much of the data, and we are particularly indebted to Miss S. H. Halsey for her constructive comments and great care in up-dating the catalogue to conform to the style required for computerized type-setting. Our librarian Miss P. M. Gilbert has been ever helpful, as have Miss M. Creer, Mr D. French, Mr C. G. J. MacPherson and Dr D. B. Williams who prepared and operated the computer programmes. The photographs used in the frontispiece were taken by Mr P. York of our Photographic Unit.

Additions to The Generic Names of Moths of the World, Volume 1

The Generic Names of Moths of the World, volume 1, published on 11th December 1975, contained the genus-group names of the families Noctuidae, Agaristidae, and Nolidae. The following genus-group names of Noctuidae and Agaristidae are mostly additions to those in volume 1. There are no additional genus-group names in the Nolidae. The names, references, type-species and other information have been included in the present catalogue.

Acanthopolia Boursin, 1943
Agoma Kiriakoff, 1976
Antigodasa Kiriakoff, 1974
Antoculeora Ichinosé, 1973
Arattatha Janse, 1917
Arboricornis Hampson, 1908
‡Aretype; Hampson, 1926
‡Astephana; Agenjo, 1946
‡Austrocarea; Holloway, 1976
Austrocarea Holloway, 1977
Batuana Laporte, 1976
Bifrontipta Berio, 1974
Bracharthrum Hampson, 1895
• '

India `
South Africa
Madagascar
India
South Africa
s Hampson, 1894
th, 1903
Holloway, 1977
Australia
Ethiopia
Ethiopia

Noctuidae, Emendation, see Bracharthron Hampson, 1891

xii	GENERIC NAMES C	OF MOTHS OF THE WO	RLD
Brandtaxia Boursin, 1943	N	octuidae, Acronictinae	Iran
Callicereon Butler, 1882	N	octuidae, Acronictinae	Madagascar
Cerviplusia Chou & Lu, 197		octuidae, Plusiinae	China
Chazaria Moore, 1881		octuidae, Noctuinae	Turkey
Chichimeca Hogue, 1963		octuidae, Acronictinae	U.S.A.
Cladocera Rambur, 1858	N	octuidae. Junior homonym,	see <i>Episema</i> Ochsenheimer, 1816
Codalithia Holloway, 1979		octuidae, Acontiinae	New Caledonia
Cofinipacia Holloway, 1979		octuidae, Hypeninae	New Caledonia
Colnettia Holloway, 1979		octuidae, Ophiderinae	New Caledonia
Cualitemoca Hogue, 1963		octuidae, Acronictinae	Mexico
Cypliocampa Harris, 1869		garistidae	U.S.A.
Cyptonychia Druce, 1899 Cyptonychia Hampson, 1900		octuidae, Acronictinae octuidae, Acronictinae	Mexico Mexico
Dargeochaeta Laporte, 1973		octuidae, Acronictinae	Cameroun
Deinhugia Laporte, 1974		octuidae, Ophiderinae	Cameroun
Diphteramonia Berio, 1961		octuidae, Acronictinae	Netherlands
Dorstiana Laporte, 1974		octuidae, Hadeninae	Et hiopia
Dufayella Capușe, 1972	No	octuidae, Sarrothripinae	U.S.S.R.
Dusponera Dognin, 1914		octuidae, Hypeninae	Ecuador
Epischausia Kiriakoff, 1976		garistidae	Kenya
Epithisanotia Kiriakoff, 1976		garistidae	America, tropical
‡Epystenie; Hübner, [1831]		garistidae. Misspelling, see E	
Eriopana Kiriakoff, 1974	A	garistidae. Objective synonyr	1895
‡Focillstis; Nye, 1975		octuidae. Misspelling, see Fo	
Godasa Walker, [1865]		garistidae	Sierra Leone
Gonepteronia Kenrick, 1917		octuidae, Ophiderinae	Madagascar
Henituerta Kiriakoff, 1976 ‡Hoplopseustis; Swinhoe, 192		garistidae octuidae. Misspelling, see <i>He</i>	Somali Republic
‡Hoplopseustrias; Swinhoe, 192		octuidae, Misspelling, see He	
‡Hypolaclina; Walker, 1865		octuidae, Misspelling, see Hy	
Hypoloclima Felder, 1861		octuidae, Sarrothripinae	Indonesia
<i>Hypliilara</i> Kirby, 1897		octuidae. Emendation, see H	
Hypotuerta Kiriakoff, 1976		garistidae	Nigeria
Iranada Wiltshire, 1977		octuidae, Ophiderinae	U.S.S.R.
Kara Matsumura, 1925 ‡Kerasia; Pagenstecher, 1909		octuidae. Junior nomonym, soctuidae. Misspelling, see <i>Ke</i>	see Parascotia Hübner, [1825]
Koffoleania Laporte, 1977		octuidae, Acronictinae	Ethiopia
Mafana Viette, 1979		octuidae, Ophiderinae	Madagascar
Marcipalina Pelletier, 1978		octuidae, Ophiderinae	Gabon
Matarum Viette, [1976]	No	octuidae, Acronictinae	Réunion
Medlerana Laporte, 1973		octuidae, Noctuinae	Nigeria
Mekrania Brandt, 1941		octuidae, Hypeninae	Iran
Michelliana Laporte, 1976		octuidae, Hadeninae garistidae	Ethiopia U.S.A.
Neotuerta Kiriakoff, 1976 Nesaegocera Kiriakoff, 1974		garistidae	Madagascar
Nocthadena Laporte, 1976		octuidae, Hadeninae	Ethiopia
Pachycerma Van Duzee, 189		octuidae. Objective synonym	
Palaeocoleus Robinson, 1975		octuidae, Hypeninae	Fiji
Palpidia Dyar, 1898		octuidae, Ophiderinae	U.S.A.
Pararothia Kiriakoff, 1974		garistidae Objective symonys	Madagascar
Parathisanotia Kiriakoff, 197		garistidae. Objective synonyi	1869
Pectinidia Holloway, 1977		octuidae, Ophiderinae	Norfolk Island
‡Peniphegostola; Hampson, 1		garistidae. Misspelling, see P	
Pempligostola Strand, 1909		garistidae	Madagascar
Pentelia Kiriakoff, 1976 Philareta Moore, 1881		garistidae. Junior homonym, octuidae, Noctuinae	Turkey
Pseudagoma Kiriakoff, 1975		garistidae	Mozambique
Pseudocoarica Holloway, 197		octuidae, Ophiderinae	New Caledonia
Pseudotuerta Kiriakoff, 1976		garistidae	Uganda ·
Ptochosiphla Meyrick, 1933		octuidae, Hypeninae	Fiji
Puriplusia Chou & Lu, 1974		octuidae, Plusiinae	Japan
Rougeotia Laporte, 1974		octuidae, Hadeninae	Ethiopia
Rougeotiana Laporte, 1974 Sarbissa Kiriakoff, 1976		octuidae, Ophiderinae garistidae	Ghana Fiji
‡Schausiana; Kiriakoff, 1976		garistidae. Misspelling, see S	
Schausilla Kiriakoff, 1974		garistidae	Madagascar
Sciatta Walker, 1869	N	octuidae, Ophiderinae	Zaire
Sclerogenia Ichinosé, 1973		octuidae, Plusiinae	Japan
Scotogramma Edwards, 1887	N	octuidae, Hadeninae	U.S.A.

Sergiusia Nye nom. n. Shensiplusia Chou & Lu, 1974 ‡Sideridia; Nye, 1975 Steneugoa Hampson, 1914 Synalamis Dognin, 1912 Synorthodes Franclemont, 1976 Syrrusoides Laporte, 1972 Tholocoleus Robinson, 1975 Thysanoplusia Ichinosé, 1973 Trichofeltia Barnes & Benjamin,

1929 September

Tuertella Kiriakoff, 1976

Agaristidae Mozambique Noctuidae, Plusiinae China Noctuidae. Misspelling, see Sideridis Hübner, [1821] Noctuidae, Sarrothripinae New Guinea Noctuidae, Ophiderinae Colombia Noctuidae, Hadeninae U.S.A. Noctuidae, Acronictinae Madagascar Noctuidae, Hypeninae Fiji Noctuidae, Plusiinae China Noctuidae. Objective synonym, see Trichofeltia

McDunnough, [1929 April]

Agaristidae

Tanzania

Summary of Inter-family Transferences

The following nominal genera are transferred from one family to another.

Nominal Genus From Abella Walker, 1862 Lymantriidae Antarctia Hübner, [1820] Arctiidae Belenmiastis Hampson, 1901 Arctiidae Caridarctia Hampson, 1901 Noctuidae Chatracharta Walker, 1862 Lasiocampidae Corticoniis van Eecke, 1924 Lymantriidae Cymaroa Hampson, 1905 Lymantriidae Arctiidae Cyptonychia Druce, 1899 Cyptonychia Hampson, 1900 Arctiidae Dactylorlivncha Hampson, [1893] Lymantriidae Dasychirinula Hering, 1926 Lymantriidae Arctiidae Diospage Walker, 1854 Godasa Walker, [1865] Arctiidae Hypolochma Felder, 1861 Lymantriidae Kawiella Roepke, 1943 Lymantriidae Lasioceros Bethune-Baker, 1904 Arctiidae Lymphorta Walker, [1858] Noctuidae Microstola Lower, 1920 Xyloryctidae Arctiidae Nesotropha Turner, 1926 Pachylaelia Butler, 1874 Lymantriidae Arctiidae Palpidia Dyar, 1898 Cossidae Pernambis Schaus, 1920 Piesta Billberg, 1820 Arctiidae Ptochosiphla Meyrick, 1933 Pyralidae Pyralopsis Boisduval, 1870 Arctiidae Arctiidae Sciatta Walker, 1869 Trichromia Hübner, [1819] Ctenuchidae

To Lasiocampidae Saturniidae Ctenuchidae Arctiidae Lymantriidae Anthelidae Arctiidae Noctuidae Noctuidae Limacodidae Lasiocampidae Ctenuchidae Agaristidae Noctuidae Pyralidae Notodontidae Notodontidae Arctiidae Glyphipterigidae Arctiidae Noctuidae Pyromorphidae Oecophoridae Noctuidae Pyralidae Noctuidae Arctiidae

Summary of New Names and New Taxa Established

The following list summarizes the junior homonyms for which new objective replacement names have been proposed in this catalogue.

Preoccupied name

Acutia Kaye, 1919 Alope Walker, 1855 Antongila Kiriakoff, 1958 Baeomorpha Turner, 1940 Blacodes Kiriakoff, 1959 Breyeria Janse, 1920 Calosonia Geyer, 1832 Chlorostola Hampson, 1898 Closteroides Kiriakoff, 1977 Coniortodes Collenette, 1955 Conosia Hampson, 1900 Crinola Leech, 1890

Cymella Felder, 1874

Replacement name

Atucia Watson nom. n. Olepa Watson nom. n. Exantongila Fletcher nom. n. Hobapromea Watson nom. n. Neoblacodes Fletcher nom. n. Eubreveria Fletcher nom. n. Osmacola Watson nom. n. Euchlorostola Watson nom. n. Closterellus Fletcher nom. n. Melgona Nye nom. n. Euconosia Watson nom. n. Halsevella Nye nom. n. Myclela Watson nom. n.

Preoccupied name

Diplonyx Hampson, 1900 Dulichia Walker, 1855 Empodisma Kiriakoff, 1964 Gynnopoda Felder, 1874 Heliocaes Turner, 1940 Ira Neumoegen, 1890 Jacksonia Collenette, 1937 Jorgensenia Schaus, 1924 Lavinia Kiriakoff, 1962 Libyssa Dognin, 1911 Mardara Walker, 1865 Micragra Hampson, 1898 Microdota Dognin, 1906 Mirobriga Walker, 1864 Ocneropsis Kozhanchikov, 1950 Paraplocia Aurivillius, 1909 Pareugoa Daniel, 1951 Pectinopliora Janse, 1920 Pentelia Kiriakoff, 1976 Pliara Walker, 1854 Pinheya Kiriakoff, 1973 Pseudoscolia Hampson, 1914 Pterygopterus Butler, 1876 Pydna Walker, 1856 Sankurua Collenette, 1960 Sarapus Walker, 1855 Scytale Felder, 1874 Sphenoptera Felder, 1874

Spilarctia Staudinger, 1891

Tricholepis Hampson, 1891

Zama Walker, 1865

Taeniopteryx Janse, 1920 Thyone Walker, 1854

Replacement Name

Didymonyx Watson nom. n. Knappetra Nye nom. n. Boscawenia Fletcher nom. n. Apeplopoda Watson nom. n. Cheliosea Watson nom, n. Cubira Watson nom. n. Jacksoniana Nye nom. n. Geriojennsa Watson nom. n. Eulavinia Fletcher nom. n. Eudognina Fletcher nom. n. Ramadra Nye nom. n. Micragrella Watson nom. n. Trocodima Watson nom. n. Birgorima Watson nom. n. Gissarus Nye nom. n. Calpoparia Watson nom. n. Apogurea Watson nom. n. Eujansea Fletcher nom. n. Sergiusia Nye nom. n. Aphra Watson nom. n. Pinlievata Nye nom. n. Parascolia Watson nom. n. Timalus Watson nom. n. Eupydna Fletcher nom. n. Kunusara Nye nom. n. Asparus Watson nom. n. Clystea Watson nom. n. Asphenoptera Watson nom. n. Nebrarctia Watson nom. n. Metopteryx Fletcher nom. n. Eutlivone Watson nom. n. Chiretolpis Watson nom. n. Zamana Fletcher nom. n.

In addition to the 46 objective replacement names listed above, the following two new nominal genera have been established for species previously combined with genus-group names proposed after 1930, which, although described, contained more than one species and were not accompanied by a definite fixation of a type-species; such genus-group names are unavailable under Article 13(b) of the *Code*. As these names have not been made available since they were proposed, the authors were contacted and the requirements of the *Code* have been fulfilled in the relevant entries in this catalogue.

Unavailable name ‡Neobourquinia Köhler, 1943 ‡Pseudohyaleucerea Rego Barros & Machado, 1971 New nominal genus

Neobourquinia Köhler gen. n.

Pseudohyaleucerea Rego Barros & Machado gen. n.

One type-species was found to be a primary homonym and, as there is neither an objective nor a subjective synonym that could be used as a replacement name, a new name has been established for this species.

Preoccupied name Dasyclira pumila Butler, 1882 Replacement name
Noliproctis milupa Nye nom. n.

Alphabetical Catalogue of Genus-group Names

SYMBOLS USED

‡. A name not nomenclaturally available under the Code.

]. Added information derived usually from external evidence.

[1825] 1816. Date of publication derived from external evidence followed by the title-page date.

ABBREVIATIONS USED FOR FAMILY-GROUP NAMES

AGAN Aganainae, Arctiidae Lyma Lymantriidae ARCT Arctiinae, Arctiidae NOTO Notodontidae

COCY Cocytiidae PERI Pericopinae, Arctiidae
CTEN Ctenuchidae STRE Strepsimanidae
DILO Dilobidae THAU Thaumetopoeidae
DIOP Dioptidae THYRET Thyretidae

LITH Lithosiinae, Arctiidae

ABAKABAKA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: LYMA

Type-species: Calliteara fuliginosa Saalmüller, 1884, Lepid. Madagascar: 193, by original designation. MADAGASCAR: N.W., Nossi-Bé Island.

ABELLA Walker, 1862. See Clathe Walker, 1855.

LASIOCAMPIDAE

ABNORMIPTERUS Orfila, 1935, Revta Soc. ent. argent. 7: 177.

Type-species: Sphecosoma abnormis Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 159, fig.71, by original designation. [Brazil]: lower Amazon, Massauary.

ABROCHIA Herrich-Schäffer, 1855. See Pseudosphex Hübner, 1818.

ABROCHOCIS Dyar, 1914, Proc. U.S. natn. Mus. 47: 170.

Type-species: Talara esperanza Schaus, 1911, Ann. Mag. nat. Hist. (8) 7: 365, by original designation. Costa Rica: Esperanza; Tuis; Sixola; and Guapiles.

ABYNOTHA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 479.

Type-species: *Liparis preussi* Mabille & Vuillot, 1892, *Novit. Lepid.*: 57, pl.9, fig. 5, by original designation. [WEST AFRICA]: [coast] côte occidentale d'Afrique.

ACANTHARCTIA Aurivillius, 1899, Ent. Tidskr. 20: 234, 241.

Type-species: Acantharctia nivea Aurivillius, 1899, ibidem 20: 242, by original designation. Congo.

ACANTHOFRONTIA Hampson, 1910, Cat. Lepid. Phalaenae Br. Mus. 10: xvii, 3, 644.

Type-species: *Acanthofrontia lithosiana* Hampson, 1910, *ibidem* 10: 644, fig.177, by original designation. [Kenya]: Athi-ya-Mawe.

Acanthofrontia was established in the Noctuidae; it was placed in the Arctiidae by Nye, 1975, Generic Names Moths World 1: 18.

ACANTHOPOLIA Boursin, 1943, Z. wien. ent. Ges. 28: 339.

Type-species: Apamea cuprina Moore, 1881, Proc. zool. Soc. Lond. 1881: 345, pl.38, fig.2, by original designation. Lectotype (but fixed as 'Type') &, INDIA: Sikkim, (MNHU, Berlin) by Boursin, 1943, ibidem 28: 338.

Acanthopolia in the Noctuidae, Hadeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

ACASINA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 140.

Type-species: Somera acasia Schaus, 1928, Proc. U.S. natn. Mus. 73 (19): 75, by original designation. INDIA: Assam.

ACCO Bethune-Baker, 1904, Novit. zool. 11: 418.

Type-species: Acco bicolora Bethune-Baker, 1904, ibidem 11: 418, pl.5, fig.21, by original designation. New Guinea: [Papua New Guinea], Dinawa and Aroa R[iver].

ACERBIA Sotavalta, 1963, Suom. hyönt. Aikak. 29: 263.

Type-species: Bombyx alpina Quensel, 1802, in Acerbi, Travels through Sweden, Finland and Lapland to North Cape, 1798 & 1799 2: 253, by original designation. LAPLAND.

ACHAERA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 337.

Type-species: Chadisra ochribasis Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 461, by original designation. [Kenya]: Laitokitok.

ACHEPYDNA Matsumura, 1925, Zool. Mag. Tokyo 37: 405.

Type-species: *Pydna formosicola* Strand, 1916, *Arch. Naturgesch.* 81 (A) 12: 158, by original designation (but cited as ‡formosana Matsumura, an incorrect subsequent spelling and an incorrect authorship). TAIWAN: Suisharyo.

Matsumura used Achepydna formosicola (Strand) as a heading, with a reference, but designated ‡'Achepydna formosana Matsumura' as type-species. In his 'List of the Formosan Notodontidae' Matsumura cited, on page 408, only the species formosicola Strand in his new genus Achepydna.

ACHERDOA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 451.

NOCTUIDAE

Type-species: *Acherdoa ferraria* Walker, 1865, *ibidem* 32: 451, by monotypy. [U.S.A.]: Florida.

Acherdoa was established in the Notodontidae; it was placed in the Noctuidae by Dyar, [1903] 1902, Bull. U.S. natn. Mus. 52: 195.

ACHROOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 90, 337.

Type-species: Achroosia nuda Hampson, 1900, ibidem 2: 338, fig.239, by original designation. BRAZIL: [Espirito Santo] Espiritu Santo.

ACIDON Hampson, 1896, Fauna Br. India (Moths) 4: 462.
NOCTUIDAE

Type-species: Acidon paradoxa Hampson, 1896, ibidem 4: 462, fig. 247, by original designation. BHUTAN.

Acidon was established in the Notodontidae; it was trans-

ferred to the Noctuidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

ACLONOPHLEBIA Butler, 1898, Proc. zool. Soc. Lond. 1898: 428.

Type-species: Aclonophlebia flavinotata Butler, 1898, ibidem 1898: 428, pl.32, fig.8, by original designation. [Kenya] British East Africa: between Voi and Ndi, Marago ya Fundi.

ACLYTIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 123.

Type-species: Sphinx heber Cramer, 1780, Uitlandsche Kapellen (Papillons exot.) 3: 169, 175, pl.287, fig.A, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 160. SURINAM.

S. heber is dated from the wrapper of the part and not from the title-page of the volume, 1782.

ACLYTOPHANES Turner, 1940, Proc. R. Soc. Qd 51: 80.

Type-species: Heliosa aedumena Turner, 1922, Proc. R. Soc. Vict. 35: 29, by monotypy. AUSTRALIA: northern Queensland, near Cairns.

ACMESHACHIA Matsumura, 1929, Insecta matsum. 4: 38,

Type-species: Acmeshachia takamukui Matsumura, 1929, ibidem 4: 39, pl.1, fig.18, by original designation (but cited as ‡takamukue, an incorrect (of a multiple) original spelling). TAIWAN: Horisha.

ACNISSA Turner, 1902 December, Trans. Proc. R. Soc. S. Aust. 26: 180.

Type-species: Acnissa pyrrhias Turner, 1902, ibidem 26: 180, by monotypy. Syntypes 18, 29, Australia: Queensland, Townsville, (ANIC, Canberra).

A. pyrrhias is a junior subjective synonym of Haplopseustis

erythrias Meyrick, 1902 April.

Acnissa was established in the Lymantriadae, now Lymantriidae; it was retained in the same family but placed as a junior subjective synonym of Haplopseustis by Turner, 1903, ibidem 27: 15; Haplopseustis, with Acnissa in synonymy, was placed in the Noctuidae, Erastrianae now Acontiinae, by Turner, 1920, ibidem 44: 166.

See also Haplopseustis Meyrick, 1902 April.

ACONTIA Ochsenheimer, 1816, Schmett. Eur. 4: 91. NOCTUIDAE

Type-species: Noctua solaris [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 90, by subsequent designation by Duponchel, 1829, in Godart & Duponchel, Hist. nat. Lépid. Papillons Fr. 7 (2): 72. [AUSTRIA]: Vienna district.

Acontia has been used as the original generic name for species now in the Arctiidae.

ACRASIELLA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg.

Type-species: Hyperaeschra curvilinea Swinhoe, 1907, Ann. Mag. nat. Hist. (7) 19: 207, by original designation. [South Africa]: [Natal], Durban.

ACRASIODES Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg.

Type-species: Fentonina io Viette, 1954, Mém. Inst. scient. Madagascar (E) 5: 75, pl.2, fig.7, by original designation. MADAGASCAR: southern, Ankarampotsy.

ACREAGRIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.83, fig.2. Type-species: Acreagris correbioides Felder, 1874, ibidem 2

(Abt.2): pl.83, fig.2, by monotypy. MEXICO: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: [1].

A junior homonym of Acreagris Koch & Berendt, 1854, in Berendt, Die im Bernstein befindlichen organischen Reste der Vorwelt 1 (2): 6, 123, pl.17, fig.157, - Collembola. The only objective replacement name Felderia Kirby, 1892, is itself a junior homonym (see below). However, the type-species of Acreagris is currently considered to be congeneric with Pyromorpha dimidiata Herrich-Schäffer, [1854], the type-species of Pyromorpha Herrich-Schäffer, [1854] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: wrapper, pl. [43], fig. 222, which is therefore available as a subjective replacement name.

Acreagris when established was not placed in a family; it was placed in the Arctiidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 163; and its objective replacement name Felderia Kirby, 1892, was placed in the Zygaenidae by Jordan, 1913, in Seitz, Gross-Schmett. Erde 6: 22.

FELDERIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera

Type-species: Acreagris correbioides Felder, 1874, by mon-

otypy (of Acreagris Felder, 1874).

Felderia was proposed as the objective replacement name for Acreagris Felder, 1874, but is itself a junior homonym of Felderia Walsingham, 1887, Trans. ent. Soc. Lond. 1887: 165, - Lepid., Tineidae.

Felderia was established in the Arctiidae; it was placed in the Zygaenidae by Jordan, 1913, in Seitz, Gross-Schmett. Erde 6: 22.

‡ACRIBIA Boisduval, 1870. See Acribia Felder, 1874. PERI

ACRIBIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.103, fig.14.

Type-species: Cocastra melanchroia Boisduval, 1870, Considérations Lépid. Guatemala: 88 (but attributed to Felder, an incorrect authorship), by monotypy. GUATEMALA.

‡ACRIBIA Boisduval, 1870, Considérations Lépid. Guatemala: 89 (but attributed to Felder, an incorrect authorship).

‡Acribia Boisduval, 1870, was first published in the synonymy of Cocastra Boisduval, 1870, ibidem: 88; and is unavailable under the Code, Article 11(d), as it has not been treated subsequently as an available name of Boisduval, 1870. Boisduval attributed ‡Acribia to Felder, and presumably had access to Felder's plates and legends (to which Boisduval refers) prior to their publication in 1874.

ACRIDOPSIS Butler, 1876, J. Linn. Soc. (Zool.) 12: 418.

Type-species: Eucereon latifascia Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1639, by original designation (but cited as ‡Eucerea, an incorrect subsequent spelling). [BRAZIL]: Para.

ACROCTENA Saalmüller, 1884, Lepid. Madagascar: 244.

Type-species: Acroctena fissura Saalmüller, 1884, ibidem: 244, pl.5, fig. 70, by monotypy. MADAGASCAR: Nossi-Bé Island.

ACROSEMA Meigen, 1830. See Phalera Hübner, [1819]. NOTO

ACSALA Benjamin, 1935, Can. Ent. 67: 195. Type-species: Acsala anomala Benjamin, 1935, ibidem 67:

196, by original designation. [U.S.A.]: Alaska, Yukon.

Acsala was established in the Arctiidae; it was transferred to the Lymantriidae by Fergusson, 1978, in Dominick, Moths Am. N. of Mexico 22 (2): 15.

‡ACTAEA; misspelling. See Mitradaemon Butler, 1878.

ACTEA Walker, 1854. See Mitradaemon Butler, 1878. DIOP

ACUTIA Kaye, 1919. See Atucia Watson nom. n. CTEN

ACYCLANIA Dognin, 1911, Hétérocères nouv. Am. Sud 3: 15.

Type-species: Acyclania tenebrosa Dognin, 1911, ibidem 3: 15, by original designation. ARGENTINA: Parana, San Ignacio.

ACYMBA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 235.

Type-species: *Noctua spectabilis* Tauscher, 1806, *Mém. Soc. Nat. Univ. imp. Moscou* 1: 212, fig., by monotypy. [U.S.S.R.] Russia: [southern] australiori.

See also Lacydes Walker, 1855.

PALPARCTIA Spuler, 1906, Schmett. Eur. 2: 133.

Type-species: Noctua spectabilis Tauscher, 1806, by monotypy.

A junior objective synonym of Acymba Rambur, 1866.

VOLGARCTIA Alphéraky, 1908, Horae Soc. ent. Ross. 38: 606.

Type-species: Noctua spectabilis Tauscher, 1806, by monotypy.

A junior objective synonym of Acymba Rambur, 1866.

ACYPHAS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 797.

Type-species: Acyphas fusca Walker, 1855, ibidem 4: 798, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 472. [AUSTRALIA] New Holland: and [Tasmania] Van Dieman's Land.

ANEPA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 478.

Type-species: Acyphas fusca Walker, 1855, by original designation.

A junior objective synonym of Acyphas Walker, 1855.

ADDUA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 254.

Type-species: Addua inclusa Walker, 1862, ibidem (3) 1: 254, by monotypy. Brazil: Rio de Janeiro neighbourhood.

A. inclusa is a junior subjective synonym of Pericopis subapicalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 352.

ADELPHONEURA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.31.

Type-species: *Adelphoneura nerias* Felder, 1874, *ibidem* 2 (Abt.2): pl.105, fig.31, by monotypy. [Brazil.]: Amazon [River], cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 1.

A. nerias is a junior subjective synonym of Josia prolifera Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2:

See also Glissa Walker, [1865].

‡ADOLPHONEURA; Bryk, 1930, in Strand, Lepid. Cat. 42:

An incorrect subsequent spelling of Adelphoneura Felder, 1874.

ADETONEURA Collenette, 1933, Stylops 2: 44. LYMA Type-species: Adetoneura lentiginosa Collenette, 1933, ibidem 2: 44, by original designation. FIJI ISLANDS.

ADITES Moore, [1882] 1882-3, Lepid. Ceylon 2: 61. LITH Type-species: Doliche hilaris Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 530, by monotypy. [SRI LANKA] Ceylon.

ADLULLIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 392.

Type-species: *Adlullia lunifera* Walker, 1865, *ibidem* 32: 392, by original designation. [SULAWESI] Celebes: Menado.

‡ADOLPHONEURA; misspelling. See Adelphoneura Felder, 1874. DIOP

ADOXOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 87, 397.

Type-species: *Adoxosia excisa* Hampson, 1900, *ibidem* 2: 398, fig. 306, by original designation. BRAZIL: Rio [de] Janeiro.

ADRALLIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 509.

Type-species: Adrallia bipunctata Walker, 1865, ibidem 32:

510, by monotypy. [South Africa]: Natal.

Adrallia was established in the Limacodidae; it was placed in the Arctiidae by Hering, 1928, in Seitz, Gross-Schmett. Erde 14: 473; and was placed in the Thaumetopoeidae by Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 22.

‡ADRAPSA; misspelling. See Adrepsa Moore, 1879. LITH

ADREPSA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 37.

Type-species: *Adrepsa stilbioides* Moore, 1879, *ibidem* (1): 38, pl.2, fig.20, by monotypy. 1NDIA: Assam.

‡ADRAPSA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 422.

An incorrect subsequent spelling of Adrepsa Moore, 1879.

ADSCITA Retzius, 1783, in Degeer, Genera Species Insect.: 8, 35. ZYGAENIDAE

Type-species: Adscita turcosa Retzius, 1783, ibidem: 35, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 84. Type-Locality: not stated, [Europe].

A. turcosa is a junior subjective synonym of Sphinx statices Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 495.

Adscita is included here as it is associated with ‡Chrysaor, the first use in the Lepidoptera of a name later established as Chrysaor Hübner, [1809], in the Ctenuchidae.

‡CHRYSAOR Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) Opinion 97: 19, also idem, 1954, Opin. Decl. int. Commin zool. Nom. 6 (Opinion 278): 140.

Only included species: Sphinx statices Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 495.

AEDOEA Turner, 1899, Trans. Proc. R. Soc. S. Aust. 23: 10.

Type-species: *Aedoea monochroa* Turner, 1899, *ibidem* **23**: 10, by monotypy. AUSTRALIA: Brisbane.

A. monochroa is a junior subjective synonym of Lithosia decreta Butler, 1877, Trans. ent. Soc. Lond. 1877: 351.

AEMENE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 541.

Type-species: Aemene taprobanis Walker, 1854, ibidem 2: 542, by monotypy. [Sri Lanka] Ceylon.

See also *Panassa* Walker, 1865, and *Autoceras* Felder, 1874.

AEMILIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1:

Type-species: Ameles rubriplaga Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 712, by monotypy (of Ameles Walker, 1855). VENEZUELA.

Aemilia was established as the objective replacement name for Ameles Walker, 1855.

AMELES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 711.

Type-species: Ameles rubriplaga Walker, 1855, by monotypy. A junior homonym of Ameles Burmeister, [1838], Handb. Ent. 2: 531, – Insecta, Orthoptera. The objective replacement name is Aemilia Kirby, 1892.

AEOLOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 89, 404.

Type-species: Aeolosia multipunctata Hampson, 1900, ibidem 2: 405, fig.315, by original designation. Burma: Hsipaw.

‡OEOLOSIA; Hampson, 1901, J. Bombay nat. Hist. Soc. 13: 502.

An incorrect subsequent spelling of Aeolosia Hampson, 1900.

‡CLEOLOSIA; Bethune-Baker, 1904, Novit. zool. 11: 424. An incorrect subsequent spelling of Aeolosia Hampson, 1900.

AETHALIDA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 320.

Type-species: Aethalida distinguenda Walker, [1865] 1864, ibidem 31: 321, by monotypy. [SULAWESI] Celebes: Menado.

AETHOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 85, 385.

Type-species: Aethosia ectrocta Hampson, 1900, ibidem 2: 385, fig. 299, by original designation. HAITI.

AETHRIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 120.

Type-species: *Sphinx haemorrhoidalis* Stoll, 1790, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* (Aanhangsel): 53, 183, pl.12, fig.1, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 152. SURINAM.

S. haemorrhoidalis is dated from the wrapper of the part

and not from the title-page of the volume, 1791.

MESOLASIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xvi, 28, 339.

Type-species: Sphinx haemorrhoidalis Stoll, 1790, by original designation.

A junior objective synonym of Aethria Hübner, [1819].

AFILIA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 261 (key), 319.

Type-species: *Afilia cinerea* Schaus, 1901, *ibidem* **1901**: 320, by original designation. Mexico: Orizaba, Guadalajara.

AFRIDA Möschler, 1886, Abh. senckenb. naturforsch. Ges. 14: 30.

Type-species: Afrida tortriciformis Möschler, 1886, ibidem 14: 30, by monotypy. JAMAICA.

AFROARCTIA Toulgoët, 1978, Nouv. Revue Ent. 8: 223.

Type-species: Spilosoma sjostedti Aurivillius, 1899, Ent. Tidskr. 20: 240, by original designation (but cited by Toulgoët, 1978, as ‡sjoestedti, an incorrect subsequent spelling). [Cameroun] Kamerun.

AFROCERURA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 213.

Type-species: Cerura leonensis Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 457, by original designation. SIERRA LEONE.

AFROGLUPHISIA Kiriakoff, 1970, Ann. Transv. Mus. 26 (9): 194. NOTO

Type-species: Afrogluphisia demas Kiriakoff, 1970, ibidem 26 (9): 195, fig.15, by original designation. South Africa: [Transvaal], Potgietersrus district, Limburg.

AFROPLITIS Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 201.

Type-species: *Hoplitis dasychirina* Gaede, 1928, *in* Seitz, *Gross-Schmett. Erde* 14: 421 (the objective replacement name for *Hoplitis dasychiroides* Rothschild, 1917; see below) [Namibia] S.W. African Protectorate: [Tsumeb] Isumeb.

Hoplitis dasychirina was established by Gaede as the objective replacement name for Hoplitis dasychiroides Rothschild, 1917, Novit. zool. 24: 250, pl.5, fig.5, then a junior secondary homonym of Stauropus dasychiroides Butler, 1898, Proc. zool. Soc. Lond. 1898: 433, pl.32, fig.12, when both were included in the genus Hoplitis Hübner, [1819]. Under the Code, Article 59(b)(i), a junior secondary homonym rejected before 1961 is permanently rejected and cannot be restored unless the use of the replacement name is contrary to existing usage.

AFROPTERYX Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 363.

Type-species: *Lophopteryx angulata* Gaede, 1928, *in* Seitz, *Gross-Schmett. Erde* **14**: 408, pl.69, row b, by original designation. [GHANA] Goldküste.

AFROPYDNA Kiriakoff, 1961, Bull. Annls Soc. r. ent. Belg. 97: 102.

Type-species: *Pydna indistincta* Gaede, 1928, *in* Seitz, *Gross-Schmett. Erde* 14: 437, pl.72, row c, by original designation. [Tanzania] Ost-Afrika: the type-specimen in the Zoological Museum in Berlin is labelled 'D.O. Afrika, Meinicke leg.' (Dr H.J. Hannemann, 27.vii.1976, personal communication).

Afropydna was established as a subgenus of Xanthodonta Gaede, 1928; it was raised to generic level by Kiriakoff, 1964,

in Wytsman, Genera Insect. 217 (A): 90.

AGABRA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 136.
NOCTUIDAE

Type-species: *Agabra trilineata* Walker, 1862, *ibidem* 6: 137, by monotypy. Borneo: Sarawak.

Agabra was established in the Notodontidae; it was transferred to the Noctuidae by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 37.

AGAGLES White, 1841, in Grey, Jls two Exped. Disc. N.W. and W. Australia . . . 1837, 38, and 39 2: 482.

Type-species: Agagles amicus White, 1841, ibidem 2: 482, by monotypy. Australia: Western Australia, King George Sound.

‡AGLAGES; Agassiz, 1846, Nomencl. zool. (Nomina syst. Generum Lepid.): 3.

An incorrect subsequent spelling of Agagles White, 1841.

AGLAGUS Agassiz, 1847, Nomencl. zool. (Index univl.): 11. An unjustified emendation of Agagles White, 1841, (but the latter cited as ‡Aglages White, an incorrect subsequent spelling).

The 'Index universalis' was published as Fascicle 12 of the Nomenclator zoologicus; Aglagus is dated from the wrapper

of Fascicle 12, not from its title-page dated 1846.

AGALTARA Toulgoët, 1979, Bull. Soc. ent. Fr. 83: 223. ARCT Type-species: Agaltara nebulosa Toulgoët, 1979, ibidem 83: 223, by original designation. RWANDA: Nyungwe Forest, Kitabi.

AGANAIS Boisduval, 1832. See Psephea Billberg, 1820.

‡AGANOPIS Herrich-Schäffer, 1855. See Aganopis Herrich-Schäffer, [1856].

AGANOPIS Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): 12, wrapper, pl.88, figs 501, 502.

Type-species: Aganopis subquadrata Herrich-Schäffer, [1856] 1850-1858, ibidem 1 (1): 12, wrapper, pl.88, figs 501, 502, by monotypy. [BANGLADESH]: [Sylhet] Silhet.

A. subquadrata is a junior subjective synonym of Hypsa orbicularis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 445.

See also Peridrome Walker, 1854.

‡AGANOPIS Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 92.

A name not nomenclaturally available as it was published without a description, and the only included name 'subquadrata m. Exot. f. 501. 502.' was also without a description or figure at that date.

‡ERIOCRYPTA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 92; [1856] 1850-1858, Samml. neuer oder

wenig bekannter aussereur. Schmett. 1 (1): 12.

A name not nomenclaturally available as it was published without a description, and the species for which it was proposed, ‡'longipennis m.', was also without a description. Herrich-Schäffer, [1858] 1850-1858, ibidem 1 (1): 70, again used ‡Eriocrypta without a description and suggested that the species for which it was proposed [‡longipennis] may represent the female of Aganopis subquadrata Herrich-Schäffer, [1856], the type-species of Aganopis Herrich-Schäffer, [1856].

‡Eriocrypta was cited as a junior synonym of Aganopis Herrich-Schäffer, [1856], by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 384, and as a junior synonym of Peridrome Walker, 1854, by Gaede, 1932, in Strand, Lepid. Cat. 52: 5.

AGAPE Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.4.

Type-species: Agape cyanopyga Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.4, by monotypy. [Philippines]: Luzon, [Indonesia]: [Moluccas], [Ambon I.] Amboina, both localities cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: [1].

A. cyanopyga Felder is a junior subjective synonym of Hypsa chloropyga Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 455.

‡AGAPEA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 426.

An incorrect subsequent spelling of Agape Felder, 1874.

‡AGAPEA; misspelling. See Agape Felder, 1874. AGAN

AGAPHTHORA Meyrick, 1886, *Proc. Linn. Soc. N.S.W.* (2) 1: 773, 774.

Type-species: Agaphthora melanora Meyrick, 1886, ibidem (2) 1: 774, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 102. AUSTRALIA: Queensland, Cape York.

A. melanora is a junior subjective synonym of Syntomis guttulosa Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 73.

AGAPOSOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.99, fig.24. ARCT

Type-species: Agaposoma marcescens Felder, 1874, ibidem 2 (Abt.2): pl.99, fig.24, by monotypy. [BRAZIL]: Amazon R., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: [1].

AGARAEA Herrich-Schäffer, 1855 December 31, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.76, fig.442.

Type-species: Agaraea longicornis Herrich-Schäffer, 1855, ibidem 1 (1): wrapper, pl.76, fig.442, by monotypy. BRAZIL.

Agaraea and A. longicornis are dated from the wrapper of the Heft and not from the title-page of the volume, 1850-1858.

‡AGORAEA; Herrich-Schäffer, 1855 December [31], Syst.

Bearbeitung Schmett. Eur. 6: 100; [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 20.

An incorrect spelling of Agaraea Herrich-Schäffer, 1855.

‡AGAREA; Schaus, 1892, Proc. zool. Soc. Lond. 1892: 288. An incorrect subsequent spelling of Agaraea Herrich-Schäffer, 1855.

‡AGAREA; misspelling. See Agaraea Herrich-Schäffer, 1855.

AGEROCHA Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 10.

Type-species: Agerocha eone Hübner, [1831] 1825, ibidem 3: 10, pl.[72], figs 417, 418, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 256. BRAZIL.

AGKONIA Dognin, 1894, Annls Soc. ent. Belg. 38: 128. LITH Type-species: Agkonia pega Dognin, 1894, ibidem 38: 128, by monotypy. [ECUADOR]: near Loja, Zamora.

‡AGLAGES; misspelling. See Agagles White, 1841. ARCT

AGLAGUS Agassiz, 1847. See Agagles White, 1841. ARCT

AGLAOSOMA Scott, 1864, Aust. Lepid. and their Transformations 1 (2): 14.

Type-species: Aglaosoma lauta Scott, 1864, ibidem 1 (2): 15, pl.5, by monotypy. AUSTRALIA.

Aglaosoma when established was not placed in a family; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 469; and was transferred to the Thaumetopoeidae by Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 39.

A. lauta is a junior subjective synonym of Teara variegata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 851.

AGLOSSOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 87, 247.

Type-species: Aglossosia flavimarginata Hampson, 1900, ibidem 2: 247, by original designation. [SOUTH AFRICA]: Natal, Mooi R. and Newcastle.

AGOMA Kiriakoff, 1976, Tierreich 98: 70. AGARISTIDAE Type-species: Aegocera trimenii Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.107, fig.15, by original designation. Type(s) [SOUTH] AFRICA: Natal, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 1, (BMNH).

Agoma in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡AGORAEA; misspelling. See Agaraea Herrich-Schäffer, 1855.

AGRISIUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 723.

Type-species: Agrisius guttivitta Walker, 1855, ibidem 3: 723, by monotypy. INDIA: northern.

AGUNAIX Schaus, 1898 June, Jl N.Y. ent. Soc. 6: 139. CTEN Type-species: Agunaix lacrumans Schaus, 1898 June, ibidem 6: 139, by monotypy. Peru.

Agunaix originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson, 1898 [December], as the name for a new genus.

AGUNAIX Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xvii, 30, 378.

Type-species: Agunaix lacrumans Schaus, 1898 June, by

monotypy (but cited as ‡lacrimans, an incorrect subsequent spelling).

A junior homonym of Agunaix Schaus, 1898 June, Jl N.Y. ent. Soc. 6: 139, — Lepid., Ctenuchidae. The objective replacement name is Agunaix Schaus, 1898 June.

AGUNAIX Hampson, 1898 [December]. See Agunaix Schaus, 1898 Junc.

AGYLLA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 552.

Type-species: Agylla fasciculata Walker, 1854, ibidem 2: 552, by monotypy. VENEZUELA.

AGYLLOIDES Strand, 1912, Arch. Naturgesch. 78 (A) 7: 186.

Type-species: Agylloides asurella Strand, 1912, ibidem 78 (A) 7: 187, by original designation. Togo: 'Bismarkberg'.

AGYRTA Hübner, [1820] 1816, Verz. bekannter Schmett.: 177.

Type-species: *Phalaena micilia* Cramer, 1779, *Uitlandsche Kapellen (Papillons exot.)* 3: 62, 175, pl.228, fig.G, by PRES-ENT DESIGNATION (see below). SURINAM.

P. micilia is dated from the wrapper of the part and not from

the title-page of the volume, 1782.

Agyrta was established for two nominal species, Sphinx auxo Linnaeus, 1767, and Phalaena coelestina Stoll, 1781 (not ‡caelestina as cited by Hübner). Under auxo, Hübner referred to his earlier interpretation of the species ([1817], Samml. exot. Schmett. 1: pl.[185], figs 1-4) and placed Phalaena micilia Cramer as a synonym of auxo.

The true Sphinx auxo Linnaeus, 1767, is a species in the Zygaenidae Chalcosiinae described from China, whereas the specimens figured as this species by Hübner represent two South American species in the Ctenuchidae. Figs 1 and 2 are of a species without a name, and figs 3 and 4 are of Phalaena micilia Cramer.

Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 121, treated micilia and auxo as different species within Agyrta, and designated auxo as the type-species. This treatment and designation (including the misidentification of auxo) was followed by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 469. Agyrta is a genus having a misidentified type-species and so should be referred to the Commission under Article 70(a) of the Code.

Tremewan, 1973, Bull. Br. Mus. nat. Hist. (Ent.) 28: 116, reviewed the case and concluded that the type-species of Agyrta was Sphinx auxo Linnaeus sensu Hübner, [1817], Samml. exot. Schmett. 1: pl.[185], figs 1, 2, by subsequent designation by Kirby, 1892, ibidem 1: 121. The taxon represented by figs 1 and 2 is without a name, and the group of species related to it requires a thorough revision.

It is here proposed that the Commission should be asked to rule that the type-species of Agyrta Hübner, [1820] is Phalaena micilia Cramer, 1779. The latter was one of the names originally included in Agyrta and was correctly represented by Hübner in figs 3 and 4. With micilia as type-species the usage of Agyrta will be maintained in the sense accepted since it was established.

CORYDALLA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1650.

Type-species: Corydalla cryptoleuca Walker, 1856, ibidem 7: 1650, by monotypy. [Brazil]: Para.

A junior homonym of *Corydalla* Vigors, 1826, *Zool. J.* 2: 397, – Aves. There is no objective replacement name but *C. cryptoleuca* is currently considered to be congeneric with *Phalaena micilia* Cramer, the proposed type-species of *Agyrta* Hübner, [1820]. The latter is therefore available for use as a subjective replacement name.

EVAGRA Boisduval, 1870, Considérations Lépid. Guatemala: 87.

Type-species: *Phalaena micilia* Cramer, 1779, by PRESENT DESIGNATION.

A junior objective synonym of Agyrta Hübner, [1820].

AGYRTIDIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xvi, 25, 315.

Type-species: Eucyane uranophila Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1874, by original designation. Peru, [Brazil]: [São Paulo] St Paul.

AGYRTIOLA Gaede, 1926, Dt. ent. Z. 1926: 136. CTEN Type-species: Agyrtiola niepelti Gaede, 1926, ibidem 1926: 136, by original designation. Brazil: [Bahia], 'Olivença'.

ALANDRIA Gaede, 1926, Dt. ent. Z. 1926: 129. CTEN Type-species: Alandria coeruleipuncta Gaede, 1926, ibidem 1926: 129, by original designation. BOLIVIA: 'S[ão] Jacinto', [SOUTH AMERICA]: 'Cillutincara'.

ALASTOR Boisduval, 1869. See Nadata Walker, 1855.

ALATANADATA Strand, 1912, Arch. Naturgesch. 78 (A) 6: 166.

Type-species: Alatanadata latipennis Strand, 1912, ibidem 78 (A) 6: 167, by original designation. CAMEROUN: Bibundi.

‡ALBARACINA; misspelling. See Albarracina Staudinger, 1883.

ALBARRACINA Staudinger, 1883, Stettin. ent. Ztg 44: 180.

Type-species: *Bombyx korbi* Staudinger, 1883, *ibidem* **44**: 179, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* **1**: 838. Spain: Provincia de Teruel, Albarracin.

B. korbi is a junior subjective synonym of Bombyx warionis Oberthür, 1881, Etudes d'Entomologie 6: 75, pl.2, fig.6

Albarracina was proposed at the end of the original description of Bombyx korbi in case a separate genus was required for it and other species.

‡ALBARACINA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 426.

An incorrect subsequent spelling of *Albarracina* Staudinger, 1883.

ALENOPHALERA Strand, 1912, Arch. Naturgesch. 78 (A) 6: NOTO

Type-species: *Phalera variegata* Aurivillius, 1904, *Ark. Zool.* **2** (4): 4, fig.12, by original designation. CAMEROUN.

ALEPISTA Toulgoët, 1976, Bull. Soc. ent. Fr. 81: 189. LITH Type-species: Alepista irregularis Toulgoët, 1976, ibidem 81: 189, fig.13, by original designation. MADAGASCAR: W., 12 km au Nord-Est de Sakaraha, réserve forestière du Zombitsy.

ALEXICLES Grote, 1883, Trans. Kans. Acad. Sci. 8: 37, 46.

Type-species: Alexicles aspersa Grote, 1883, ibidem 8: 37, 46, by original designation. U.S.A.: New Mexico.

ALIMALA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 132.

Type-species: *Alimala limacodoides* Walker, 1862, *ibidem* 6: 133, by monotypy. Borneo: Sarawak.

Alimala was established in the Notodontidae; it was placed in the Noctuidae by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 38.

ALINA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 230.

Type-species: *Dasychira ochroderoea* Mabille, 1898, *Annls Soc. ent. Fr.* **66**: 223, by original designation. MADAGASCAR: Diego-Suarez.

ALLATA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 140.
NOTO

Type-species: *Allata argentifera* Walker, 1862, *ibidem* 6: 140, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 621. BORNEO: Sarawak.

ALLOCHRISTA Roepke, 1946, Tijdschr. Ent. 87: 85. LITH Type-species: Allochrista toxopei Roepke, 1946, ibidem 87: 86, fig., by monotypy. [SULAWESI] Celebes: Todjambu.

ALLODONTA Staudinger, 1887, in Romanoff, Mém. Lépid. 3: 223.

Type-species: *Notodonta tristis* Staudinger, 1887, *ibidem* 3: 223, pl.13 [recte 17], fig.7, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Askold [Island].

N. tristis is a junior subjective synonym of Notodonta plebeja Oberthür, 1880, Etudes d'Entomologie 5: 65, pl.8, fig.7

Allodonta was established as a subgenus of Notodonta Ochsenheimer, 1810.

ALLODONTINA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 409.

Type-species: *Allodonta apicalis* Kiriakoff, 1974, *ibidem* 17: 410, text-fig.25, pl.4, fig.5, by original designation. SUMATRA: northern, Deli, Dolok Merangir.

ALLODONTOIDES Matsumura, 1922, Zool. Mag. Tokyo 34 402: 523.

Type-species: *Hyperaeschra discoidalis* Matsumura, 1920, *ibidem* 32: 148, by original designation. TAIWAN.

H. discoidalis is a junior subjective synonym of Hyperaeschra dentata Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 165.

See also Polyaeschra Kiriakoff, 1967.

ALLOTOMA Roepke, 1944, Natuurh. Maandbl. 33: 4. LYMA Type-species: Allotoma cornifrons Roepke, 1944, ibidem 33: 4, fig.18, by monotypy. BORNEO: south-eastern.

Allotoma was established in the Notodontidae; it was transferred to the Lymantriidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

ALOA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 699.

Type-species: *Phalaena lactinea* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 58, 149, pl.133, fig.D, by subsequent designation by Moore, [1883] 1882-3, *Lepid. Ceylon* 2: 74. [Java]: [Djakarta] Batavia.

P. lactinea is dated from the wrapper of the part and not from the title-page of the volume, 1779.

ALOPE Walker, 1855. See Olepa Watson nom. n. ARCT

ALPENUS Walker, 1855, List Specimens lepid. Insects Colln Br.. Mus. 3: 686.

Type-species: Alpenus aequalis Walker, 1855, ibidem 3: 686,

by monotypy. WEST AFRICA.

A. aequalis is a junior subjective synonym of Phalaena maculosa Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 156, 250. Dated from the wrapper of the part and not from the title-page of the volume, 1782.

ALPHAEA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 683.

Type-species: *Alphaea fulvohirta* Walker, 1855, *ibidem* 3: 684, by monotypy. Type-locality: not stated, [INDIA].

ALPIS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1094.

MEGALOPYGIDAE

Type-species: Alpis defoliata Walker, 1855, ibidem 5: 1094, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 541. MEXICO.

Alpis was established in the Notodontidae; it was placed in the Megalopygidae by Berg, 1882, An. Soc. cient. argent. 13: 264.

ALTIMAENAS Dyar, 1913, Proc. U.S. natn. Mus. 45: 642.

Type-species: *Altimaenas tapina* Dyar, 1913, *ibidem* **45**: 642, by monotypy. Peru: Coropuna.

ALYSIDOTA Agassiz, 1847. See Halysidota Hübner, [1819].

ALYTARCHIA Wallengren, 1863, Wien. ent. Monatschr. 7: 147.

Type-species: *Euchelia amanda* Boisduval, 1847, *in* Delegorgue, *Voyage Afr. australe* 2: 597, by subsequent designation by Hampson, 1905, *Ann. S. Afr. Mus.* 3: 416. [SOUTH AFRICA]: [Zululand] Pays de Amazoulous, [WEST AFRICA]: 'Cote de Guinée' and 'Pays de Galam'.

Alytarchia was redescribed as new by Wallengren in 1865, K. svenska Vetensk Akad. Handl. 5 (4): 47.

AMALEA Herrich-Schäffer, 1855. See Nudaria Haworth, [1809].

AMALODETA Meyrick, 1889, Trans. ent. Soc. Lond. 1889: 462.

Type-species: *Amalodeta electraula* Meyrick, 1889, *ibidem* **1889**: 463, by monotypy. New Guinea: Papua New Guinea, Port Moresby.

AMANA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 588 (key), 661.

Type-species: Amana angulifera Walker, 1855, ibidem 3: 662, by monotypy. Type-locality: not known. Specimens associated with the holotype & in BMNH are from N. India and Bhutan.

Amana was established in the Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 474; and was placed in the Epiplemidae by Swinhoe, 1894, Trans. ent. Soc. Lond. 1894: 164.

AMASTUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 723.

Type-species: *Phaegoptera collaris* Herrich-Schäffer, [1853] 1850-1858, *Samml. neuer oder wenig bekannter aussereur.* Schmett. 1 (1): wrapper, pl.15, fig.65, by monotypy. Venezuela.

AMATA Fabricius, 1807, Magazin Insektenk. (Illiger) 6: 289.

Type-species: Zygaena passalis Fabricius, 1781, Species Insect. 2: 159, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 59. Type-locality: not stated, [India or Sri Lanka].

COENOCHROMIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 121.

Type-species: Zygaena passalis Fabricius, 1781, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 59, 86 (but designated as 'Sphinx creusa, Cram.'; see below).

A junior objective synonym of Amata Fabricius, 1807.

Coenochromia was established having two originally included nominal species. The first of these was cited by Hübner as '1316. Coenochromia Creusa Linn. Syst. Sph. 39. Cram. 248.F.' Hampson, 1898, ibidem 1: 304 treated Sphinx creusa

Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 494, as denoting a valid species in the genus Euchromia Hübner, [1819]. On page 86, however, Hampson treated Sphinx creusa Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 94, 174, pl.248, fig. F, as a junior synonym of Zygaena passalis Fabricius, 1781. Hampson thus indicated that Cramer had misidentified S. creusa Linnaeus, and that Hübner had followed Cramer. This is a case of a misidentified type-species which should be referred to the Commission under Article 70(a) of the Code.

AMATHES Gistl, 1848. See Nyctemera Hübner, [1820].

AMATISSA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 138.

PSYCHIDAE

Type-species: Amatissa inornata Walker, 1862, ibidem 6: 138, by monotypy. Borneo: Sarawak.

Amatissa was established in the Notodontidae; it was placed in the Psychidae by Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 292.

AMAXIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 721.

Type-species: *Amaxia pardalis* Walker, 1855, *ibidem* 3: 721, by monotypy. [Brazil]: Amazon, [Tefé] Ega.

AMBADRA Moore, 1883, Proc. zool. Soc. Lond. 1883: 16.

Type-species: Eumeta rafflesi Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 430, by subsequent designation by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 92. JAVA.

Ambadra was established as the objective replacement name

for Corma Moore, 1881.

CORMA Moore, 1881, Proc. zool. Soc. Lond. 1881: 326.

Type-species: Eumeta rafflesi Moore, [1860], by subsequent designation by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 92 (for Ambadra Moore, 1883).

A junior homonym of *Corma* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 124, – Lepid., Zygaenidae. The objective replacement name is *Ambadra* Moore, 1883.

AMBINA Kiriakoff, 1962, Tijdschr. Ent. 105: 242. NOTO Type-species: Ichthyura spissicornis Mabille, 1900, Annls Soc. ent. Fr. 68: 723, by original designation. [MADAGASCAR]: Antongil.

AMBLYTHYRIS Mabille, 1879, Bull. Soc. philomath. Paris (7) 3: 137.

Type-species: Amblythyris radama Mabille, 1879, ibidem (7) 3: 137, by subsequent designation by Hampson, 1900, Ann. S. Afr. Mus. 2: 60, 61, but cited as Aganais vitripennis Blanchard, 1861, a nominal species not originally included in Amblythyris. Hampson, on page 61, placed vitripennis as the senior synonym of radama, a nominal species originally included in Amblythyris. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. MADAGASCAR.

A. radama is a junior subjective synonym of Chelonia madagascariensis Boisduval, 1847, in Delegorgue, Voyage Afr.

australe 2: 598.

See also Phryganeomorpha Wallengren, 1858.

AMBRYLLIS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 750.

Type-species: Glaucopis boisduvallii van der Hoeven, 1840, in van der Hoeven & de Vriese, Tijdschr. Natuurl. Gesch. Physiol. 7: 279, by monotypy. Brazil.

‡AMBRYSIIS; Burmeister, 1878, Descr. phys. Répub. Argent. 5 (1): 437.

An incorrect subsequent spelling of Ambryllis Walker, 1855.

‡AMBRYSIIS; misspelling. See Ambryllis Walker, 1855.

AMELES Walker, 1855. See Aemilia Kirby, 1892. ARCT

AMELETA Turner, 1940, Proc. R. Soc. Qd 51: 79. LITH Type-species: Ameleta panochra Turner, 1940, ibidem 51: 79, by monotypy. AUSTRALIA: N. Queensland, Kuranda.

AMERILA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 725.

Type-species: Sphinx astreus Drury, 1773, Illust. nat. Hist. exot. Insects 2: index, 49, pl.28, fig.4 (but cited as ‡astraea, an incorrect subsequent spelling), by subsequent designation by Hampson, 1900, Ann. S. Afr. Mus. 2: 60 (but cited as ‡astreas, an incorrect subsequent spelling). [INDIA]: Bengal [formerly included much of northern India].

CANOPUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 747.

Type-species: Canopus bubo Walker, 1855, ibidem 3: 747, by subsequent designation by Hampson, 1900, Ann. S. Afr.

Mus. 2: 60. [ZAIRE] Congo.

A junior homonym of *Canopus* Fabricius, 1803, *Syst. Rhyngotorum*: 127, – Insecta, Hemiptera. There is no objective replacement name but the type-species of *Canopus* Walker, 1855, is currently considered to be congeneric with the type-species of *Amerila* Walker, 1855. The latter is therefore available for use as a subjective replacement name.

V AMMALO Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 622.

Type-species: Ammalo fervidus Walker, 1855, ibidem 3: 622, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 222. Type-locality: not stated, [South America].

A. fervidus is a junior subjective synonym of Phalaena helops Cramer, 1775, Uitlandsche Kapellen (Papillons exot.) 1: 113, pl.72, fig.C. Dated from the wrapper of the part and not from the title-page of the volume, 1779.

AMMATHO Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 759.

Type-species: Ammatho cuneonotatus Walker, 1855, ibident 3: 759, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 308. [Sri Lanka] Ceylon.

AMMOBIOTA Wallengren, 1885. See Eucharia Hübner, [1820].

AMPHEKES Collenette, 1936, Ann. Mag. nat. Hist. (10) 17: 332.

Type-species: *Amphekes gymna* Collenette, 1936, *ibidem* (10) 17: 333, by original designation. [CHINA]: North Yunnan, Likiang.

V AMPHELARCTIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 42.

Type-species: Automolis priscilla Schaus, 1911, Ann. Mag. nat. Hist. (8) 7: 183, by original designation. Costa Rica: Juan Viñas.

AMPHICALLIA Aurivillius, 1900, Ent. Tidskr. 20: 235.

Type-species: *Callimorpha bellatrix* Dalman, 1823, *Analecta Ent.*: 50, by original designation. Type-locality: not stated, [SOUTH AFRICA].

AMPHIPHALERA Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 477.

Type-species: Amphiphalera leuconephra Hampson, 1910, ibidem (8) 5: 478, by original designation. [GHANA] Gold Coast: Volta River.

AMPHISSA Walker, 1855. See Phissama Moore, 1858.

AMPLICINCIA Field, 1950, Proc. U.S. natn. Mus. 100: 312.

Type-species: *Cincia pallida* Butler, 1878, *Proc. zool. Soc. Lond.* **1878**: 484, by original designation. JAMAICA.

AMSACTA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 804.

Type-species: Amsacta marginalis Walker, 1855, ibidem 4:

805, by monotypy. SIERRA LEONE.

Amsacta was established in the Liparidae, now Lymantriidae; it was transferred to the Arctiidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498.

AMSACTARCTIA Berio, 1938, Annali Mus. civ. Stor. nat. Giacomo Doria 58: 197.

Type species: Amsacta radiosa Pagenstecher, 1903, Jb. nassau. Ver. Naturk. 56: by original designation. [ETHIOPIA]: Darassum [Approx. 5°50'N 41°10'E].

AMSACTOIDES Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19 (1): 53.

Type-species: Creatonotus formosae Strand, 1915, Ent. Mitt. 4: 13, by original designation. [TAIWAN] Formosa: Kankau.

AMYCLES Herrich-Schäffer, [1855]. See Anycles Walker, 1854.

‡AMYCLES; misspelling. See Anycles Walker, 1854.

AMYOPS Karsch, 1895, Ent. Nachr. Berlin 21: 362.

Type-species: Amyops ingens Karsch, 1895, ibidem 21: 362, pl.3, fig.4, by original designation. Togo: Misahöhe.

See also Melebaeas Distant, 1903.

ANACE Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1719.

THYRETIDAE

Type-species: Anace rubra Walker, 1856, ibidem 7: 1720, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 221, but cited as Automolis lateritia Herrich-Schäffer, [1855], a nominal species not originally included in Anace. Kirby placed lateritia, incorrectly, as the senior synonym of rubra, a nominal species originally included in Anace. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. [SOUTH AFRICA]: Cape.

Anace was established in the Arctiidae; it was placed, correctly, in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 47, but used with an incorrect typespecies Anace perpusilla Walker, 1856, designated by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 143.

ANAEMOSIA Hampson, 1918, Novit. zool. 25: 101. LITH Type-species: Anaemosia albida Hampson, 1918, ibidem 25: 101, by original designation. [MALAWI] Br. C. Africa: Mt Mlanje.

ANAENE Dyar, 1914, Proc. U.S. natn. Mus. 47: 171. LITH Type-species: Anaene spurca Dyar, 1914, ibidem 47: 171, by original designation. PANAMA: La Chorrera, Canal Zone, Corozal.

‡ANAGINA; misspelling. See Anagnia Walker, 1854.

ANAGNIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 446.

Type-species: *Hypsa subfascia* Walker, 1854, *ibidem* 2: 446, by monotypy. 1NDIA: northern, EAST INDIES.

Anagnia was established as a subgenus of Hypsa Hübner, [1819].

‡ANAGINA; Swinhoe, 1890, Trans. ent. Soc. Lond. 1890: 179. An incorrect subsequent spelling of Anagnia Walker, 1854.

ANALAMA Kiriakoff, 1960, Revue fr. Ent. 27: 189. NOTO Type-species: Scalmicauda perinetensis Kiriakoff, 1958, Mém. Inst. scient. Madagascar (E) 9: 225, fig.7, by original designation. MADAGASCAR: eastern, Perinet district, [Analamazaotra] Analamazoatra forest.

ANAPHE Walker, 1855 [August 11], List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 856.

THAU

Type-species: *Anaphe reticulata* Walker, 1855 [August 11], *ibidem* **4**: 856, by monotypy. [SOUTH AFRICA]: [Durban] Port Natal.

See also Arctiomorpha Herrich-Schäffer, 1855 December 31.

‡ARAPHA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 429.

An incorrect subsequent spelling of Anaphe Walker, 1855.

ANAPHELA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 751.

Type-species: Anaphela luctifera Walker, 1855, ibidem 3: 751, (but generic and specific names attributed to Boisduval, an incorrect authorship), by monotypy. MADAGASCAR.

A. luctifera is a junior subjective synonym of Agarista stellata Guérin-Méneville, [1844], Icon. Règne anim. Cuvier: 493. See also Arnophila Mabille, 1879.

ANAPHLEBIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2):,pl.102, fig.36. CTEN Type-species: Anaphlebia caudatula Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.36, by monotypy. [Brazil]: Amazon River, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: [1].

ANAPHLEPS Rothschild, 1916, Novit. zool. 23: 326. LITH Type-species: Anaphleps angustipennis Rothschild, 1916, ibidem 23: 326, by monotypy. New Guinea: Papua New Guinea, Vulcan Island.

ANAPHOSIA Hampson, 1903, Ann. Mag. nat. Hist. (7) 11: 344.

Type-species: *Anaphosia cyanogramma* Hampson, 1903, *ibidem* (7) 11: 344, by monotypy. [RHODESIA]: Mashonaland, Mpodzi River.

ANAPISA Kiriakoff, 1952, Revue Zool. Bot. afr. 46: 174.

Type-species: *Apisa monotica* Holland, 1893, *Psyche, Camb.* 6: 395, by original designation. [GABON]: valley of the [Ogooué] Ogové River.

ANATOLIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.10.

Type-species: Anatolis subtilis Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.10, by monotypy. Type-Locality: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 1, as 'Columbia?' [Colombia]. The series associated with the type-specimen in the BMNH is from Venezuela.

Anatolis, when established, was not placed in a family; it was placed in the Zygaenidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 86; and was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 418.

ANATOLMIS Packard, 1864, Proc. Essex Inst., Salem, Mass. 4: 45.

Type-species: Anatolmis grotei Packard, 1864, ibidem 4: 47, by monotypy. U.S.A.: Colorado, Pike's Peak.

ANAULOSIA Schaus, 1911, Ann. Mag. nat. Hist. (8) 7: 368.

Type-species: Anaulosia impolita Schaus, 1911, ibidem (8) 7: 368, by monotypy. Costa Rica: Guapiles.

ANAXILA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 810.

Type-species: Anaxila plagiata Walker, 1855, ibidem 4: 810. by monotypy. TYPE-LOCALITY: not known, [WEST AFRICA].

ANAXITA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 748. ARCT

Type-species: Anaxita decorata Walker, 1855, ibidem 3: 748, by monotypy. MEXICO.

See also Echemis Boisduval, 1870.

ANCHYNEURA Felder, 1861, Sber. Akad. Wiss. Wien 43 (Abt.1): 33. LYMA

Type-species: Anchyneura praeusta Felder, 1861, ibidem 43 (Abt.1): 33, by monotypy. [INDONESIA]: [Moluccas], [Ambon 1.] Amboina.

ANDALA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 774.

Type-species: Andala unifascia Walker, 1855, ibidem 3: 774, by monotypy. Type-locality: stated to be 'East Indies' but Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 181 designated a lectotype from India. All the specimens in BMNH are from India.

ANDOCIDIA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 51.

Type-species: Andocidia tabernaria Kiriakoff, 1958, ibidem **66**: 51, pl.4, fig.1, pl.5, fig.7, pl.9, fig.30, pl.10, fig.39, by original designation. [GHANA] Cote d'Or: [Bibiani] Bibianaha.

ANDRACA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 581.

Type-species: Andraca bipunctata Walker, 1865, ibidem 32:

582, by monotypy. [INDIA]: Hindostan.

Andraca was established in the Bombycidae; it was placed in the Notodontidae by Eecke, 1921, Tijdschr. Ent. 64 (Verslag): xlix; and was transferred to the Eupterotidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

ANDRENIMORPHA Butler, 1876, J. Linn. Soc. (Zool.) 12: 382.

Type-species: Glaucopis xanthogastra Perty, [1833] 1830-1834, in Martius, Delectus Anim. articulatorum Bras.: 156, fig., by original designation. BRAZIL: Minarum.

‡ANDRIA Hübner, [1806]. See Cerura Schrank, 1802. NOTO

ANDRIA Hübner, 1822. See Cerura Schrank, 1802. NOTO

ANDROCHARTA Felder & Felder, 1862. See Euclera Herrich-Schäffer, 1855. CTEN

ANDROSTIGMA Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 13, 82.

Type-species: Didugu albicosta Hampson, 1891, ibident 8: 5, 53, pl.140, fig.17, by monotypy. INDIA: Nilgiri platcau.

ANEPA Swinhoe, 1903. Sec Acyphas Walker, 1855. LYMA

ANESTIA Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 691, 745. LITH

Type-species: Anestia ombrophanes Meyrick, 1886, ibidem (2) 1: 746, by monotypy. AUSTRALIA: Victoria, Melbourne.

‡ANEUROCAMPA; misspelling. See Anurocampa Herrich-Schäffer, [1854].

ANEXOTAMOS Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 157.

Type-species: Anexotamos flavibasis Hering, 1926, ibideni 14: 157, pl.22, row h, by monotypy. TYPE-LOCALITY: stated as not known presumably East African lake district. [Specimens in BMNH from Uganda and Zaire].

The date of publication of the part of Seitz containing page 157 is stated on page 153 as 5.xi.1926 in both the German and

the English editions.

ANGUSTIALA Bryk, 1950, Ark. Zool. 42 (A) 19: 3. Type-species: Angustiala cryptocephala Bryk, 1950, ibidem 42 (A) 19: 4, pl.1, fig.4, by original designation. Burma:

A. cryptocephala is a junior subjective synonym of Notodonta nodyna Swinhoe, 1907, Ann. Mag. nat. Hist. (7) 19: 206.

See also Macroshachia Matsumura, 1925.

ANIGRAEA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 139. NOCTUIDAE

Type-species: Anigraea rubida Walker, 1862, ibidem 6: 139,

by monotypy. Borneo: Sarawak.

Anigraea was established in the Notodontidae; it was placed in the Noctuidae by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 93.

ANIMOMYIA Dyar, 1908, Proc. ent. Soc. Wash. 10: 53. GEOMETRIDAE

Type-species: Animomyiu morta Dyar, 1908, ibidem 10: 53, by monotypy. [U.S.A.]: California, San Diego.

Animoniyia was established in the Liparidae, now Lymanand was transferred to the Geometridae by McDunnough, 1917, Ent. News 28: 233.

ANITA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 263 (key), 332.

Type-species: Anita basipuncta Schaus, 1901, ibidem 1901: 332, pl.12, fig.8, by original designation. VENEZUELA: Aroa.

ANKOVA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 227.

Type-species: Nioda lignea Butler, 1879, Ann. Mag. nat. Hist. (5) 4: 241, by original designation. MADAGASCAR: Fianarantsoa.

ANODONTA Herrich-Schäffer, [1854]. See Arhacia Herrich-Schäffer, 1855. NOTO

ANOTODONTA Strand, 1912, Arch. Naturgesch. 78 (A) 6: 164.

Type-species: Anotodonta bivittata Strand, 1912, ibidem 78 (A) 6: 164, by original designation. [EQUATORIAL GUINEA]: [Rio Muni], Benito region, Makomo.

ANSONIA Kiriakoff, 1967. See Oligoclona Felder, 1874.

ANTAEA Hübner, [1823] 1816, Verz. bekannter Schmett.: 266.

Type-species: Phalaena juturna Cramer, 1777, Uitlandsche Kapellen (Papillons exot.) 2: 48, pl.129, fig.E, by subsequent designation by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 342. SURINAM.

P. juturna is dated from the wrapper of part 11, not from the title-page of volume 2 which is dated 1779.

ANTARCTIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 191.

Type-species: *Phalaena metabus* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 117, pl.74, fig.D, by PRESENT DESIGNATION. SURINAM.

P. metabus is dated from the wrapper of the part and not

from the title-page of the volume, 1779.

Incorrect type-species designations: (1) Antarctia vulpina Hübner, [1825], a nominal species not originally included in Antarctia Hübner, [1820], and not linked with one of the originally included species when designated by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 225.

(2) Antarctia brunnea Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: [7], pl.70, figs 403, 404, designated by Hampson, 1900, Ann. S. Afr. Mus. 2: 60; and repeated by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 444; and by Gaede, 1926, in Seitz, Gross-Schmett. Erde 14: 106. A. brunnea was a nomen nudum at the time Antarctia Hübner, [1820], was established.

Antarctia has been used as the generic name for several species of Arctiidae, which are probably best placed in Palustra Bar, 1873. The use of Antarctia in the Arctiidae has arisen through the acceptance of the above incorrect type-species designations. Antarctia is here transferred from the Arctiidae to the Saturniidae, subfamily Hemileucinae, where it will become a junior subjective synonym of Hylesia Hübner, [1820], Verz. bekannter Schmett.: 186.

‡ANTARCTIS; Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 6: 1381.

An incorrect subsequent spelling of Antarctia Hübner, [1820].

‡ANTARCTIS; misspelling. See Antarctia Hübner, [1820].
SATURNIIDAE

ANTAXIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 475.

Type-species: *Amaxia pyrgion* Druce, 1897, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 2: 370, pl.74, fig.7, by original designation. Panama: Chiriqui Volcano.

ANTHELA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 853.

ANTHELIDAE

Type-species: Anthela ferruginosa Walker, 1855, ibidem 4: 854, by monotypy. AUSTRALIA: [New South Wales], Sydney.

Anthela was established in the Liparidae, now Lymantridae. Turner, 1904, Trans. ent. Soc. Lond. 1904: 479, placed it in the Lymantridae, subfamily Anthelinae, and subsequently raised Anthelinae to family level in 1920, ibidem 1919: 415.

ANTHELYMANTRIA Strand, 1925, in Seitz, Gross-Schmett. Erde 10: 363.

Type-species: Anthelymantria bistigmalis Strand, 1925, ibidem 10: 363, pl.57, row e, by original designation. [AUSTRALIA]: N.S.W., Richmond River, Cooktown.

The date of publication of the part of Seitz containing page 363 is stated on page 361 as xi.1925 in both the German and the English editions.

ANTHEUA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 687.

Type-species: Antheua simplex Walker, 1855, ibidem 3: 687, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 582. [SOUTH AFRICA]: [Natal], [Durban] Port Natal, Cape [Province].

Antheua was established in the Arctiidae; it was placed in the Notodontidae by Moore, [1883] 1882-3, Lepid. Ceylon 2: 111.

See also Sirenopyga Wallengren, 1858.

DIASTEMA Herrich-Schäffer, [1855] 1850-1858, Samml.

neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.66, fig.379.

Type-species: *Diastema straminea* Herrich-Schäffer, [1855] 1850-1858, *ibidem* 1(1): wrapper, pl.66, fig.379, by monotypy. [SOUTH AFRICA]: Natal.

A junior homonym of *Diastema* Guenée, 1852, in Boisduval & Guenée, *Hist. nat. Insectes* (Spec. gén. Lépid.) 6: 317, – Lepid., Noctuidae. There is no objective replacement name, but *D. straminea* is a junior subjective synonym of *Antheua simplex* Walker, 1855, the type-species of *Antheua* Walker, 1855; the latter is therefore available for use as a subjective replacement name.

‡ANTHEURA; Druce, 1888, Proc. zool. Soc. Lond. 1887: 685. An incorrect subsequent spelling of Antheua Walker, 1855.

DIASTEMA Aurivillius, 1904, Ark. Zool. 2 (4): 4.

Type-species: Antheua simplex Walker, 1855, by original designation.

A junior homonym of *Diastema* Guenée, 1852, *in* Boisduval & Guenée, *Hist. nat. Insectes* (Spec. gén. Lépid.) 6: 317, – Lepid., Noctuidae. The objective replacement name is *Antheua* Walker, 1855.

DIASTEMINA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 431.

Diastemina was established, unnecessarily, as an objective replacement name for Diastema Aurivillius, 1905.

ANTHEUELLA Kiriakoff, 1961, Bull. Annls Soc. r. ent. Belg. 97: 112.

Type-species: *Pydna flavida* Hampson, 1910, *Ann. Mag. nat. Hist.* (8) 5: 485, by original designation. [SOUTH AFRICA]: Natal, Durban.

Antheuella was established as a subgenus of Xanthodonta Gaede, 1928; it was raised to generic level by Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 92.

‡ANTHEURA; misspelling. See Antheua Walker, 1855.

ANTHEUSINA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 433.

Type-species: *Antheusina incompleta* Gaede, 1928, *ibidem* **14**: 433, pl.71 row i, by monotypy. EAST AFRICA.

The date of publication of the part of Seitz containing page 433 is cited on that page as 30.v.1928 in the German edition and as 26.xi.1928 in the English edition.

ANTHOMYZA Swainson, 1833. See Notophyson Boisduval, 1870.

ANTHORA Walker, 1855. See Laelia Stephens, 1828. LYMA

ANTHYPERAESCHRA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 642.

Type-species: *Hyperaeschra trichosticha* Hampson, 1897, *J. Bombay nat. Hist. Soc.* 11: 283, by original designation. [INDIA]: Sikkim.

The date of publication of the part of Seitz containing page 642 is cited on page 641 as 8.iv.1930 in both the German and the English editions.

ANTICHADISRA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 416.

Type-species: *Antichadisra dentata* Kiriakoff, 1974, *ibidem* 17: 417, text-fig.29, pl.5, fig.3, by original designation. SUMATRA: Dolok Merangir.

ANTICHERA Snellen, 1896, Tijdschr. Ent. 38: 180. AGAN Type-species: Neochera tortuosa Moore, 1872, Proc. zool. Soc. Lond. 1872: 570, pl.33, fig.2, by monotypy. INDIA.

ANTICHLORIS Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 9, 24.

Type-species: Zygaena eriphia Fabricius, [1777], Genera Insect.: 276, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 158. SURINAM.

See also *Illipula* Walker, 1854, and *Copaena* Herrich-Schäffer, [1855].

ERIPHIA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 96, 97; [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 16, 17.

Type-species: Antichloris phemonoe Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 9, by PRESENT DESIGNATION. BRAZIL.

The type designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 399 of Zygaena eriphia Fabricius is invalid as Hampson linked more than one of the originally included species with eriphia in the synonymy of the latter.

A junior homonym of *Eriphia* Latreille, 1817, *Nouv. Dictionnaire Hist. nat.* (Edn 2) **10**: 404, – Crustacea. There is no objective replacement name but *A. phemonoe* is a junior subjective synonym of *Zygaena eriphia* Fabricius, [1777], the typespecies of *Antichloris* Hübner, 1818. The latter is therefore available for use as a subjective replacement name.

Eriphia Herrich-Schäffer was placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1721.

ANTICLEAPA Kiriakoff, 1970, Ann. Transv. Mus. 26 (9): 192.

Type-species: Anticleapa varii Kiriakoff, 1970, ibidem 26 (9): 193, figs 12, 13, by original designation. South Africa: Natal, Impetyeni Forest.

ANTICOREURA Prout, 1918, Novit. zool. 25: 396 (key), 427.

Type-species: Coreura salmoni Druce, 1885, Proc. zool. Soc. Lond. 1885: 521, pl.32, fig.6, by original designation. Colombia: Antioquia, Frontino.

ANTICYRA Walker, 1855. See Dinara Walker, 1856. NOTO

ANTIFENTONIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 96.

Type-species: *Pseudofentonia cineraria* Kiriakoff, 1964, *Bonn. zool. Beitr.* 14: 277, abb.4, fig.34, pl.3, fig.36, by original designation (for *Pseudofentonia* Kiriakoff, 1964). China: Chekiang, west Tien-mu-Shan.

Antifentonia was established as the objective replacement name for *Pseudofentonia* Kiriakoff, 1964.

PSEUDOFENTONIA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 277.

Type-species: *Pseudofentonia cineraria* Kiriakoff, 1964, by original designation.

A junior homonym of *Pseudofentonia* Strand, 1912, *Fauna exot*. 2: 40, – Lepid., Notodontidae. The objective replacement name is *Antifentonia* Kiriakoff, 1967.

ANTIGODASA Kiriakoff, 1974, in Kiriakoff & Viette, Faune Madagascar 39: 6 (key), 94.

AGARISTIDAE

Type-species: Godasa rufodiscalis Rothschild, 1896, in Rothschild & Jordan, Novit. zool. 3: 47, pl.14, fig.23, by original designation. Holotype & MADAGASCAR, (BMNH).

Antigodasa in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

ANTILOBA Weymer, 1895, Stettin. ent. Ztg 55: 328. ARCT Type-species: Antiloba carnea Weymer, 1895, ibidem 55: 329, by monotypy. Brazil: Rio Grande do Sul.

A. carnea is a junior subjective synonym of Neritos repanda Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 711.

See also Neritos Walker, 1855.

ANTIMIMA Turner, 1917, Proc. R. Soc. Qd 29: 73. NOTO Type-species: Antimima cryptica Turner, 1917, ibidem 29: 73, by monotypy. AUSTRALIA: Western Australia, Quindalup.

ANTIOPHA Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 259 (key), 275.

Type-species: Antiopha multilinea Schaus, 1901, ibidem 1901: 275, pl.11, fig.3, by original designation. BRAZIL: Rio Janeiro, [Parana], Castro.

ANTIORA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1769.

Type-species: Antiora subfulva Walker, 1856, ibidem 7: 1769, by monotypy. BRAZIL.

Antiora was established in the Cossidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 266.

‡ANTIOZA; Dalla Torre, 1923, in Strand, Lepid. Cat. 29: 27. An incorrect subsequent spelling of Antiora Walker, 1856.

ANTIORA Möschler, 1883. See Pseudantiora Kirby, 1892.

ANTIOTRICHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.27. ARCT Type-species: Antiotricha vexata Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.27, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-

A. vexata is a junior subjective synonym of Mennis integra Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 181.

‡ANTIOZA; misspelling. See Antiora Walker, 1856. NOTO

ANTIPHA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 806.

Type-species: Antipha costalis Walker, 1855, ibidem 4: 806, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 456. [SRI LANKA] Ceylon.

Incorrect type-species designation: Lacida antica Walker, 1855, a nominal species not originally included in Antipha Walker, and not linked with one of the originally included names when cited by Moore, [1883] 1882-3, Lepid. Ceylon 2: 80.

ANTIPHALERA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 614.

Type-species: *Phalera bilineata* Hampson, 1896, *Fauna Br. India* (Moths) 4: 455, by monotypy. BHUTAN, INDIA: Khasia Hills.

The date of publication of the part of Seitz containing page 614 is cited on page 609 as 10.iii.1930 in both the German and English editions.

‡ANTIPHELEA; misspelling. See Antiphella Walker, 1856.

ANTIPHELLA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1743.

Type-species: *Liparis crocicollis* Herrich-Schäffer, [1854] 1850-1858, *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.24, fig.110, but included by Walker as '*Liparis crocicollis* Boisd.' an incorrect authorship, by monotypy. [SOUTH AFRICA]: [Cape of Good Hope] Cap.

‡ANTIPHELEA; Lucas, 1900, Arch. Naturgesch. 63 (2) 2: 489.

An incorrect subsequent spelling of Antiphella Walker, 1856.

ANTISCRANCIOLA Kiriakoff, 1967, Biologia gabon. 3: 342.

Type-species: Antiscranciola bisignata Kiriakoff, 1967, ibidem 3: 343, fig.29, by original designation. GABON: Bélinga.

ANTISTAURA Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 312.

Type-species: *Antistaura decorata* Kiriakoff, 1965, *ibidem* **101**: 312, figs 1, 2, by monotypy. ETHIOPIA: Diredawa, northwest of Harar.

ANTITHEMERASTIS Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 216.

Type-species: *Themerastis acrobela* Turner, 1922, *Proc. Linn. Soc. N.S.W.* 47: 383, by original designation. AUSTRALIA: Queensland, Cape York.

ANTITROTONOTUS Kiriakoff, 1966, Biologia gabon. 2: 132.

Type-species: *Antitrotonotus gabonensis* Kiriakoff, 1966, *ibidem* 2: 133, pl.A, fig.1, pl.1, fig.1, by original designation. GABON: Ivindo.

ANTIZANA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 437.

Type-species: *Antizana sparsata* Gaede, 1928, *ibidem* 14: 437, pl.72 row c, by monotypy. [ZIMBABWE-]RHODESIA.

The date of publication of the part of Seitz containing page 437 is cited on page 433 as 30.v.1928 in the German edition; and as 26.xi.1928 in the English edition.

ANTOCULEORA Ichinosé, 1973, Kontyu 41: 136.

Type-species: *Plusia ornatissima* Walker, 1858, *List Specimens lepid. Insects Colln Br. Mus.* 15: 1786, by original designation. Holotype 9, INDIA: North Hindostan, (BMNH).

Erythroplusia Ichinosé, 1962, was raised from subgeneric to generic rank, and on the same page Antoculeora was established as a subgenus of it.

Antoculeora in the Noctuidae, Plusiinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

ANTONA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 479.

Type-species: Antona subluna Walker, 1854, ibidem 2: 480, by monotypy. BRAZIL: Rio de Janeiro.

ANTONGILA Kiriakoff, 1958. See Exantongila Fletcher nom.

ANTOROKA Kiriakoff, 1969, Faune Madagascar 29: 152.

Type-species: Antoroka munda Kiriakoff, 1969, ibidem 29: 153, text-fig.100, pl.5, fig.55, by original designation. MADAGASCAR: eastern, Maroantsetra district, Farankaraina forest station.

ANTSALOVA Kiriakoff, 1960, Revue fr. Ent. 27: 174. NOTO Type-species: Antsalova musculus Kiriakoff, 1960, ibidem 27: 175, fig.2, by original designation. MADAGASCAR: western, Antsalova district, Andobo.

ANUROCAMPA Herrich-Schäffer, [1854] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.18, figs 83, 84.

Type-species: Anurocampa mingens Herrich-Schäffer, [1854] 1850-1858, ibidem 1(1): wrapper, pl.18, figs 83, 84, (but attributed to Boisduval, an incorrect authorship), by monotypy. Brazil.

‡ANEUROCAMPA; Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1047.

An incorrect subsequent spelling of *Anurocampa* Herrich-Schäffer, [1854].

ANYCLES Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 253.

Type-species: *Euchromia anthracina* Walker, 1854, *ibidem* 1: 253, by subsequent designation by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 398. VENEZUELA, [BRAZIL]: [Tefé] Ega.

The name Anycles probably originated from Boisduval but was published and made available by Walker. The name Anycles probably had the same origin and was made available by Herrich-Schäffer.

Anycles was established as a subgenus of Euchromia Hübner, [1819], in the 'Zygaenides'; it was placed in the Arctiidae as a valid genus by Butler, 1876, J. Linn. Soc. (Zool.) 12: 425; and was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 398.

AMYCLES Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.49, fig.259.

Type-species: Euchromia anthracina Walker, 1854, by monotypy.

A junior objective synonym of Anycles Walker, 1854.

‡AMYCLES; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1633.

An incorrect subsequent spelling of *Anycles* Walker, 1854, as Walker was referring back to his earlier usage, not to *Amycles* Herrich-Schäffer, [1855].

ANZABE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1093.

Type-species: Anzabe sinensis Walker, 1855, ibidem 5: 1093, by monotypy. Hong Kong.

Anzabe was established in the Notodontidae; it was placed in the Limacodidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 531.

AOBA Kiriakoff, 1965, Bull. Inst. r. Sci. nat. Belg. 41 (8): 13.
NOTO

Type-species: *Aoba tosta* Kiriakoff, 1965, *ibidem* 41 (8): 15, fig.4, by original designation. [ZAIRE]: Kivu, Kabunga.

AOSHACHIA Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19: 9. GEOMETRIDAE

Type-species: Aoshachia virescens Matsumura, 1927, ibidem 19: 9, pl.5, fig.37, by original designation. Talwan: Baibara. Aoshachia was established in the Notodontidae; it was transferred to the Geometridae by Inoue, 1955, Kontyu 23: 71.

APAIDIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xviii, 84, 467.

Type-species: Callimorpha mesogona Godart, [1824] 1822, Hist. nat. Lépid. Papillons Fr. 4: 396, pl.40, fig.6, by original designation. France: Paris region.

Apaidia was established in order to name the concept *Paida* Hübner sensu Herrich-Schäffer, [1847] 1845, *Syst. Bearbeitung Schmett. Eur.* 2: 156.

APANTESIS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 631.

Type-species: *Apantesis radians* Walker, 1855, *ibidem* 3: 632, by monotypy. [U.S.A.]: Georgia.

APELA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1092.

Type-species: *Apela divisa* Walker, 1855, *ibidem* 5: 1093, by monotypy. Type-locality: stated to be Nepal, but this was a mistake, for the species occurs in Trinidad and the northern part of Tropical America.

‡APELLA; Rothschild, 1917, Novit. zool. 24: 262.
An incorrect subsequent spelling of Apela Walker, 1855.

‡APELLA; misspelling. See Apela Walker, 1855. NOTO

APEPLOPODA Watson nom. n. for Gymnopoda Felder, 1874.

Type-species: *Gymnopoda ochracea* Felder, 1874, *in* Felder & Rogenhofer, *Reise öst. Fregatte Novara* (Zool.) 2 (Abt.2): pl.102, fig.22, by subsequent designation (for *Gymnopoda* Felder, 1874) by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 137. [Colombia]: Bogota, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 5.

GYMNOPODA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, figs 22, 23.

Type-species: Gymnopoda ochracea Felder, 1874, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 137.

A junior homonym of *Gymnopoda* Macquart, 1835, *Hist.* nat. Insectes Diptères 2: 502, – Insecta, Diptera. The objective replacement name is Apeplopoda Watson nom. n.

APHISAON Boisduval, 1870, Considérations Lépid. Guatemala: 86.

Type-species: *Phalaena tricolor* Sulzer, 1776, *Geschichte Insecten*: 160, fig., by original designation (but attributed to Cramer, an incorrect authorship). Type-Locality: not stated, [AMERICA]: [tropical].

See also Thebrone Boisduval, 1870.

APHOMOEOMA Collenette, 1959, Annls Soc. ent. Fr. 128: LYMA

Type-species: *Aphomoeoma mesembrinum* Collenette, 1959, *ibidem* **128**: 118, pl.2, fig.5, by original designation. MADAGASCAR: S., Bekily.

APHRA Watson nom. n. for Phara Walker, 1854. CTEN Type-species: Dioptis trivittata Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 322, by monotypy (of Phara Walker, 1854). Type-locality: not known, [Brazil].

PHARA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 321.

Type-species: *Dioptis trivittata* Walker, 1854, by monotypy. A junior homonym of *Phara* Walker, 1850, *in* Saunders, *Insecta Saundersiana* 1: 9, – Insecta, Diptera. The objective replacement name is *Aphra* Watson nom. n.

OLINA Walker, 1869, Characters undescr. Lepid. Heterocera:

Type-species: Olina nyctemeroides Walker, 1869, ibidem: 6, by monotypy. Type-locality: not stated, [AMERICA]: [tropical].

A junior homonym of Olina Robineau-Desvoidy, 1830, Mém. prés. div. Sav. Acad. Sci. Inst. Fr. 2: 812, — Insecta, Diptera. There is no objective replacement name but O. nyctemeroides is congeneric with Dioptis trivittata the type-species of Aphra Watson nom. n. The latter is therefore available for use as a subjective replacement name.

APHYARCTIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 50.

Type-species: *Automolis surinamensis* Rothschild, 1912, *Novit. zool.* 18: 158, by original designation. Surinam: Maroewym Valley, Aroewarwa Creek.

APHYLE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 720.

Type-species: *Aphyle margaritaceus* Walker, 1855, *ibidem* 3: 720, by monotypy. [Brazil]: Para.

APICONOMA Butler, 1876, J. Linn. Soc. (Zool.) 12: 422.

Type-species: *Euchromia opposita* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 260, by original designation. BRAZIL.

APINA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 587 (key), 756.

AGARISTIDAE

Type-species: *Apina callisto* Walker, 1855, *ibidem* 3: 756, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 442.

Apina was established in the Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, *ibidem* 1: 442; it was transferred to the Noctuidae by Swinhoe, 1903, *Trans. ent. Soc. Lond.* 1903: 497; and was placed in the Agaristidae by Jordan, 1912, *in* Seitz, *Gross-Schmett. Erde* 11: 26.

APISA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 916.

THYRETIDAE

Type-species: *Apisa canescens* Walker, 1855, *ibidem* **4**: 917, by monotypy. South Africa.

Apisa was established in the Liparidae, now Lymantriidae; it was placed in the Arctiidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 220; and was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 49.

APISTAESCHRA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 272.

Type-species: *Apistaeschra substyxana* Kiriakoff, 1964, *ibidem* 14: 272, abb.3, fig.29, pl.3, fig.30, by original designation. China: Kiangsu, Lungtan near Nanking.

‡APISTASIA; misspelling. See Apistosia Hübner, 1818.

APISTOSIA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 13.

Type-species: *Apistosia judas* Hübner, 1818, *ibidem* 1: 13, pl.[9], figs 51, 52, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 286. BRAZIL.

‡HIPOCRITA Hübner, 1808, Erste Zuträge Samml. exot. Schmett.: 4, 5.

Included in a work rejected for nomenclatural purposes by Int. Commn zool. Nom., 1966, Bull. zool. Nom. 23 (Opinion 789): 213. Placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1858.

Two included species: ‡Hipocrita judas Hübner, 1808, ibidem: 4, subsequently published and made available as Apistosia judas Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 13, pl.[9], figs 51, 52, — Arctiidae, Lithosiinae; and ‡Hipocrita tendinosa Hübner, 1808, ibidem: 5, subsequently published and made available as Phaeochlaena tendinosa Hübner, 1818, ibidem 1: 18, pl.[16], figs 89, 90, — Dioptidae.

‡APISTASIA; Hübner, [1826] 1816, Verz. bekannter Schmett. (Anzeiger): 27.

An incorrect subsequent spelling of Apistosia Hübner, 1818.

APISTUS Agassiz, 1847, Nomencl. zool. (Index univl.): 29. An unjustified emendation of Apistosia Hübner, 1818.

A junior homonym of *Apistus* Cuvier, 1829, *in* Cuvier & Valenciennes, *Hist. nat. Poissons* 4: 391, – Pisces:

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Apistus* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

APISTUS Agassiz, 1847. See Apistosia Hübner, 1818. LITH

APOCEREA Dyar, 1905, Proc. U.S. natn. Mus. 29: 174.

Type-species: *Apocerea sobria* Schaus, 1905, *ibidem* **29**: 193, by original designation (in anticipation of Schaus's description of *sobria*). French Guiana: Maroni River, St Jean.

Apocerea was available but was without an available included

species between the date of its publication 1905 October 3rd and that of its type-species 1905 October 11th by Schaus.

APOCERURA Lattin, Becker & Bender, 1974, Ent. Z., Frankf.a.M. 84: 206.

Type-species: *Phalaena erminea* Esper, 1783, *Die Schmett*. 3: 100, pl.19, figs 1, 2, by original designation. GERMANY: Frankfurt am Main.

Apocerura was established as a subgenus of Cerura Schrank, 1802.

P. erminea is dated from the wrapper of Heft 18, not from the title-page of the volume which is dated 1782.

APOCINESIA Bryk, 1930. See Scedros Walker, 1854. DIOP

APOCRISIAS Franclemont, 1966, Proc. ent. Soc. Wash. 68: 255.

Type-species: *Apocrisias thaumasta* Franclemont, 1966, *ibidem* **68**: 256, fig., by original designation. U.S.A.: Arizona, Santa Cruz County, Santa Rita Mts, Madera Canyon.

APOGUREA Watson nom. n. for Pareugoa Daniel, 1951.

Type-species: *Pareugoa grisescens* Daniel, 1951, *Bonn. zool. Beitr.* **2**: 299, pl.1, fig.6, by original designation (for *Pareugoa* Daniel, 1951). China: Chekiang, Hangchow.

PAREUGOA Daniel, 1951, Bonn. zool. Beitr. 2: 299.

Type-species: *Pareugoa grisescens* Daniel, 1951, by original designation.

designation.

A junior homonym of *Pareugoa* Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* 2: 405, – Lepid., Arctiidae. The objective replacement name is *Apogurea* Watson nom. n.

‡APOSURA Costa, [1836] 1832-1836, Fauna Regno Napoli (Lepid.): [148].

‡Aposura is not nomenclaturally available as a genus-group name as it was used by Costa in a suprageneric sense.

APOTHOSIA Hampson, 1918, Novin. zool. 25: 100. LITH Type-species: Apothosia conformis Hampson, 1918, ibidem 25: 100, by original designation. [MALAWI] Br. C. Afr.: lower Shire [River] valley.

APROSDOCETOS Kiriakoff, 1962, Revue Zool. Bot. afr. 66:
15. NOTO

Type-species: *Phalera inexpectata* Rothschild, 1917, *Novit. zool.* **24**: 253, pl.7, fig.10, by original designation. NIGERIA: southern, Ilesha.

APTEROGYNIS Guenée, 1867. See Orgyia Ochsenheimer, 1810.

APTILOSIA Schaus, 1911, Ann. Mag. nat. Hist. (8) 7: 357.

Type-species: Aptilosia crocea Schaus, 1911, ibidem (8) 7: 358, by monotypy. Costa Rica: Juan Viñas; Tuis; and Guapiles.

APYRE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 490.

Type-species: Apyre separata Walker, 1854, ibidem 2: 491, by monotypy. [BRAZIL]: Amazon, [Tefé] Ega.

ARACHNIS Geyer, 1837, in Hübner, Zuträge Samml. exot. Schmett. 5: 28.

Type-species: Arachnis aulaea Geyer, 1837, ibidem 5: 28, pl.[157], figs 913, 914, by monotypy. MEXICO.

ARAEOMOLIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 2, 38.

Type-species: Araeomolis rhodographa Hampson, 1901, ibi-

dem 3: 38, fig.31, by original designation. Colombia: Amazon[a]s, Rio Negro, [Tomar] Thomar.

‡AREOMOLIS; Rothschild, 1909, Novit. zool. 16: 269.

An incorrect subsequent spelling of Araeomolis Hampson, 1901.

‡ARAPHA; misspelling. See Anaphe Walker, 1855. THAU

ARARA Walker, 1855. See Lepidokirbyia Travassos, 1949.

ARATTATHA Janse, 1917, Ann. Durban Mus. 1: 476.
NOCTUIDAE

Type-species: *Arattatha superba* Janse, 1917, *ibidem* 1: 476, by monotypy. Holotype & South Africa: Barberton.

Arattatha in the Noctuidae, Catocalinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

ARBORICORNIS Hampson, 1908. See Arboricornus Hampson, 1894.

ARBORICORNUS Hampson, 1894, Fauna Br. India (Moths) 2: 164, 258.

Type-species: *Arboricornus ruber* Hampson, 1894, *ibidem* 2: 258, fig.146, by original designation.

Further information in Nyc, 1975, Generic Names Moths World 1: 56.

ARBORICORNIS Hampson, 1908, Cat. Lepid. Phalaenae Br. Mus. 7: 358.

An unjustified emendation of *Arboricornus* Hampson, 1894. *Arboricornis* in the Noctuidae, Acronictinae is cited here as it was not included in Nye, 1975, *Generic Names Moths World* 1.

ARCHIGARGETTA Kiriakoff, 1967, Tijdschr. Ent. 110: 38.

Type-species: Archigargetta cyclopea Kiriakoff, 1967, ibidem 110: 38, fig.2, by original designation. New Guinea: [Irian Jaya], Waigeo [Island], Camp Nok.

ARCHILEMA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 72.

Type-species: *Lepista subumbrata* Holland, 1893, *Psyche, Camb.* 6: 411, by original designation. [GABON]: valley of the [Ogooué] Ogové River.

ARCHINADATA Kiriakoff, 1969 January, Bull. Inst. fond. Afr. noire (A) 30 (4): 1454. NOTO

Type-species: *Macronadata aurivilliusi* Kiriakoff, 1954, *Bull. Inst. r. Sci. nat. Belg.* **30** (14): 5, fig.4, by original designation. [ZAIRE] Congo Belge: Kinshasa.

Archinadata is dated from the foot of page 1762 of part 4, not from its title-page which is dated October 1968.

ARCHISTILBIA Kiriakoff, 1954, Revue Zool. Bot. afr. 49: 317.

Type-species: *Archistilbia cineracea* Kiriakoff, 1954, *ibideni* 49: 317, pl.1, fig.13, pl.2, fig.17, by original designation. [ZAIRE] Congo Belge: Katanga, [Lubumbashi] Elisabethville.

ARCHITESMIA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 28.

Type-species: Archithosia sordida Birket-Smith, 1965, ibidem 1: 29, figs 18, 19, by original designation. CAMEROUN: Ja River, Bitje.

Architesmia was established as a subgenus of Archithosia Birket-Smith, 1965.

ARCHITHOSIA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 11.

Type-species: *Lithosia costimacula* Mabille, 1879, *Bull. Soc. zool. Fr.* 3: 87, by original designation. CONGO.

ARCIERA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 20.

Type-species: Arciera roseiventris Kiriakoff, 1962, ibidem 114: 20, fig.21, by original designation. [ZAIRE]: Sankaru, Katako-Kombe.

ARCILASISA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 470.

Type-species: Arcilasisa sobria Walker, 1865, ibidem 32:

470, by monotypy. [INDIA]: South Hindostan.

Arcilasisa was established in the Notodontidae; it was placed in the Noctuidae by Cotes & Swinhoe, 1888, Cat. Moths India (3): 267.

ARCTAGYRTA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: x, 4, 173.

Type-species: Agyrta nana Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1652, by original designation. [Brazil]: Amazon valley.

ARCTATA Roepke, 1949, Trans. R. ent. Soc. Lond. 100: 50.

Type-species: Nyctemera arctata Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1664, by monotypy. [INDIA]: Hindostan; [Assam], [Cherrapunji] Cherra Poonjee. Arctata was established as a subgenus of Nyctemera Hübner, [1820].

ARCTIA Schrank, 1802, Fauna Boica 2 (2): 152. ARCT Type-species: Phalaena caja Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 500, by subsequent designation by Westwood, 1840, Introd. mod. Classif. Insects 2 (Synopsis Genera Br. Insects): 92. Type-locality: not stated, [Europe].

Neave, 1939, Nomencl. zool. 1: 283, also cited 'Arctia Huebner [non Schrank] [1820], Verz. bekannt. Schmett., (12): 183'. This was an error, as Hübner was referring to Arctia Schrank, 1802.

‡HYPERCOMPE Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) Opinion 97: 19; also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140.

Only included species: Phalaena caja Linnaeus, 1758.

EYPREPIA Ochsenheimer, 1810, Schmett. Eur. 3: 299.

Type-species: *Phalaena caja* Linnaeus, 1758, by subsequent designation by Curtis, 1824, *Br. Ent.* 1: 21 (but included by Ochsenheimer as *caia*, an unjustified emendation by Ochsenheimer, 1810).

A junior objective synonym of Arctia Schrank, 1802.

‡EUPREPIA; Hübner, [1819] 1816, Verz. bekannter Schmett.: 181.

An incorrect subsequent spelling of *Eyprepia* Ochsenheimer, 1810.

ZOOTE Hübner, [1820] 1816, Verz. bekannter Schmett.: 181.
Type-species: Phalaena caja Linnaeus, 1758, by subsequent designation by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus.
3: 463 (but cited as caia, an unjustified emendation).

A junior objective synonym of Arctia Schrank, 1802.

CHELONIA Godart, [1823] 1822, Hist. nat. Lépid. Papillons Fr. 4: 299.

Type-species: *Phalaena caja* Linnaeus, 1758, by subsequent designation by Duponchel, 1843, *Dict. univl Hist. nat.* 3: 460.

A junior homonym of Chelonia Brongniart, 1800, Bull. Soc.

philomath. Paris 2: 89, – Reptilia. The objective replacement name is Arctia Schrank, 1802.

‡ARCTINIA; Eichwald, 1830, Zool. spec. 2: 195.

An incorrect subsequent spelling of Arctia Schrank, 1802.

‡ARETIA; Poey, 1832, Centurie Lépid. Cuba: pl.[17].

An incorrect subsequent spelling of Arctia Schrank, 1802.

EUPREPIA Agassiz, 1846, Nomencl. zool. (Nomina syst. Generum Lepid.): 29.

An unjustified emendation of Eyprepia Ochsenheimer, 1810.

CALLARCTIA Packard, 1864, Proc. ent. Soc. Philad. 3: 114. Type-species: Phalaena caja Linnaeus, 1758, by subsequent designation (for Chelonia Godart, [1823]) by Duponchel, 1843, Dict. univl Hist. nat. 3: 460.

A junior objective synonym of Arctia Schrank, 1802.

Callarctia was proposed as the objective replacement name for Chelonia Godart, [1823].

ARCTIARPIA Travassos, 1951, Revta bras. Biol. 11: 249.

Type-species: *Idalus melanopasta* Dognin, 1907, *Annls Soc. ent. Belg.* **51**: 226, by original designation. FRENCH GUIANA: St Laurent du Maroni.

ARCTIITES Rebel, 1898, Sber. K. Akad. Wiss. Wien 107 (Abt.1): 732.

Type-species: *Arctiites deletus* Rebel, 1898, *ibidem* **107** (Abt.1): 732, pl.1, fig.6, by monotypy. ITALY: Gabbro near Pisa. Miocene.

‡ARCTINIA; misspelling. See Arctia Schrank, 1802. ARCT

ARCTIOMORPHA Herrich-Schäffer, 1855 December 31, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.75, fig.434.

Type-species: *Arctiomorpha euprepiaeformis* Herrich-Schäffer, 1855 December 31, *ibidem* 1 (1): wrapper, pl.75, fig.434, by monotypy. [SOUTH AFRICA]: Natal.

A. euprepiaeformis is a junior subjective synonym of Anaphe reticulata Walker, 1855 [August 11], List Specimens lepid. Insects Colln Br. Mus. 4: 856.

Arctiomorpha and A. euprepiaeformis are both dated from the wrapper of the part, not from the title-page of the volume which is dated 1850-1858.

Arctionorpha was also included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' published in 1855 December [31], Syst. Bearbeitung Schmett. Eur. 6: 91, and again in [1856 September], Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 11. In both these text references the figure of the type-species is listed in error as '438'.

See also Anaphe Walker, 1855 [August 11].

ARCTIONEURA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.107, fig.16. ARCT

Type-species: Arctioneura lorquinii Felder, 1874, ibidem 2 (Abt.2): pl.107, fig.16, by monotypy. [SULAWESI] Celebes: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 2.

A. lorquinii Felder is a junior subjective synonym of Satara aequata Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 160.

See also Satara Walker, [1865].

ARCTOPROCTUS Herrich-Schäffer, 1855. See Trichiocercus Stephens, 1835. THAU

ARCTORNIS Germar, 1810, Diss. sistens Bombycum Species:

Type-species: *Bombyx v-nigrum* Fabricius, 1775, *Syst. Ent.*: 577, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 432, but cited as *Phalaena l-nigrum*

Müller, 1764, a nominal species not originally included in Arctornis. Kirby, however, placed l-nigrum as the senior synonym of v-nigrum, a nominal species originally included in Arctornis. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. [D.D.R.]: Saxony.

B. v-nigrum is a junior subjective synonym of Phalaena (Bombyx) l-nigrum Müller, 1764, Fauna Insect. Fridrichsdal-

ina . . .: 40.

ARCTURUS Curtis, 1830. See Trichiocercus Stephens, 1835.

ARDICES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 709.

Type-species: Ardices fulvohirta Walker, 1855, ibidem 3: 710, by monotypy. [AUSTRALIA]: [Tasmania] Van Dieman's

‡ARDONCA; misspelling. See Ardonea Walker, 1854. LITH

ARDONEA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 482.

Type-species: Ardonea morio Walker, 1854, ibidem 2: 482, by monotypy. VENEZUELA.

See also Doracis Boisduval, 1870.

‡ARDONCA; Dognin, 1894, Annls Soc. ent. Belg. 38: 238. An incorrect subsequent spelling of Ardonea Walker, 1854.

ARDONIPSA Dyar, 1912, Proc. U.S. natn. Mus. 42: 50.

Type-species: Ardonipsa melas Dyar, 1912, ibidem 42: 50,

by original designation. MEXICO: Cuernavaca.

A. melas is a junior subjective synonym of Ctenucha dolens Druce, 1897, Biologia cent.-am. (Zool.) Lepid.-Heterocera 2: 360, pl.73, fig.4.

See also Stenucha Hampson, 1901, and Zamolis Dyar, 1910.

ARDONISSA Dognin, 1907, Annls Soc. ent. Belg. 51: 226.

Type-species: Ardonissa adscitina Dognin, 1907, ibidem 51: 226, by original designation. PERU: Cuzco.

ARE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 758.

Type-species: Phalaena marginata Drury, 1773, Illust. nat. Hist. exot. Insects 2: index, 38, fig., by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 186. JAMAICA.

AREAS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 658.

Type-species: Areas orientalis Walker, 1855, ibidem 3: 658, by monotypy. INDIA: northern.

‡AREOMOLIS; misspelling. See Araeomolis Hampson, 1901. ARCT

ARESIA Barnes & McDunnough, 1913, Contr. nat. Hist. Lepid. N. Am. 2: 167.

Type-species: Aresia parva Barnes & McDunnough, 1913, ibidem 2: 167, pl.1, fig.1, by original designation. U.S.A.: Florida, Everglade[s]; and Texas, Brownsville.

A. parva is a junior subjective synonym of Afrida ydatodes Dyar, 1913, Insecutor Inscit. menstr. 1: 31.

ARESTHA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key), 805.

Type-species: Arestha antica Walker, 1855, ibidem 4: 805, by monotypy. JAVA.

A. antica is a junior subjective synonym of Bombyx pennatula Fabricius, 1793, Ent. Syst. 3 (1): 465.

‡ARETIA; misspelling. See Arctia Schrank, 1802.

ARETYPA Smith, 1903, Trans. Am. ent. Soc. 29: 222. NOCTUIDAE.

Type-species: Eucalyptera pectinicornis Smith, 1899, Jl N.Y. ent. Soc. 7: 232, by original designation. Syntypes 28, 39, [U.S.A.]: Arizona, Phoenix.

E. pectinicornis is a junior subjective synonym of Pseudorgyia

russula Grote, 1883, Papilio 3: 75.

In Nye, 1975, Generic Names Moths World 1: 58, the typespecies was incorrectly cited as Feltia pectinicornis Smith, 1890, a different species.

‡ARETYPE; Hampson, 1926, Descr. new Genera Species Lepid. Phalaenae Subfamily Noctuinae Br. Mus.: 243.

An incorrect subsequent spelling of Aretypa Smith, 1903. ‡Aretype is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡ARETYPE; misspelling. See Aretypa Smith, 1903. NOCTUIDAE

AREVA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 525.

Type-species: Areva laticilia Walker, 1854, ibidem 2: 526, by monotypy. VENEZUELA.

ARGILA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 810.

Type-species: Argila basalis Walker, 1855, ibidem 4: 811, by monotypy. SIERRA LEONE.

ARGINA Hübner, [1819] 1816, Verz. bekannter Schmett.: 167.

Type-species: Phalaena cribraria Clerck, 1764, Icon. Insect. rariorum 2: pl. [54], fig. 4 & register (a junior primary homonym, see below; attributed by Hübner to Cramer, an incorrect authorship), by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 350. INDIA.

Phalaena cribraria Clerck, 1764, is a junior primary homonym of Phalaena cribraria Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 507, - Arctiidae, Arctiinae. There is no objective replacement name but Phalaena astrea Drury, 1773, Illust. nat. Hist. exot. Insects 2: index, 11, pl.6, fig.3, is the oldest of its junior subjective synonyms and is therefore available for use as a subjective replacement name.

LOMASPILIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.100, fig.27.

Type-species: Lomaspilis pantheraria Felder, 1874, ibidem 2 (Abt.2): pl.100, fig.27, by monotypy. [South Africa]: 'Caffraria', cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

A junior homonym of *Lomaspilis* Hübner, [1825] 1816, Verz. bekannter Schmett.: 306, - Lepid., Geometridae. There is no objective replacement name but L. pantheraria is a junior subjective synonym of *Phalaena cribraria* Clerck, 1764, the type-species of Argina Hübner, [1819]. The latter is therefore available for use as a subjective replacement name.

ARGYROEIDES Butler, 1876, J. Linn. Soc. (Zool.) 12: 403.

Type-species: Glaucopis ophion Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 191, by original designation. VENEZUELA, BRAZIL: 'Sta Carthagena'.

ARGYROSTAGMA Aurivillius, 1904, Ark. Zool. 2 (4): 56.

Type-species: Porthetria niobe Weymer, 1896, Berl. ent. Z.

41: 89, by original designation. [Tanzania]: [Tanganyika], [Kilimanjaro] Kilimandscharo.

ARGYROTHYRIS Bryk, 1913, Societas ent. 28: 101. NOTO Type-species: Argyrothyris allocota Bryk, 1913, ibidem 28: 101, by monotypy. CAMEROUN: Lolodorf.

ARHABDOSIA Dyar, 1905, Proc. U.S. natn. Mus. 29: 174.

Type-species: *Arhabdosia subvarda* Schaus, 1905, *ibidem* **29**: 201, by original designation (in anticipation of Schaus's description of *subvarda*). French Guiana: St Jean.

Arhabdosia was available but was without an available included species between its date of publication 1905 October 3rd and that of its type-species 1905 October 11th by Schaus.

ARHACIA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 91.

Type-species: Anodonta combusta Herrich-Schäffer, [1854] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.18, figs 81, 82 (but attributed to Boisduval, an incorrect authorship), by monotypy. Brazil.

Arhacia was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

ANODONTA Herrich-Schäffer, [1854] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.18, figs 81, 82.

Type-species: Anodonta combusta Herrich-Schäffer, [1854],

by monotypy.

A junior homonym of Anodonta Lamarck, 1799, Mém. Soc. Hist. nat. Paris: 87, – Mollusca; placed (as a junior homonym) on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.961, in 1957, Opin. Decl. int. Commn zool. Nom. 1 (E) Direction 72: 168, 186. The objective replacement name is Arhacia Herrich-Schäffer, 1855.

‡ARRHACIA; Bergroth, 1908, Zoologist (4) 12: 252.

An incorrect subsequent spelling of *Arhacia* Herrich-Schäffer, 1855.

ARINA Walker, 1856. See Polypoetes Druce, 1885. DIOP

ARISTODAEMA Wallengren, 1858, Ofvers. K. VetenskAkad. Förh. Stockh. 15: 136.

Type-species: Aristodaema arauna Wallengren, 1858, ibidem 15: 136, by original designation. BRAZIL.

A. arauna is a junior subjective synonym of Laemocharis hanga Herrich-Schäffer, [1854] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.[47], fig.246

Aristodaema was redescribed as new by Wallengren, 1861, K. svenska Fregatten Eugenies Resa omkring Jorden 2 (1): 359.

ARMIANA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 141.

Type-species: *Armiana lativitta* Walker, 1862, *ibidem* 6: 142, by monotypy. Borneo: Sarawak.

A lectotype [male] of Armiana lativitta was designated and figured by Swinhoe, 1892, Cat. east. and Aust. Lepid. Heter-

ocera Colln Oxf. Univ. Mus. 1: 296, pl.8, fig.1. Swinhoe's designation was accompanied by an incorrect original reference, repeated by subsequent authors, relating to a species of Noctuidae described from a female less than half the size of the specimen that was figured.

ARNA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1159 (key), 1176.

Type-species: *Arna apicalis* Walker, 1855, *ibidem* 5: 1177, by monotypy. [SRI LANKA] Ceylon.

Arna was established in the Drepanulidae, now Drepanidae; it was placed in the Liparidae, now Lymantriidae, by Moore, [1883] 1882-3, Lepid. Ceylon 2: 84.

ARNOPHILA Mabille, 1880, Annls Soc. ent. Fr. (5) 9: 309 (but attributed to Guérin-Méneville, an incorrect authorship).

Type-species: Agarista stellata Guérin-Méneville, 1829-1838 [1844], Icon. Règne anim. Cuvier 3 (Insectes): 493, by monotypy (but cited as ‡stella, an incorrect subsequent spelling). MADAGASCAR.

See also Anaphela Walker, 1855.

AROA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key), 791.

Type-species: *Aroa discalis* Walker, 1855, *ibidem* 4: 792, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 463. SOUTH AFRICA: Cape, and [Durban] Port Natal

See also Ornithopsyche Wallengren, 1863.

‡ARVA; Grünberg, 1913, Arch. Naturgesch. 78 (B) 7: 110. An incorrect subsequent spelling of Aroa Walker, 1855.

ARPEMA Schaus, 1915, Trans. Am. ent. Soc. 41: 5. NOTO Type-species: Arpema megalopia Schaus, 1915, ibidem 41: 5, by original designation. BRAZIL: south-eastern, Joinville.

‡ARPYIA; misspelling. See Harpyia Ochsenheimer, 1810.

‡ARRHACIA; misspelling. See Arhacia Herrich-Schäffer, 1855.

ARRHYTHMICA Turner, 1940, Proc. R. Soc. Qd 51: 72.

Type-species: Arrhythmica semifusca Turner, 1940, ibidem 51: 73, by monotypy. Australia: Western Australia, Denmark.

ARTANASA Kiriakoff, 1960, Revue fr. Ent. 27: 175. NOTO Type-species: Artanasa viettei Kiriakoff, 1960, ibidem 27: 176, fig.3, by original designation. MADAGASCAR: south-western, Sakaraha district, Lambomakandro forest.

ARTAXA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key), 794.

Type-species: Artaxa guttata Walker, 1855, ibidem 4: 795, by subsequent designation by Moore, [1883] 1882-3, Lepid. Ceylon 2: 84, but cited as Bombyx digramma Guérin-Méneville, [1844], a nominal species not originally included in Artaxa. Moore on the next page placed digramma as the senior synonym of guttata, a nominal species originally included in Artaxa. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. A. guttata is now regarded as a valid species. [Bangladesh]: [Sylhet] Silhet, India: North, [Nepal] Nepaul, EAST INDIES.

ARTIMELIA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 248. ARCT

Type-species: Chelonia latreillii Godart, [1823] 1822, Hist. nat. Lépid. Papillons Fr. 4: 318, by monotypy. Spain.

Artimelia was established as a subgenus of Pachylischia Rambur, 1866.

ARUNDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 466.

Type-species: *Arunda opponens* Walker, 1865, *ibidem* 32: 467, by monotypy. [INDIA]: South Hindostan.

ARUTA Swinhoe, 1922, Ann. Mag. nat. Hist. (9) 10: 471.

Type-species: Cispia flavipes Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 493, by monotypy. INDIA: [Sikkim] Sikhim.

‡ARVA; misspelling. See Aroa Walker, 1855.

ASCAPTESYLE Dyar, 1905, Proc. U.S. natn. Mus. 29: 175.

Type-species: Ascaptesyle submarginata Schaus, 1905, ibidem 29: 201, by original designation (in anticipation of Schaus's description of submarginata). TRINIDAD.

Ascaptesyle was available but was without an available included species between its date of publication 1905 October 3rd and that of its type-species 1905 October 11th by Schaus.

ASINUSCA Wallengren, 1862, Ofvers. K. Vetensk Akad. Förh. Stockh. 19: 197, 199. Available but without included species until Wallengren, 1863, Wien ent. Monatschr. 7: 139 (but cited as ‡Asinutea, an incorrect subsequent spelling).

Type-species: Asinusca atricornis Wallengren, 1863, ibidem 7: 139, by subsequent monotypy (but cited as ‡Asinutea, an incorrect subsequent spelling). South Africa: [eastern] Caffragia orientali

A. atricornis (as a nomen nudum) was mentioned by Wallengren, 1862, ibidem 19: 199, but was first made available by Wallengren, 1863, ibidem 7: 139. It was redescribed, as new, by Wallengren, 1865, K. svenska VetenskAkad. Handl. 5 (4): 13.

Asinusca was redescribed, as new, by Wallengren, 1865, ibidem 5 (4): 13.

‡ASINUTEA; Wallengren, 1863, Wien. ent. Monatschr. 7: 139. An incorrect subsequent spelling on the evidence of the material of the type-species of Asinusca which is labelled Asinusca atricornis (Dr P.I. Persson, in litt.) and that the spelling Asinusca was used again by Wallengren in 1865, K. svenska VetenskAkad. Handl. 5 (4): 13.

‡ASINUTEA; misspelling. See Asinusca Wallengren, 1862.

ASOTA Hübner, [1819] 1816, Verz. bekannter Schmett.: 164.

Type-species: *Phalaena javana* Cramer, 1780, *Uitlandsche Kapellen (Papillons exot.)* 3: 146, 175, pl.274, fig.C, by subsequent designation by Jordan, 1896, *Novit. zool.* 3: 203. JAVA: Samarang.

P. javana is dated from the wrapper of the part and not from the title-page of the volume, 1782.

PETALIA Walker, 1854, List Specimens lepid. Insects Colln. Rr. Mus. 2: 457

Br. Mus. 2: 457.

Type-species: Hypsa plagiata Walker, 1854, ibidem 2: 457,

by monotypy. [AUSTRALIA] New Holland.

A junior homonym of *Petalia* Gray, 1838, *Mag. Zool. Bot.* 2: 494, – Mammalia. There is no objective replacement name but *H. plagiata* is currently considered to be congeneric with *Phalaena javana* Cramer, 1780, the type-species of *Asota* Hübner, [1819]. The latter is therefore available for use as a subjective replacement name.

Petalia was established as a subgenus of Hypsa Hübner, [1819].

ASOTOMORPHA Hulstaert, 1923. See Caryatis Hübner, [1819].

ASPA Walker, 1854. See Hypsa Hübner, [1819]. AGAN

ASPARUS Watson nom. n. for Sarapus Walker, 1855. LITH Type-species: Sarapus bicolor Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 761, by monotypy (of Sarapus Walker, 1855). [SOUTH AFRICA]: Cape.

SARAPUS Walker, 1855, List Specimens lepid. Insects Colln Br., Mus. 3; 760.

Type-species: Sarapus bicolor Walker, 1855, by monotypy. A junior homonym of Sarapus Fischer, 1821, Entomographia Imperii Russici 1 (Genera Coleopterorum): 39, 87, – Insecta, Coleoptera. The objective replacement name is Asparus Watson nom. n.

ASPHENOPTERA Watson nom. n. for Sphenoptera Felder, CTEN

Type-species: *Sphenoptera triangulifera* Felder, 1874, *in* Felder & Rogenhofer, *Reise öst. Fregatte Novara* (Zool.) **2** (Abt.2): pl.102, fig.34, by subsequent designation by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 332 (for *Sphenoptera* Felder, 1874). [BRAZIL]: Amazon R., cited by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 9.

SPHENOPTERA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, figs 34, 35.

Type-species: Sphenoptera triangulifera Felder, 1874, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 332.

A junior homonym of *Sphenoptera* Dejean, 1833, *Cat. Coléoptères* 2: 81. The objective replacement name is *Asphenoptera* Watson nom. n.

ASTACOSIA Toulgoët, 1958, Mém. Inst. scient. Madagascar (E) 9: 179.

Type-species: *Philenora oblonga* Toulgoët, 1955, *Revue fr. Ent.* **22**: 200, pl.6, figs 20, 21, by original designation. MADAGASCAR: E. Moramanga.

ASTAPA Dognin, 1911, Hétérocères nouv. Am. Sud 2: 40.

Type-species: Astapa signata Dognin, 1911, ibidem 2: 41, by original designation. Colombia: San Antonio, Alto de las Cruces, near Cali.

‡ASTOPA; Grünberg, 1913, Arch. Naturgesch. 78 (B) 7: 89. An incorrect subsequent spelling of Astapa Dognin, 1911.

ASTAURA Kiriakoff, 1964, Revue Zool. Bot. afr. 69: 64.

Type-species: *Astaura umbrosa* Kiriakoff, 1964, *ibidem* 69: 65, fig.1, by original designation. [ZAIRE]: Uele, Paulis.

‡ASTEPHANA Agenjo, 1946, Graellsia 4 (6): Cat. Lepid. España, Family 59, Agrotidae page [7].

‡Astephana was attributed to Boursin and used in a checklist as a subgenus of Leucania Ochsenheimer, 1816. This use of ‡Astephana did not establish the name under the Code, Article 13(a) or (b), as it was neither accompanied by a statement differentiating the taxon, nor by a bibliographic reference to such a statement, nor was it accompanied by the fixation of a type-species. Of the six originally included nominal species Phalaena comma Linnaeus, 1761, is the type-species of Leucania Ochsenheimer, 1816; and Noctua loreyi Duponchel, 1827, is the type-species of Acantholeucania Rungs, 1953.

No other published reference to ‡Astephana has been found. ‡Astephana in the Noctuidae, Hadeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡ASTOPA; misspelling. See Astapa Dognin, 1911.

ASTRALARCTIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 63.

Type-species: Automolis pulverosa Schaus, 1905, Proc. U.S. natn. Mus. 29: 215 SURINAM: Surinam River, Geldersland.

ASTRIDIA Kiriakoff, 1948. See Pleurosoma Orfila, 1935.

ASTYLIS Boisduval, 1872. See Crinodes Herrich-Schäffer, 1855. NOTO

ASURA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 484.

Type-species: Asura cervicalis Walker, 1854, ibidem 2: 484, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 299. [AUSTRALIA] New Holland: and [Tasmania] Van Dieman's Land.

‡AZUREA; Strand & Ramme, 1911, Arch. Naturgesch. 75 (2) 2: 301.

An incorrect subsequent spelling of Asura Walker, 1854.

ASURGYLLA Kiriakoff, 1958, Ruwenzori Exped. 1952 1: 7.

Type-species: Asurgylla collenettei Kiriakoff, 1958, ibidem 1: 8, figs 14, 62, by original designation. UGANDA: Fort Portal.

ASURIDIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: vi, 89, 412.

Type-species: Ammatho carnipicta Butler, 1877, Trans. ent. Soc. Lond. 1877: 342, by original designation. MONGOLIA.

ASURIDOIDES Daniel, 1951, Bonn. zool. Beitr. 2: 308. LITH Type-species: Asuridoides atuntseica Daniel, 1951, ibidem 2: 308, text-fig. 10, by original designation. China: N. Yunnan, A-tun-tse.

ASUROPSIS Matsumura, 1927, J. Coll. Agric. Hokkaido imp.

Type-species: Asuropsis ranruna Matsumura, 1927, ibidem 19 (1): 70, pl.5, fig.5, by original designation. [TAIWAN] Formosa: Ranrun.

ASYTHOSIA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 67.

Type-species: Asythosia velutina Birket-Smith, 1965, ibidem 1: 69, figs 53-55, by original designation. NIGERIA: Mt Cameroon, Buea.

ATASCA Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 139.

Type-species: Phalaena pellex Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 510, by subsequent designation by Roepke, 1949, Trans. R. ent. Soc. Lond. 100: 47. [INDIA] 'Indiis'.

ATELOPHLEPS Turner, 1940, Proc. R. Soc. Qd 51: 74.

Type-species: Atelophleps tridesma Turner, 1940, ibidem 51: 74, by monotypy. Australia: N. Queensland, Lake Barrine.

ATEUCHETA Turner, 1940, Proc. R. Soc. Qd 51: 65. LITH Type-species: Poliosia zatesima Hampson, 1914 [November], Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: 463, pl.25, fig.26, by monotypy (but included as zetesima Turner, a junior objective synonym, see below). AUSTRALIA: near Cairns, Kuranda.

The name zatesima originated from Turner but was used in conjunction with a description and made available by Hampson prior to its proposal for a new species (but spelt as zetesima)

by Turner, 1914 December, Proc. R. Soc. Od 26: 16. Both nominal species were based on the same type-specimen and so are treated as objective synonyms.

See Athrula Walker, 1862. ‡ATHRALA; misspelling. LYMA

ATHRULA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 82.

Type-species: Athrula saturnioides Walker, 1862, ibidem 1: 83, by monotypy. BRAZIL.

A. saturnioides is a junior subjective synonym of Rolepa delineata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 770.

Athrula was established in the Limacodidae; it was transferred to the Lymantriidae by Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 557.

See also Rolepa Walker, 1855, and Olenoptera Herrich-Schäffer, [1856].

‡ATHRALA; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 558.

An incorrect subsequent spelling of Athrula Walker, 1862.

ATOLMIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 164.

Type-species: Phalaena rubricollis Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 511, by monotypy. EUROPE.

Atolmis when established contained the name of a second species ‡flavicapilla, at that time a nomen nudum (see Hemming, 1937, Hübner 1: 494; 2: 160) and therefore not eligible for selection as type-species. The name of the species was later established as Atolmis flavicapilla Hübner, [1819], Samml. exot. Schmett. 2: pl.[180], - Lepid., Dioptidae.

GNOPHRIA Stephens, 1829 February, Illust. Br. Ent. (Haustellata) 2: 89, 98.

Type-species: Phalaena rubricollis Linnaeus, 1758, by

A junior objective synonym of *Atolmis* Hübner, [1819].

The name Gnophria was proposed again as new by Stephens, 1829 [June], Nom. Br. Insects: 40, and by Stephens, 1829 [July], Syst. Cat. Br. Insects 2: 61.

ATOPOMIMA Meyrick, 1936, Exot. Microlepid. 5: 61. LITH Type-species: Atopomima philocosma Meyrick, 1936, ibidem 5: 61, by monotypy. Solomon Islands: New Georgia, Arundel.

A. philocosma is a junior subjective synonym of Scaptesyle caerulescens Butler, 1889, Trans. ent. Soc. Lond. 1889: 390.

ATORNOPTERA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 373.

Type-species: Atornoptera discocellularis Kiriakoff, 1974, ibidem 17: 374, text-fig.1, pl.1, fig.1, by original designation. SUMATRA: Dolok Merangir.

ATRASANA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1749.

Type-species: Atrasana postica Walker, 1856, ibidem 7: 1750, by monotypy. [South Africa]: [Natal], [Durban] Port Natal.

ATUCIA Watson nom. n. for Acutia Kaye, 1919. Type-species: Acutia bidensis Kaye, 1919, Ann. Mag. nat. Hist. (9) 4: 90, by original designation (for Acutia Kaye, 1919). Borneo: Sarawak, Bidi.

ACUTIA Kaye, 1919, Ann. Mag. nat. Hist. (9) 4: 89.

Type-species: Acutia bidensis Kaye, 1919, by original

A junior homonym of Acutia Ragonot, 1891, Annls Soc. ent. Fr. (6) 10: 539, - Lepid., Pyralidae. The objective replacement name is Atucia Watson nom. n.

ATYPHOPSIS Butler, 1878, Trans. ent. Soc. Lond. 1878; 49.

Type-species: Atyphopsis modesta Butler, 1878, ibidem 1878: 49, by original designation. [BRAZIL]: 'Boaventura, Rio Jatahi' and [Para], Santarém.

‡AUGE Oken, 1815. See Cosmosoma Hübner, [1823]. CTEN

‡AUSTROCAREA Holloway, 1976. See Austrocarea Holloway, 1977.

AUSTROCAREA Holloway, 1977, Series Ent. 13: 75, 139.
NOCTUIDAE

Type-species: *Maurilia iocephala* Turner, 1902, *Proc. Linn. Soc. N.S.W.* **27**: 94, by original designation. Holotype \mathfrak{P} , Australia: Queensland, Townsville.

Austrocarea in the Noctuidae, Chloephorinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡AUSTROCAREA Holloway, 1976, Norfolk Island Philatelic Bull. 1976 (2): [3].

An unavailable name under the Code, Article 13(a)(i), as it was published after 1930 and was not accompanied by a statment that purported to differentiate the taxon.

AUTHYALA Warren, 1905, Novit. zool. 12: 311.

Type-species: Authyala obliquaria Warren, 1905, ibidem 12: 312, by original designation. PERU: Cuzco.

Authyala was established in the Cyllopodinae, now Geometridae, Sterrhinae; it was transferred to the Dioptidae by Prout, 1918, Novit. zool. 25: 414.

AUTOCERAS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.24. LITH

Type-species: *Autoceras grammophora* Felder, 1874, *ibidem* **2** (Abt.2): pl.106, fig.24, by monotypy. [SRI LANKA] Ceylon: cited by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 2.

A. grammophora is a junior subjective synonym of Aemene taprobanis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 542.

See also Aemene Walker, 1854, and Panassa Walker, 1865.

AUTOCHLORIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 125.

Type-species: *Sphinx almon* Cramer, 1779, *Uitlandsche Kapellen (Papillons exot.)* 3: 52, pl.224, fig.F, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 125. SURINAM.

S. almon is dated from the wrapper of the part and not from the title-page of the volume, 1782.

AUTOMOLIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 170.

Type-species: *Sphinx meteus* Stoll, 1781, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 109, pl.347, fig.B, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 220. [SOUTH AFRICA]: [Cape of Good Hope] Kaap de Goede Hoop.

S. meteus is dated from the wrapper of the part and not from the title-page of the volume, 1782.

Incorrect type-species: Glaucopis sphingidea Perty, 1834, a name not originally included in Automolis and not linked with one of the originally included names when designated by Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7:

Automolis was established in the 'Phalaenae'; it was placed in the 'Zygaenides' by Walker, 1856, ibidem 7: 1634; was placed in the Arctiidae by Kirby, 1892, ibidem 1: 220; and its type-species was placed in the Thyretidae by Kiriakoff, 1956, Lambillionea 56: 26.

AUTOMOLIS Walker, 1854. See Cyanopepla Clemens, 1861.

AVELA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1658.

Type-species: Avela diversa Walker, 1856, ibidem 7: 1659, by monotypy. BRAZIL.

AXANA Swinhoe, 1922, Ann. Mag. nat. Hist. (9) 10: 471.

Type-species: *Naxa puncticilia* Moore, 1872, *Proc. zool. Soc. Lond.* **1872**: 575, by monotypy. INDIA: [S.], [Nilgiri Hills] Nilghiris.

AXIOCLETA Turner, 1911, Ann. Qd Mus. 10: 133. THAU Type-species: Axiocleta perisema Turner, 1911, ibidem 10: 134, by original designation. AUSTRALIA: Queensland, Evelyn Scrub, near Herberton.

Axiocleta was established in the Eupterotidae; it was placed in the Notodontidae, Cnethocampinae, now Thaumetopoeidae, by Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 364.

AXIOLOGA Turner, 1904, Trans. ent. Soc. Lond. 1904: 477.

Type-species: *Teara pura* Lucas, 1892, *Proc. R. Soc. Qd* 8: 75, by monotypy. AUSTRALIA: [Queensland], Brisbane.

‡AXIOPAENELLA; misspelling. See Axiopoeniella Strand, 1909.

‡AXIOPENA; misspelling. See Axiopoena Ménétriés, 1842.

AXIOPOENA Ménétriés, 1842, Bull. scient. Acad. imp. Sci. St. Petersb. 9; 42.

Type-species: *Bombyx maura* Eichwald, 1830, *Zool. spec.* 2: 196, by monotypy. [U.S.S.R.]: [Turkmeniya], eastern coast of Caspian Sea, near Krasnovodsk.

‡AXIOPENA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 421.

An incorrect subsequent spelling of Axiopoena Ménétriés, 1842.

AXIOPOENIELLA Strand, 1909, Annls Soc. ent. Belg. 53: 357.

Type-species: *Lithosia laymerisa* Grandidier, 1867, *Revue zool.* (2) **19**: 274, by original designation. MADAGASCAR: S.W., Tullear.

‡AXIOPAENELLA; Toulgoët, 1956, Bull. Soc. ent. Fr. 61: 183.

An incorrect subsequent spelling of Axiopoeniella Strand, 1909.

AXUENNA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 382.

Type-species: *Eloria discalis* Walker, 1856, *ibidem* 7: 1726, by monotypy. [Brazil]: [R. Amazon], [Tefé] Ega.

AZA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 384.

APATELODIDAE

Type-species: *Aza micacea* Walker, 1865, *ibidem* **32**: 384, by monotypy. [Colombia]: Bogota.

A junior homonym of Aza Mulsant, 1850, Annls Soc. Agric. Lyon (2) 2: 424, 1085, – Insecta, Coleoptera. There is no objective replacement name, but A. micacea is congeneric with Colla glaucescens Walker, 1865, the type-species of Colla Walker, 1865, ibidem 32: 580: the latter is thus available for use as a subjective replacement name.

Aza was established in the Liparidae, now Lymantriidae. It was placed in the Epiinae, now included in the Apatelodidae,

by Schaus, 1929, in Seitz, Gross-Schmett. Erde 6: 679, who

included the type-species in Colla.

Schaus, 1927, in Seitz, ibidem 6: 537, included Aza in the Lymantriidae and stated that the type of Aza micacea was lost. Two syntypes, one male and one female, have since been found in the BMNH and A. micacea is placed as a senior subjective synonym of Colla albescens Schaus, 1905, Proc. U.S. natn. Mus. 29: 299.

AZATREPHES Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 430.

Type-species: *Halysidota discalis* Walker, 1856, *List Specimens lepid. Insects Colln Br. Mus.* 7: 1706 (but cited by Walker as ‡*Halesidota*, an incorrect subsequent spelling), by monotypy (but included by Hampson as ‡*Halisidota*, an incorrect subsequent spelling). [BRAZIL]: Amazon valley.

Azatrephes was established for H. discalis Walker, 1856, and its junior synonym Zatrephes paradisea Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: xii, 52, fig. Hampson's 1905, ibidem (7) 15: 430 opening remarks 'Page 474. Insert Azatrephes, n.n. for Zatrephes, Hmpsn. (nec Hübn.)' [referring to Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 474] are misleading. Hampson, 1905, ibidem (7) 15: 430 was not proposing a replacement name but was establishing the genus-group name of Azatrephes for the nominal species H. discalis Walker, 1856.

AZAXIA Dyar, 1908, Proc. ent. Soc. Wash. 9: 46.

Type-species: Heterocampa luteilinea Druce, 1904, Ann. Mag. nat. Hist. (7) 13: 249, by original designation. Peru: south-eastern, Santo Domingo.

‡AZUREA; misspelling. See Asura Walker, 1854. LITH

‡BAENASA; misspelling. See Boenasa Walker, [1865].

BAEOMORPHA Turner, 1940. See Hobapromea Watson nom. n.

BAGISARA Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1791. See Caroia Walker, 1858.

BAHAIA Dyar, 1924, Insecutor Inscit. menstr. 12: 17. NOTO Type-species: Bahaia sceletaria Dyar, 1924, ibidem 12: 17, by monotypy. MEXICO: Colima, Jalapa.

BALACA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 129.

Type-species: *Balaca picaria* Walker, [1865] 1864, *ibidem* 31: 130, by monotypy. Holotype ♀ [not male as stated by Walker], [New Guinea]: [Irian Jaya], [Waigeo I.] Waigiou, (BMNH).

The holotype is the only known specimen of *B. picaria*. It is in very poor condition but sufficient remains to show that it is probably not correctly placed in the Arctiidae.

BALACRA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1721.

Type-species: *Balacra caeruleifascia* Walker, 1856, *ibidem* 7: 1721, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 221. Type-locality: not stated, [Sierra Leone], [Ghana].

Balacra was established in the Arctiidae; it was placed in the Thyretidae by Kiriakoff, 1953, Annls Mus. r. Congo Belge

26: 57

BALACRELLA Kiriakoff, 1957, Bull. Annls Soc. r. ent. Belg. 93: 154.

Type-species: *Pseudapiconoma affinis* Rothschild, 1910, *Novit. zool.* 17: 442, by original designation. [ZAIRE] Congo Free State: Kassai District.

Balacrella was established as a subgenus of Balacra Walker, 1856.

BALBURA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 481.

Type-species: *Balbura dorsisigna* Walker, 1854, *ibidem* **2**: 481, by monotypy. VENEZUELA, HONDURAS.

BALIA Doubleday, 1869, in Scudder, Occ. Pap. Boston Soc. nat. Hist. 1: 132.

Type-species: *Stauropus menas* Doubleday, 1869, *ibidem* 1: 134, by PRESENT DESIGNATION. [U.S.A.]: East Florida, St John's Bluff.

S. menas is a junior subjective synonym of Heterocampa astarte Doubleday, 1841, Entomologist 1: 57.

Doubleday did not make clear whether he intended to establish *Balia* for only one nominal species, *Stauropus menas* Doubleday, 1869 (his species no.60), or whether he intended to include also *Stauropus athereo* Doubleday, 1869 (his species no.56) and *Stauropus manteo* Doubleday, 1869 (his species no.57). For reasons of clarity the above designation has been made.

Balia Doubleday, 1869, is not preoccuppied by ‡Balia Swainson, 1840, Treatise Malacology: 182, 334, an incorrect subsequent spelling of Balea Gray, 1824.

See also *Heterocampa* Doubleday, 1841, and *Trichotis* Felder, 1874.

BALIOPTERYX Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 84.

Type-species: *Spatalia argyrophora* Hampson, 1910, *Proc. zool. Soc. Lond.* **1910**: 465, pl.39, fig.14, by original designation. [ZAIRE] Congo: S.E. Katanga.

BARADESA Moore, 1883, Proc. zool. Soc. Lond. 1883: 16.
NOTO

Type-species: *Baradesa lithosioides* Moore, 1883, *ibidem* 1883: 17, pl.5, fig.2, by monotypy. [INDIA]: Darjeeling.

BARBAROSSULA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 285.

Type-species: *Barbarossula rufibarbis* Kiriakoff, 1964, *ibidem* 14: 286, abb.5, fig.47, pl.5, fig.50, by original designation. China: south Shensi, Tapaishan im Tsinling.

BARDAXIMA Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 14: 1343 (key), 1349.

Type-species: *Bardaxima lucilinea* Walker, 1858, *ibidem* 14: 1349, by monotypy. Brazil.

Bardaxima was established in the Noctuidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 616.

GOZARTA Walker, 1869, Characters undescr. Lepid. Heterocera: 18.

Type-species: Gozarta fulgurifera Walker, 1869, ibidem: 18,

by monotypy. [Honduras]: [Limas] Limar.

A junior homonym of Gozarta Walker, [1868] 1870, J. Linn. Soc. (Zool.) 10: 116, — Insecta, Hemiptera. There is no objective replacement name, but G. fulgurifera is a junior subjective synonym of Bardaxima lucilinea Walker, 1858, the type-species of Bardaxima Walker, 1858. The latter is therefore available for use as a subjective replacement name.

‡BARECONIA; misspelling. See Boreconia Walker, 1865.

BARHONA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 55.

Type-species: *Barhona carneola* Moore, 1879, *ibidem* (1): 56, by monotypy. INDIA: [W. Bengal, Darjeeling] Darjiling.

BARITIUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 712.

Type-species: Baritius discalis Walker, 1855, ibidem 3: 713. by monotypy. [BRAZIL]: Rio [de] Janeiro.

BARLOWIA Talbot, 1929, Bull. Hill Mus. Witley 3: 128.

LYMA

Type-species: Barlowia zelotes Talbot, 1929, ibidem 3: 129, text-fig.2, pl.3, figs 3, 4, by original designation. [MALAWI] Nyasaland: Zomba.

BAROA Moore, 1878, Proc. zool. Soc. Lond. 1878: 28. ARCT Type-species: Cycnia punctivaga Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 682, by original designation. JAVA.

BAROA Hampson, 1896, Fauna Br. India (Moths) 4: 493.

Type-species: Cycnia punctivaga Walker, 1855, by original

A junior homonym of Baroa Moore, 1878, Proc. zool. Soc. Lond. 1878: 28, - Lepid., Arctiidae. The objective replacement name is Baroa Moore, 1878.

BAROA Hampson, 1896. See Baroa Moore, 1878. ARCT

BAROBATA Karsch, 1895, Ent. Nachr. Berlin 21: 371. LYMA Type-species: Barobata trocta Karsch, 1895, ibidem 21: 371, pl.3, fig.3, by original designation. Togo: [approx. 250 km NNW of Lome] Station Bismarckburg.

BARSINE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 546.

Type-species: Barsine defecta Walker, 1854, ibidem 2: 546, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 306. [NEPAL] Nepaul.

BARSINELLA Butler, 1878, Proc. zool. Soc. Lond. 1878: 385.

Type-species: Barsinella mirabilis Butler, 1878, ibidem 1878: 385, fig., by original designation. BRAZIL: [Espirito Santo] Espiritu Santo.

BARYAZA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 45. LYMA

Type-species: Baryaza cervina Moore, 1879, ibidem (1): 45, pl.3, fig.1, by monotypy. INDIA: [W. Bengal, Darjeeling] Darjiling.

BARYDIA Guenée, 1852, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 6: 409.

Type-species: Barydia bufo Guenée, 1852, ibidem 6: 410, pl.18, fig.1, by monotypy. French Guiana: Cayenne.

Barydia was established in the Noctuidae; it was placed in the Notodontidae by Nye, 1975, Generic Names Moths World 1: 76.

BATEMPA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 40.

Type-species: Batempa plana Kiriakoff, 1962, ibidem 114: 40, fig.51, by original designation. [ZAIRE]: Sankaru, Km 45, route Lusambo - Batempa.

BATHMOCHTHA Karsch, 1895, Ent. Nachr. Berlin 21: 368.

Type-species: Bathmochtha albilunulata Karsch, 1895, ibidem 21: 368, pl.4, fig.12, by original designation. [CAMEROUN] Kamerun: Hinterland, Yaunde-Station.

B. albilunulata is a junior subjective synonym of Dediama antica Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1074.

Bathmochtha was established in the 'Arctiiden'; it was

placed in the Lymantriidae by Bryk, 1934, in Strand, Lepid. Cat. 62: 6.

See also Dediama Walker, 1855.

BATUANA Laporte, 1976, Bull. mens. Soc. linn. Lyon 45: 131.

Type-species: Batuana rougeoti Laporte, 1976, ibidem 45: 131, fig.1, by original designation. Holotype &, Етнюріл: environs du Mont Batu, (MNHN, Paris).

Batuana in the Noctuidae. Hadeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

BAZISA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 398.

Type-species: Bazisa detecta Walker, 1865, ibidem 32: 399, by monotypy. [INDIA]: Hindostan.

‡BAZIZA; Hampson, [1893] 1892, Fauna Br. India (Moths)

An incorrect subsequent spelling of Bazisa Walker, 1865; used again by Hampson, 1905, Ann. S. Afr. Mus. 3: 393, and by Hampson, 1910, Proc. zool. Soc. Lond. 1910: 455.

‡BAZIZA; misspelling. See Bazisa Walker, 1865. LYMA

BEERIA Hartig, 1963. See Casama Walker, 1865. LYMA

BELCIANA Walker, [1865]. See Dialephtis Felder, 1874. DIOP

BELEMNIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 211.

Type-species: Zygaena eryx Fabricius, 1775, Syst. Ent.: 552, by monotypy. [South America]: [tropical] America meridionalis.

Belemnia was established as a subgenus of Euchromia Hübner, [1819], in the 'Zygaenides'; it was placed in the Arctiidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 170; and was transferred to the Ctenuchidae by Kiriakoff, 1950, Lambillionea 50: 64.

CHRYSAOR Hübner, [1809], Samml. exot. Schmett. 1: pl.[161].

Type-species: Zygaena eryx Fabricius, 1775, by monotypy. A junior homonym of Chrysaor Montfort, 1808, Conch. 1: 378, - Mollusca, Cephalopoda. The objective replacement name is Bélemnia Walker, 1854.

Chrysaor Hübner, [1809], has been placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1832.

BELEMNIASTIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 5, 213.

Type-species: Belemnia whiteleyi Druce, 1882, Entomologist's mon. Mag. 25: 62, by original designation. [GUYANA]

Belemniastis was established in the 'Arctiadae' and is here transferred to the Ctenuchidae.

BELISARIA Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 331.

Type-species: Belisaria camerunica Kiriakoff, 1965, ibidem 101: 332, fig.1, by original designation. Cameroun: Lomié.

BELLURA Walker, 1865, List Specimens lepid. Insects Colln NOCTUIDAE Br. Mus. 32: 465.

Type-species: Bellura gortynoides Walker, 1865, ibidem 32: 465, by monotypy. U.S.A.

Bellura was established in the Notodontidae; it was placed in the Noctuidae by Smith, 1893, Bull. U.S. natn. Mus. 44: 181.

BEMBINA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 505.

Type-species: Bembina apicalis Walker, 1865, ibidem 32:

505, by monotypy. [SRI LANKA] Ceylon.

Bembina was established in the Limacodidae; it was placed in the Liparidae, now Lymantriidae, by Moore, [1883] 1882-3, Lepid. Ceylon 2: 87.

BENBOWIA Kiriakoff, 1967, Tijdschr. Ent. 110: 52. NOTO Type-species: Stauropus virescens Moore, 1879, Proc. zool. Soc. Lond. 1879: 404 [INDIA]: Darjeeling.

BEPARA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 143.

Type-species: Bepara egaca Walker, [1865] 1864, ibidem 31: 143, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 400, but cited as Phaeoclaena subapicalis Walker, 1854, a nominal species not originally included in Bepara. Kirby, however, placed subapicalis as the senior synonym of egaca, a nominal species originally included in Bepara. Under the Code, Article 69 (a)(iv), this designation constitutes the fixation of the originally included name as the type-species. [Brazil]: [Tefé] Ega.

BERALADE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 852. LASIOCAMPIDAE

Type-species: Beralade perobliqua Walker, 1855, ibidem 4:

853, by monotypy. [South Africa]: Caffraria.

Beralade was established in the Liparidae, now Lymantridae; it was placed in the Lasiocampidae by Distant, 1897, Ann. Mag. nat. Hist. (6) 20: 206.

BERGERIA Kiriakoff, 1952, Revue Zool. Bot. afr. 46: 397.

Type-species: Bergeria haematochrysia Kiriakoff, 1952, ibidem 46: 397, figs 1-3, by original designation. [ZAIRE] Congo Belge: Sankuru, Lusambo.

BERNARDITA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 191.

Type-species: *Peratodonta albiplagiata* Gaede, 1928, *in* Seitz, *Gross-Schmett. Erde* **14**: 406, pl.68 row i, by original designation. [Ghana] Goldküste.

BERNATHONOMUS Fragoso, 1953, Revta bras. Biol. 13: 179.

Type-species: *Phaegoptera piperita* Herrich-Schäffer, [1855] 1850-1858, *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.53, fig.293, by original designation. BRAZIL.

BERTHOLDIA Schaus, 1896, Jl N.Y. ent. Soc. 4: 136. ARCT Type-species: Phaegoptera specularis Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.14, fig.59 (but attributed to Boisduval, an incorrect authorship), by original designation. Venezuela.

BESAIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 458.

Type-species: *Besaia rubiginea* Walker, 1865, *ibidem* 32: 458, by monotypy. [INDIA]: Hindostan.

BESIDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 456.

Type-species: *Besida xylinata* Walker, 1865, *ibidem* **32**: 456, by monotypy. JAVA.

BETASHACHIA Matsumura, 1925, Zool. Mag. Tokyo 37: 398.

Type-species: *Betashachia angustipennis* Matsumura, 1925, *ibidem* 37: 398, pl.7, fig.7, by original designation. TAIWAN.

BETOLA Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 260 (key), 289.

Type-species: *Betola aroata* Schaus, 1901, *ibidem* **1901**: 290, pl.11, fig.6, by original designation. VENEZUELA: Aroa.

BETTONIA Butler, 1898, Proc. zool. Soc. Lond. 1898: 418.

Type-species: *Bettonia ferruginea* Butler, 1898, *ibidem* 1898: 418, pl.32, fig.5, by original designation. [Kenya]: Voi.

B. ferruginea is a junior subjective synonym of Crocota unicolor Hopffer, 1858, Mber. K. preuss. Wiss. 1857: 422. See also Ochrota Hopffer, 1862.

BICELLUPHORA Janse, 1915, Ann. Transv. Mus. 5: 16.

Type-species: *Bicelluphora argentea* Janse, 1915, *ibidem* 5: 17, pl.2, fig.4, pl.3, fig.2, by original designation. SOUTH AFRICA: [Transvaal], Barberton.

‡BIERA; misspelling. See Sericochroa Felder, 1874. NOTO

BIFRONTIPTA Berio, 1974, Annali Mus. civ. Stor. nat. Giacomo Doria 80: 223.

NOCTUIDAE

Type-species: *Bifrontipta saskai* Berio, 1974, *ibidem* 80: 224, by original designation. Holotype 9, [ETHIOPIA] Abessinien: Addis Abeba, (NR, Stockholm).

Bifrontipta in the Noctuidae, Noctuinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

BIFURCA Joannis, 1930, Annls Soc. ent. Fr. 98: 781. LYMA Type-species: Bifurca longinasus Joannis, 1930, ibidem 98: 782, by original designation. [VIETNAM]: Tonkin, Cha pa.

Bifurca and B. longinasus are dated from the wrapper of the

part and not from the title-page of the volume, 1929.

Bifurca was established in the Lymantriidae; was transferred to the Noctuidae, Hypeninae by Collenette, 1935, Stylops 4: 242; and was transferred back to the Lymantriidae by Nye, 1975, Generic Names Moths World 1: 80.

BIFURCIFER Ebert, 1968, Reichenbachia 10: 203. NOTO Type-species: Bifurcifer afghanus Ebert, 1968, ibidem 10: 204, fig.4, by original designation. Afghanistan: south-eastern, Safed Koh, Kotkai.

BILLETIA Joannis, 1900, Bull. Soc. ent. Fr. 1900: 281.

Type-species: *Billetia rubripuncta* Joannis, 1900, *ibidem* 1900: 281, by monotypy. [VIETNAM]: Tonkin, Cao-Bang.

B. rubripuncta is a junior subjective synonym of Cerasana anceps Walker, 1862, J. Linn. Soc. (Zool.) 6: 123.

See also Cerasana Walker, 1862.

BILULUA Kiriakoff, 1954, Revue Zool. Bot. afr. 49: 319.

Type-species: *Bilulua atricollis* Kiriakoff, 1954, *ibidem* 49: 320, pl.1, fig.14, pl.2, fig.14, by original designation. [ZAIRE] Congo Belge: Katanga, Elisabethville.

BINNA Walker, [1865] 1864, List Specimens lepid. Insects
Colln Br. Mus. 31: 319.

ARCT
Type-species: Binna penicillata Walker, [1865] 1864, ibidem

Type-species: *Binna penicillata* Walker, [1865] 1864, *ibidem* 31: 319, by monotypy. SIERRA LEONE.

BIOTES Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 91. NOTO

Type-species: *Phalaena bitia* Stoll, 1780, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* **4**: 69, pl.322, fig.F, by monotypy. [West Africa]: Coast of Guinea.

Biotes was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume

6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the

Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

BIRAIA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 152 (key), 174.

Type-species: Ceira postica Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 66, by original designation. INDIA: Darjeeling.

Biraia was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 75.

BIRETA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1754.

Type-species: Bireta longivitta Walker, 1856, ibidem 7: 1754, by monotypy. INDIA: Hindostan.

See also Menapia Walker, 1865.

BIRGORIMA Watson nom. n. for Mirobriga Walker, 1864.

Type-species: Mirobriga pulchripicta Walker, 1864, J. Proc. Linn. Soc. (Zool.) 7: 196, by monotypy (of Mirobriga Walker, 1864). [BORNEO]: Sarawak.

MIROBRIGA Walker, 1864, J. Proc. Linn. Soc. (Zool.) 7: 196. Type-species: Mirobriga pulchripicta Walker, 1864, by

A junior homonym of Mirobriga Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 27: 131, - Lepid., Pyralidae. The objective replacement name is Birgorima Watson nom. n.

BIRNARA Butler, 1879, Trans. Linn. Soc. Lond. (2) Zool. 1:

Type-species: Birnara nubila Butler, 1879, ibidem (2) Zool. 1: 560, by original designation. [MALAYSIA]: [Malaya], Malacca.

BISOLITA Kiriakoff, 1961, Bull. Annls Soc. r. ent. Belg. 97:

Type-species: Ramesa strigata Aurivillius, 1906, Ark. Zool. 3(1): 7, by original designation. [ZAIRE] Congo: Mukimbungu.

Bisolita was established as a subgenus of Xanthodonta Gaede, 1928; it was raised to generic level by Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 91.

BITECTA Heylaerts, 1891, Annls Soc. ent. Belg. 35 (Bull.): ccccxi.

Type-species: Lithosia murina Heylaerts, 1891, ibidem 35 (Bull.): ccccxi, by monotypy. JAVA: Préanger.

Bitecta was established as a subgenus of Lithosia Fabricius, 1798.

BITHRA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 621.

Type-species: Bithra spilosomoides Walker, 1865, ibidem 32: 621, by monotypy. South Africa.

BITHRA Fawcett, 1918. See Galtara Walker, [1863]. ARCT

BITURIX Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 717. ARCT

Type-species: Biturix diversipes Walker, 1855, ibidem 3: 718, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 204. VENEZUELA.

‡BITURYX; Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 175.

An incorrect subsequent spelling of Biturix Walker, 1855.

‡BITURYX; misspelling. See Biturix Walker, 1855. ARCT

BIULA Walker, 1857, List Specimens lepid. Insects Colln Br. Mus. 11: 714.

Type-species: Biula lignilutea Walker, 1857, ibidem 11: 714, by monotypy. JAVA.

Biula was established in the Noctuidae; it was placed in the Notodontidae by Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 213.

BIZONE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 548.

Type-species: Bizone perornata Walker, 1854, ibidem 2: 548, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 303. [BANGLADESH]: [Sylhet] Silhet,

BLABIOIDES Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 84, 279.

Type-species: Pseudoblabes snelleni Ritsema, 1875, Petites Nouv. Ent. 7: 479, by original designation. JAVA: Ambarawa.

BLACODES Kiriakoff, 1959. See Neoblacodes Fletcher nom.

BLAKEIA Kiriakoff, 1967, Tijdschr. Ent. 110: 39. Type-species: Blakeia marmorata Kiriakoff, 1967, ibidem 110: 41, fig.4, by original designation. BORNEO: south-eastern, Samarinda.

BLAVIA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 108.

Type-species: Blavia caliginosa Walker, 1862, ibidem 6: 108, by monotypy. [BORNEO]: Sarawak.

BLAVIODES Bethune-Baker, 1910, Ann. Mag. nat. Hist. (8) 6: 441.

Type-species: Blaviodes ochracea Bethune-Baker, 1910, ibidem (8) 6: 441, by original designation. NEW GUINEA: [Irian Jaya], Arfak Mts.

BLAZIA Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 536 (key), 555.

Type-species: Orgya lixivia Dognin, 1923, Hétérocères nouv. Am. Sud 21: 10, by original designation. BOLIVIA: rio Songo. O. lixivia was established in Orgya, an unjustified emendation by Zetterstedt, [1839], of Orgyia Ochsenheimer, 1810.

The date of publication of the part of Seitz containing page 536 is cited on page 529 as 17.v.1927 in both the German and the English editions.

BLENNEMA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 152 (key), 190.

Type-species: Pydna griseodivisa Bryk, 1950, Ark. Zool. 42 (A) 19: 14, pl.4, fig.9, by original designation. Burma: north, Kambaiti.

Blennema was established as a subgenus of Bireta Walker. 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 98.

BLERA Walker, 1862. See Sericochroa Felder, 1874. NOTO

BOADICEA Tams, 1930, Ann. Mag. nat. Hist. (10) 5: 484.

Type-species: *Boadicea pelecoides* Tams, 1930, *ibidem* (10) 5: 485, pl.18, fig.1, by original designation. [Congo] French Congo: 'Popos'.

BODOSA Machado & Rego Barros, 1969, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 270: 2.

Type-species: *Euchromia tina* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 233, by original designation. [BRAZIL]: [Amazonas], Rio Grande [do Sul].

Bodosa was established as a subgenus of Euchromia Hübner,

[1819]

BOENASA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 266.

Type-species: *Boenasa nigrorosea* Walker, [1865] 1864, *ibidem* 31: 266, by monotypy. [HISPANIOLA]: [Dominican Republic], S[an]t[o] Domingo.

‡BAENASA; Druce, 1894, Ann. Mag. nat. Hist. (6) 13: 177. An incorrect subsequent spelling of Boenasa Walker, [1865].

BOMBILIODES Hampson, 1898. See *Gymnelia* Walker, 1854.

‡BOMBISATUR; misspelling. See ‡Bombysatur Corporaal, 1921.

BOMBOPSYCHE Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xii, 24, 161.

Type-species: Antichloris ruatana Druce, 1888, Ann. Mag. nat. Hist. (6) 2: 241, by original designation. Honduras: [Roatan Island] Ruatan Island.

‡BOMBYLIODES; misspelling. See Gymnelia Walker, 1854.

‡BOMBYSATUR Corporaal, 1921, De Thee 2: 114. NOTO Both the generic name and the only included specific name, ‡Bombysatur corporaali, are nomina nuda. Both names were attributed by Corporaal to Van Eecke, but no published reference by Van Eecke to these names has been found. Neither Dr R. de Jong nor Dr A. Diakonoff, both of the Rijksmuseum, Leiden, found any indication that Van Eecke ever published the names ‡Bombysatur and ‡corporaali (personal communication to D.S. Fletcher on 28.iii.1978).

‡BOMBISATUR; Bernard, 1921, De Thee 2: 115.

An incorrect subsequent spelling of ‡Bombysatur Corporaal, 1921.

BORECONIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 459.

Type-species: *Boreconia subviridis* Walker, 1865, *ibidem* 32: 460, by monotypy. South Africa.

Boreconia was established in the Notodontidae; it was placed in the Lymantriadae, now Lymantriidae, by Hampson,

1905, Ann. S. Afr. Mus. 3: 397.

‡BARECONIA; Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 298.

An incorrect subsequent spelling of *Boreconia* Walker, 1865.

BORIZA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 261 (key), 310.

Type-species: *Heterocampa crossoea* Druce, 1894, *Ann. Mag. nat. Hist.* (6) **13**: 357, by original designation. MEXICO: Orizaba.

BORSEBA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 318.

Type-species: *Borseba surgens* Walker, [1865] 1864, *ibidem* 31: 318, by monotypy. [COLOMBIA]: Bogota.

BOSCAWENIA Fletcher nom. n. for Empodisma Kiriakoff, 1964.

Type-species: Stenostaura polioplaga Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 472, by original designation (for Empodisma Kiriakoff, 1964). NIGERIA: southern, Arochuku.

Prof. Dr S.G. Kiriakoff was informed of the homonymy and has suggested this replacement name.

EMPODISMA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 34.

Type-species: Stenostaura polioplaga Hampson, 1910, by original designation.

A junior homonym of *Empodisma* Yamaguti, 1963, *Syst. helminthum* 5: 114, – Vermes, Acanthocephala. The objective replacement name is *Boscawenia* Fletcher nom. n.

BOSTRYCHOGYNA Kiriakoff, 1960, Revue Zool. Bot. afr. 61: 50.

Type-species: *Peratodonta bella* Bethune-Baker, 1913, *Ann. Mag. nat. Hist.* (8) **12**: 63, by original designation. NIGERIA: Oni, 70 miles E. of Lagos.

BRACHAROA Hampson, 1905, Ann. S. Afr. Mus. 3: 392.

Type-species: Orgyia quadripunctata Wallengren, 1875, Ofvers. K. Vetensk Akad. Förh. Stockh. 32 (1): 99, by original designation. South Africa: Transvaal.

BRACHARTHRON Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 25, 99. NOCTUIDAE

Type-species: Bracharthron maculapex Hampson, 1891, ibidem 8: 25, 99, pl.148, figs 4, 11, by monotypy.

Further information in Nye, 1975, Generic Names Moths World 1: 86.

BRACHARTHRUM Hampson, 1895, Fauna Br. India (Moths) 3: 50.

An unjustified emendation of *Bracharthron* Hampson, 1891. *Bracharthrum* in the Noctuidae, Hypeninae is cited here as it was not included in Nye, 1975, *Generic Names Moths World* 1.

BRACHARTHRUM Hampson, 1895. See Bracharthron Hampson, 1891. NOCTUIDAE

BRACHIONYCOIDES Marumo, 1920. See Shaka Matsumura, 1920. NOTO

BRACHIOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 89, 334.

Type-species: *Brachiosia castaneola* Hampson, 1900, *ibidem* 2: 335, fig.235, by original designation. [INDONESIA]: [Sangihe I.] Sangir.

BRACHYCHIRA Aurivillius, 1905, Ark. Zool. 2 (12): 28.
NOTO

Type-species: *Brachychira ferruginea* Aurivillius, 1905, *ibidem* 2 (12): 28, text-fig.4, pl.4, fig.6, by monotypy. [NIGERIA]: Niger.

BRACHYGLENE Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 99, 100.

Type-species: *Phaeochlaena bracteola* Geyer, 1832, *in* Hübner, *Zuträge Samml. exot. Schmett.* **4**: 17, pl.[113], figs 655, 656, by monotypy. SOUTH AMERICA.

Brachyglene was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855;

the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig

bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

Brachyglene was established in the 'Lithosina', now Arctiidae; it was placed in the Cyllopodidae, now Geometridae Sterrhinae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 407; and was placed in the Dioptidae by Prout, 1918, Novit.

zool. 25: 418.

BRADYTERA Toxopeus, 1948, Treubia 19: 478.

Type-species: Euproctis lodra Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 349, pl.9A, fig.6, by original designation. Java.

BRANDTAXIA Boursin, 1943, Z. wien. ent. Ges. 28: 314.
NOCTUIDAE

Type-species: Sidemia discalis Brandt, 1941, Mitt. münch. ent. Ges. 31: 850, pl.26, fig.37, by original designation. Type(s) 8, IRAN: Fort Sengan.

Brandtaxia in the Noctuidae, Acronictinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

BRAURA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 464.

LASIOCAMPIDAE

Type-species: Braura ligniclusa Walker, 1865, ibidem 32:

464, by monotypy. [South Africa]: Natal.

Braura was established in the Notodontidae; it was included in the Lasiocampidae by Aurivillius, 1927, in Seitz, Gross-Schmett. Erde 14: 246.

BRECONTIA Dyar, 1908, Proc. ent. Soc. Wash. 9: 49. NOTO Type-species: Brecontia plagipennis Dyar, 1908, ibidem 9: 49, by original designation (in accordance with the Code, Article 68(a)(i)). PERU: Callao.

BREYERIA Janse, 1920. See Eubreyeria Fletcher nom. n. NOTO

BRIACHISIA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 140.

Type-species: Stauropus briachisia Schaus, 1928, Proc. U.S. natn. Mus. 73 (19): 75, by original designation. Philippines: Mindanao, Surigao.

BRUCEIA Neumoegen, 1893, Jl N.Y. ent. Soc. 1: 35.
Type-species: Bruceia pulverina Neumoegen, 1893, ibidem
1: 36, by monotypy. U.S.A.: western Colorado.

‡BRUCIA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 421.

An incorrect subsequent spelling of *Bruceia* Neumoegen, 1893.

‡BRUCIA; misspelling. See Bruceia Neumoegen, 1893.

BRUNIA Moore, 1878, Proc. zool. Soc. Lond. 1878: 15. LITH Type-species: Lithosia antica Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 505, by PRESENT DESIGNATION. [SRI LANKA] Ceylon.

BRYANTIA Schaus, 1922, Insecutor Inscit. menstr. 10: 24.

Type-species: *Bryantia caudata* Schaus, 1922, *ibidem* 10: 25, by original designation. JAVA: Pelaboean.

BRYCEA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 563.

Type-species: *Brycea disjuncta* Walker, 1854, *ibidem* 2: 563, by monotypy. BRAZIL.

BRYKIA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 644.

Type-species: *Allata albonotata* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) **6**: 140, by monotypy. [BORNEO]: Sarawak.

A. albonotata is a junior subjective synonym of Eumeta horsfieldi Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 430. The title of volume 1 of the work, [1858] 1857 is Cat. lepid. Insects Mus. Hon. East-India Company

The date of publication of the part of Seitz containing page 644 is cited on page 641 as 8.iv.1930 in both the German and

the English editions.

BUCAEA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1983.

Type-species: Savara simplex Walker, [1865] 1864, ibidem 31: 320, by monotypy (of Savara Walker, [1865]). [INDIA]: Hindostan.

Bucaea was established as the objective replacement name for Savara Walker, [1865].

SAVARA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 320.

Type-species: Savara simplex Walker, [1865], by monotypy. A junior homonym of Savara Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 184, – Lepid., Noctuidae. The objective replacement name is Bucaea Walker, 1866.

BUGANDITA Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 316.

Type-species: *Bugandita bisignata* Kiriakoff, 1965, *ibidem* 101: 317, fig.6, by monotypy. UGANDA: Bwamba.

‡BUGISARA; misspelling. See Caroia Walker, 1858. NOTO

BUMETOPON Aurivillius, 1920, Ark. Zool. 13 (2): 30. GEOMETRIDAE

Type-species: *Bumetopon bicorne* Aurivillius, 1920, *ibidem* 13 (2): 30, pl.1, fig.5, by monotypy. AUSTRALIA: [Western Australia] north-west, Broome.

Bumetopon was established in the Notodontidae; it was transferred to the Geometridae, Oenochrominae by Prout, 1934, in Seitz, Gross-Schmett. Erde 12: 135.

BURTIA Grote, 1866, Proc. ent. Soc. Philad. 6: 185. CTEN Type-species: Burtia rubella Grote, 1866 July, ibidem 6: 186, pl.5, fig.1, by monotypy. Cuba.

GUNDLACHIA Herrich-Schäffer, 1866, CorrespBl. zool.-min. Ver. Regensburg 20: 107, 108.

Type-species: Gundlachia cruenta Herrich-Schäffer, 1866 July, ibidem 20: 108, by monotypy. CUBA.

G. cruenta is a junior subjective synonym of Burtia rubella Grote, 1866, Proc. ent. Soc. Philad. 6: 186. Grote, 1867, ibidem 6: 309, as first reviser, chose the name rubella for the taxon for which the names rubella and cruenta were published in the same month.

A junior homonym of *Gundlachia* Pfeiffer, 1850, *Z. Malakozool.* **1849**: 97, – Mollusca. There is no objective replacement name but *G. cruenta* is a junior subjective synonym of *Burtia rubella* Grote, 1866 (see above) the type-species of *Burtia* Grote, 1866. The latter is therefore available for use as a subjective replacement name.

‡BUTHISIA; misspelling. See Buthysia Wallengren, 1863.

BUTHYSIA Wallengren, 1863, Wien. ent. Monatschr. 7: 139.

Type-species: Buthysia sangaris Wallengren, 1863, ibidem

7: 139, by monotypy (but attributed to Dalman, an incorrect

authorship). JAVA.

B. sangaris is a junior subjective synonym of Syntomis huebneri Boisduval, [1828], Essai Monogr. Zygénides: 127, pl.8, fig.4 (but cited as ‡hubneri [without umlaut], an incorrect original spelling).

‡BUTHISIA; Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1866.

An incorrect subsequent spelling of *Buthysia* Wallengren, 1863.

BYRDIA Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 536 (key), 537.

Type-species: Laria rossii Curtis, 1835, in Ross, Narrative of a second voyage in search of a N.W. Passage (Appendix): lxx, pl.A, fig.10, by original designation. [Canada]: [N.W. Territories], Fury Beach.

The date of publication of the part of Seitz containing page 536 is cited on page 529 as 17.v.1927 in both the German and

the English editions.

See also Konokareha Matsumura, [1928].

BYRSIA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 193.

Type-species: *Byrsia dotata* Walker, [1865] 1864, *ibidem* 31: 193, by monotypy. [Indonesia]: Timor [I.] and [Moluccas], [Batjan I.] Batchian.

BYSSOPHAGA Stretch, 1872, Illust. Zygaenidae Bombycidae N. Am. 1: 48.

Type-species: *Lithosia nexa* Boisduval, 1869, *Annls Soc. ent. Belg.* 12: 74, by subsequent designation by Grote, 1874, *Bull. Buffalo Soc. nat. Sci.* 2: 151. [U.S.A.]: California.

CABARDA Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 435.

Type-species: *Cabarda molliculana* Walker, 1863, *ibidem* **28**: 435, by monotypy. Borneo: Sarawak.

‡CACHARA; misspelling. See Caschara Walker, 1862.

CACOETHES Hübner, [1819] 1816, Verz. bekannter Schmett.: 170. CTEN

Type-species: Sphinx polidamon Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 75, pl.237, fig.B, by monotypy. [SOUTH AFRICA]: [Cape of Good Hope] Kaap de Goede Hoop.

S. polidamon is dated from the wrapper of the part and not from the title-page of the volume, 1782.

CACOLYCES Warren, 1906, Proc. U.S. natn Mus. 30: 409.

Type-species: Lyces plagifera Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1656, by original designation. [BRAZIL]: Valley of the Amazon.

Cacolyces was established in the Cyllopodinae, now Geometridae Sterrhinae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 404.

CACOSOMA Felder, 1874, in Felder & Rogenhoffer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.13. CTEN

Type-species: Cacosoma naclioides Felder, 1874. ibidem 2 (Abt.2): pl.102, fig.13, by monotypy. [SOUTH AFRICA]: Port Natal, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 2.

C. naclioides is a junior subjective synonym of Naclia gnatula Boisduval, 1847, in Delegorgue, Voyage Afr. australe 2: 596.

CACOSTATIA Hampson, 1898, Cat. Lepid. Phalaenae Br.
Mus. 1: xviii, 26, 430.

Typc-species: Agyrta saphira Staudinger, 1876, Verh. zool.-

bot. Ges. Wien 25 (Abh.): 122, by original designation (but cited as sapphira, an unjustified emendation by Hampson, 1898). [PANAMA]: Chiriqui.

CADORELA Griveaud, 1973, Bull. Soc. ent. Fr. 78: 136.

Type-species: Cadorela translucida Griveaud, 1973, ibidem 78: 137, figs 11, 12, by original designation. MADAGASCAR: [W.], E. de Sakaraha, réserve spéciale du Zombitsy, Matsabory.

CADRUSIA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S.Atkinson (1): 54.

Type-species: *Cadrusia virescens* Moore, 1879, *ibidem* (1): 54, pl.3, fig.16, by monotypy. INDIA: [W. Bengal, Darjeeling] Darjiling.

CADURCA Swinhoe, 1906, Ann. Mag. nat. Hist. (7) 17: 546.

Type-species: *Cadurca venata* Swinhoe, 1906, *ibidem* (7) **17**: 546, by monotypy. [ANGOLA]: Bihé.

CAELICOLA Dognin, 1923, Hétérocères nouv. Am. Sud 22: 4.

Type-species: Caelicola rostrata Dognin, 1923, ibidem 22: 4, by original designation. [BRAZIL]: Amazones, São Paulo de Olivença.

CAENERESSA Obraztsov, 1957, Bull. Mus. comp. Zool. Harv. 116: 392.

Type-species: *Syntomis diaphana* Kollar, [1844] 1848, *in* Hügel, *Kaschmir und das Reich der Siek* 4 (2): 460, pl.19, fig.7, by PRESENT DESIGNATION. [INDIA]: [Mussooree] Massuri.

CAENINA Felder, 1861, Sber. Akad. Wiss. Wien 43 (Abt.1): 33.

Type-species: Caenina lymantrioides Felder, 1861, ibidem 43 (Abt.1): 34, by monotypy. [Indonesia]: [Moluccas], [Ambon I.] Amboina.

‡CAENOSTEGIA; misspelling. See Coenostegia Mabille, 1891.

CALAMIDIA Butler, 1877, Trans. ent. Soc. Lond. 1877: 358.

Type-species: *Lithosia hirta* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 2: 510, by original designation. [AUSTRALIA]: Sydney.

‡CALANOTOS; misspelling. See Calonotos Hübner, [1819].

CALAPANA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 214.

Type-species: *Dediama basivacua* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 134, by original designation. Borneo: Sarawak.

CALEPIDOS Boisduval, 1870, Considérations Lépid. Guatemala: 89. PERI

Type-species: *Calepidos celina* Boisduval, 1870, *ibidem*: 89, by subsequent designation by Bryk, 1931, *Lepid. Cat.* 45: 18. GUATEMALA.

C. celina is a junior subjective synonym of Phalaena pylotis Drury, 1773, Illust. nat. Hist. exot. Insects 2: 9, pl.5, fig.3, index.

CALIDOTA Dyar, 1901, Proc. U.S. natn. Mus. 23: 268. ARCT Type-species: Arctia strigosa Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 615, by monotypy. Type-LOCALITY: not known, [AMERICA]: [Central].

‡CALLIDOTA; Neave, 1939, Nomencl. zool. 1: 532.

An incorrect subsequent spelling, by Neave who attributed it to Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 432.

‡CALIGENIA; misspelling. See Calligenia Duponchel, [1845].

CALIGULA Aurivillius, 1879. See Paralacydes Aurivillius, 1889.

CALLAGRA Butler, 1876. See Euagra Walker, 1854. CTEN

CALLALUCIA Grote, 1865. See Omoiala Grote, 1863. PERI

CALLARCTIA Packard, 1864. See Arctia Schrank, 1802.

CALLARCTIA Leech, 1899. See Euleechia Dyar, 1900.

CALLEDEMA Butler, 1878, Trans. ent. Soc. Lond. 1878: 65.

Type-species: Calledema marmorea Butler, 1878, ibidem 1878: 65, by original designation. [BRAZIL]: Amazon region, Lake Cararaucu.

CALLIA Walker, 1855 [December 8]. See Porela Walker, 1855 [June 9].

CALLICARUS Grote, 1866, Proc. ent. Soc. Philad. 6: 182.

Type-species: *Callicarus pennipes* Grote, 1866, *ibidem* 6: 182, pl.5, fig.3, by subsequent designation by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 417. CUBA.

‡CALLICORUS; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 129.

An incorrect subsequent spelling of Callicarus Grote, 1866.

CALLICEREON Butler, 1882, Cistula ent. 3: 2. NOCTUIDAE Type-species: Callicereon affine Butler, 1882, ibidem 3: 3, by monotypy. Type(s) & MADAGASCAR: Betsileo Country, [Ankafina forest] Ankafana, (BMNH).

C. affine is a junior subjective synonym of Deiopeia heterochroa Mabille, 1879, Bull. Soc. zool. Fr. 3: 88 (but proposed

in ‡Dejopeia, an incorrect subsequent spelling).

Callicereon was established in the Arctiidae; it was placed in the Noctuidae by Viette, 1965, Bull. Soc. ent. Fr. 70: 151.

Callicereon in the Noctuidae, Acronictinae was not included in Nye, 1975, Generic Names Moths World 1.

‡CALLICORUS; misspelling. See Callicarus Grote, 1866.

CALLIDESMA Agassiz, 1847. See Calodesma Hübner, [1820].

‡CALLIDOTA; misspelling. See Calidota Dyar, 1901.

‡CALLIGENEA; misspelling. See Calligenia Duponchel, [1845].

CALLIGENIA Duponchel, [1845], Cat. méth. Lépid. Eur.: 59.

Type-species: *Bombyx rosea* Fabricius, 1775, *Syst. Ent.*: 587, by monotypy. Europe.

B. rosea Fabricius is a junior subjective synonym of Phalaena miniata Forster, 1771, Nova Species Insectorum: 75.

See also Miltochrista Hübner, [1819].

‡CALLIGENEA; Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 575.

An incorrect subsequent spelling of *Calligenia* Duponchel, [1845].

‡CALIGENIA; Staudinger & Rebel, 1901, Cat. Lepid. palaearct. Faunengeb. 1: 374.

An incorrect subsequent spelling of *Calligenia* Duponchel, [1845].

CALLIMORPHA Latreille, 1809, Genera Crust. Insect. 4: 220.

ARCT

Type-species: *Phalaena dominula* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 509, by subsequent designation by Desmarest, 1857, *in* Chenu, *Encycl. Hist. nat.* (Papillons nocturnes): 33. Type-LOCALITY: not stated, [EUROPE].

The type-species designations of Latreille, 1810, Considérations générales sur l'Ordre naturel des Animaux: 365, were accepted by the International Commission on Zoological Nomenclature, 1939, Opin. Decl. int. Commn zool. Nom. 2 (Opinion 136): 15, for those genera in which Latreille in his 'Table des genres avec l'indication de l'espèce qui leur sert de type' cited only one of the species included in the genus by the original author. In the case of Callimorpha four such species were cited.

Curtis, 1834, Br. Ent. 11: 499, and Westwood, 1840, Introd. mod. Classif. Ins. 2 (Synopsis Genera Br. Ins.): 92, both designated Phalaena jacobaeae Linnaeus, 1758, a nominal species not originally included in Callimorpha and therefore not eligible to be the type-species.

PANAXIA Tams, 1939, Entomologist 72: 73.

Type-species: *Phalaena dominula* Linnaeus, 1758, by original designation.

A junior objective synonym of Callimorpha Latreille, 1809.

CALLINDRA Röber, 1925, Stettin. ent. Ztg 86 (1): 172.

Type-species: Callindra gigantea Röber, 1925, ibidem 86 (1): 171, by monotypy. [INDIA]: Naga Hills.

C. gigantea Röber is a junior subjective synonym of Areas arginalis Hampson, 1894, Fauna Br. India (Moths) 2: 26.

CALLINOTUS Agassiz, 1847. See Calonotos Hübner, [1819].

CALLISTHENIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 89, 362.

Type-species: *Maepha plicata* Butler, 1877, *Trans. ent. Soc. Lond.* 1877: 328, by original designation. [BRAZIL]: [Espirito Santo] Espiritu Sancto.

CALLITEARA Butler, 1881, Trans. ent. Soc. Lond. 1881: 12.

Type-species: *Dasychira argentata* Butler, 1881, *ibidem* 1881: 12, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 470. Japan: Yokohama.

CALLITOMIS Butler, 1876, J. Linn. Soc. (Zool.) 12: 351.

Type-species: *Callitomis syntomoides* Butler, 1876, *ibidem* 12: 351, by original designation. [INDIA/ PAKISTAN]: [Kashmir] Cashmere.

‡CALLITORNIS; Neave, 1939, Nomencl. zool. 1: 540.

An incorrect subsequent spelling by Neave who incorrectly attributed it to Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 33.

‡CALLITORNIS; misspelling. See Callitomis Butler, 1876.

CALLOBALACRA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 70.

Type-species: *Metarctia rubrostriata* Aurivillius, 1898, *Ent. Tidskr.* 19: 185, by original designation. Togo.

Callobalacra was established as a subgenus of Balacra Walker, 1856.

CALLOPEPLA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xix, 29, 437.

Type-species: *Charidea inachia* Schaus, 1892 August, *Proc. zool. Soc. Lond.* **1892**: 277, by original designation. BRAZIL: Petropolis.

C. inachia was redescribed by Schaus, 1892 November, American Lepid.: 6, pl.1, fig.19.

CALLOPIS Billberg, 1820. See Spiris Hübner, [1819].

CALODESMA Hübner, [1820] 1816, Verz. bekannter Schmett.: 177. PERI

Type-species: *Phalaena amica* Stoll, 1781, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* **4**: 158, pl.370, fig.H, by monotypy. SURINAM.

P. amica is dated from the wrapper of the part and not from the title-page of the volume, 1782.

See also Leucopsumis Hübner, [1820]

CALLIDESMA Agassiz, 1847, Nomencl. zool. (Index univl.): 58, 60.

An unjustified emendation of Calodesnia Hübner, [1820]

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Callidesma* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

PYRODESMA Boisduval, 1870, Considérations Lépid. Guatemala: 85.

Type-species: *Phalaena amica* Stoll, 1781, by PRESENT DESIGNATION.

A junior objective synonym of Calodesma Hübner, [1820].

CALOMATHETES Dyar, 1913, Proc. U.S. natn. Mus. 44: 303.

Type-species: Calomathetes halmaphylla Dyar, 1913, ibideni 44: 303, by original designation. MEXICO: Tehuacan.

‡CALONOTA; misspelling. See Calonotos Hübner, [1819].

CALONOTOS Hübner, [1819] 1816, Verz. bekannter Schmett.: 123.

Type-species: *Sphinx phlegmon* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 57, pl.35, fig.H, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 126. Surinam.

S. phlegmon is dated from the wrapper of the part and not from the title-page of the volume, 1779.

CALLINOTUS Agassiz, 1847, Nomencl. zool. (Index univl.): 59, 60.

An unjustified emendation of *Calonotos* Hübner, [1819] (but cited by Agassiz as ‡*Calonotus* an incorrect subsequent spelling).

A junior homonym of *Callinotus* Schoenherr, 1825, *Isis* (*Oken*) 1825: 584, – Insecta, Coleoptera. The objective replacement name is *Calonotos* Hübner, [1819].

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Callinotus* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

‡CALONOTUS; Agassiz, 1847, Nomencl. zool. (Index univl.): 59, 60.

An incorrect subsequent spelling of *Calonotos* Hübner, [1819].

‡CALONOTA; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1627.

An incorrect subsequent spelling of *Calonotos* Hübner, [1819].

‡CALANOTOS; Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 283, 551.

An incorrect subsequent spelling of *Calonotos* Hübner, [1819].

‡CALONOTUS; misspelling. See Calonotos Hübner, [1819].

CALOSOMA Geyer, 1832. See Osmacola Watson nom. n.

CALPENIA Moore, 1872, Proc. zool. Soc. Lond. 1872: 571.

Type-species: *Calpenia saundersi* Moore, 1872, *ibidem* **1872**: 571, pl.33, fig.1, by monotypy. INDIA.

CALPOPARIA Watson nom. n. for Paraplocia Aurivillius, 1909.

Type-species: *Lithosia imparepunctata* Oberthür, 1890, *Etudes d'Entomologie* 13: 15, pl.4, figs 19, 20, by original designation (for *Paraplocia* Aurivillius, 1909). Comoro ISLANDS: [Indian Ocean], Anjouan [I.].

PARAPLOCIA Aurivillius, 1909, in Voeltzkow, Wiss. Ergebn. Reise Ostafr. 2: 336.

Type-species: *Lithosia imparepunctata* Oberthür, 1890, by original designation.

A junior homonym of *Paraplocia* Pomel, [1872], *Paléontologie Prov. Oran.* (Zoophytes) 5: 104, – Porifera. The objective replacement name is *Calpoparia* Watson nom. n.

CALTURA Moore, 1879, Proc. zool. Soc. Lond. 1879: 401.

Type-species: *Caltura alba* Moore, 1879, *ibideni* **1878**: 401, by monotypy. [Sri Lanka] Ceylon.

CALYBIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 446.

Type-species: *Phryne immaculata* Grote, 1865, *Proc. ent. Soc. Philad.* **5**: 246, by monotypy (of *Phryne* Grote, 1865). CUBA.

Calybia was established as the objective replacement name for *Phryne* Grote, 1865.

Calybia was established in the Liparidae, now Lymantriidae; it was transferred to the Eucleidae, now Limacodidae, by Dyar, 1897, Jl N.Y. ent. Soc. 5: 121.

PHRYNE Grote, 1865, Proc. ent. Soc. Philad. 5: 246.

Type-species: Phryne immaculata Grote, 1865, by monotypy. A junior homonym of Phryne Fitzinger, 1843, Systema Reptilium (Fasc. Primus): 32, — Amphibia. The objective replacement name is Calybia Kirby, 1892. Further details of nomenclaturally unavailable usages of Phryne before 1843, and another usage in the Lepidoptera after 1843, will be given in Generic Names Moths World 4.

Phryne Grote was established in the Dasychirae, now Lymantriidae; it was transferred to the Eucleidae, now Limacodidae, by Dyar, 1897, Jl N.Y. ent. Soc. 5: 121.

CALYPTRONOTUM Roepke, 1944, Natuurh. Maandbl. 33:

Type-species: Calyptronotum confusum Roepke, 1944, ibidem 33: 5, figs 19a, 19b, by monotypy. [Java]: G. Gedeh.

CAMPTOLOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.93, fig.7. ARCT

Type-species: Camptoloma erythropygum Felder, 1874, ibidem 2 (Abt.2): pl.93, fig.7, by monotypy. [CHINA]: Shanghai, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 2.

C. erythropygum is a junior subjective synonym of Numenes interiorata Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 290.

‡CAMPYLOCTIS; misspelling. See Campyloctys Janse, 1920. NOTO

CAMPYLOCTYS Janse, 1920, Ann. Transv. Mus. 7: 219.

Type-species: Campyloctys gladstonei Janse, 1920, ibidem 7: 220, pl.5, fig.20, pl.10, figs 1-7, pl.14, fig.13, by original designation. South Africa: Capetown.

‡CAMPYLOCTIS; Gaede, 1934, in Strand, Lepid. Cat. 59:

An incorrect subsequent spelling of Campyloctys Janse, 1920.

CAMPYLONA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 658.

Type-species: Campylona bicolor Möschler, 1878, ibidem 27 (Abh.): 659, pl.9, fig.27, by monotypy. Surinam: interior. Campylona was established in the 'Lithosidae', now Arctiidae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 398.

CANDIDATA Toxopeus, 1948, Treubia 19: 452 (key), 453.

Type-species: Leucoma subargentea Felder, 1861, Sber. Akad. Wiss. Wien 43 (Abt.1): 32, by original designation. [Indonesia]: [Moluccas], [Ambon I.] Amboina.

CANODIA Guenée, 1852, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 6: 377.

Type-species: Canodia carmelitoides Guenée, 1852, ibidem

6: 378, by monotypy. Brazil: Nova Friburgo.

Canodia was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 343.

CANOPUS Walker, 1855. See Amerila Walker, 1855. ARCT

CAPISSA Moore, 1878, Proc. zool. Soc. Lond. 1878: 19.

Type-species: Lithosia vagesa Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 304, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 130. INDIA: Darjeeling.

CAPRIMA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 128.

Type-species: Caprima gelida Walker, [1865] 1864, ibidem 31: 129, by monotypy. New Guinea: [Irian Jaya], [Misool I.] Mysol.

Caprima has been used as the original generic name for species now placed in the Arctiidae.

CAPRIMIMA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 561.

Type-species: Hypocrita albicollis Pagenstecher, 1886, Jb. nassau. Ver. Naturk. 39: 125, by original designation by Hampson on page 286 but cited as 'Caprima gelida Walker' (see below). [Indonesia]: [Moluccas], Aru Islands, New Guinea: south-western.

Hampson on page 286 used Caprima Walker, [1865] 1864; cited C. gelida Walker, [1865] 1864 as type-species; and included Hypocrita albicollis Pagenstecher, 1886, as a junior subjective synonym of gelida. In the Addenda on page 561, Hampson stated that C. gelida was a zygaenid; that the valid name of the arctiid species should be albicollis; and established a new genus Caprimima for the concept for which he had incorrectly used Caprima on page 286.

H. albicollis is a junior subjective synonym of Damias elegans

Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 259, fig. See also Damias Boisduval, 1832.

CAPRIMIMODES Rothschild, 1913, Novit. zool. 20: 220.

Type-species: Caprimimodes mimetica Rothschild, 1913, ibidem 20: 220, by monotypy. New Guinea: [Irian Jaya], Snow Mts, near Oetakwa River.

CARAGOLA Moore, 1879. See Caragolina Strand, 1928.

CARAGOLINA Strand, 1928, Arch. Naturgesch. 92 (A) 8: 50.

Type-species: Caragola costalis Moore, 1879, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 46, pl.2, fig.21, by monotypy (of Caragola Moore, 1879). INDIA: [W. Bengal, Darjeeling Darjiling.

Caragolina was established as the objective replacement name for Caragola Moore, 1879.

CARAGOLA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 46.

Type-species: Caragola costalis Moore, 1879, by monotypy. A junior homonym of Caragola Gray, 1851, Proc. zool. Soc. Lond. 19: 235, 239, – Pisces. The objective replacement name is Caragolina Strand, 1928.

CARALES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 729.

Type-species: Carales albicans Walker, 1855, ibidem 3: 730, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 207, but cited as Phalaena astur Cramer, 1777, a nominal species not original included in Carales. Kirby, however, placed astur as the senior synonym of albicans, a nominal species originally included in Carales. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. VENEZUELA.

CARALISA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1660.

Type-species: Caralisa editha Walker, 1856, ibidem 7: 1661, by monotypy. [HISPANIOLA] [Haiti] Hayti: [Dominican Republic], S[an]t[o] Domingo.

CARANUSCA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6:

Type-species: Caranusca rubrifera Walker, 1862, ibidem 6: 101, by monotypy. [Borneo]: Sarawak.

CARATHIS Grote, 1865, Proc. ent. Soc. Philad. 5: 253.

Type-species: Carathis gortynoides Grote, 1865, ibidem 5: 253, pl.4, fig.8, by monotypy. CUBA.

CARBISA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 41.

Type-species: Carbisa venosa Moore, 1879, ibidem (1): 41, pl.2, fig.10, by monotypy. INDIA: [Darjeeling] Darjiling.

CARCINARCTIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xvi, 7, 472.

Type-species: Carcinarctia metamelaena Hampson, 1901, ibidem 3: 472, fig. 206, by original designation. [Kenya] Br. E. Afrika: Lari.

CARCINOPODIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 86, 249.

Type-species: Gnophria furcifasciata Butler, 1895, Proc.

zool, Soc. Lond. 1895: 268, by original designation. [MALAWI]: Fwambo.

‡CARSINOPODIA; Neave, 1939, Nomencl. zool. 1: 592.

An incorrect subsequent spelling of Carcinopodia Hampson, 1900.

CARCINOPYGA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.101, fig.3. ARCT

Type-species: Carcinopyga lichenigera Felder, 1874, ibidem 2 (Abt.2): pl. 101, fig.3, by monotypy. [INDIA/PAKISTAN]: [Kashmir], Ladak [h], cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 2.

CARGIDA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 258 (key), 267.

Type-species: *Heterocampa pyrrha* Druce, 1898, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 2: 459, pl.91, figs 4, 5, by original designation. MEXICO: Presidio de Mazatlan, Jalisco State.

CARIDARCTIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xvi, 7, 452.

Type-species: Arctia albicancellata Burmeister, 1878, Descr. phys. Répub. Argent. 5: 449, by original designation. ARGENTINA: Buenos Aires.

Caridarctia was established in the 'Arctiadae'; it was transferred to a new subfamily Xanthopastinae, in the Noctuidae, by Biezanko, 1971, in Biezanko & Guerra, Archos Mus. nac. Rio de J. 54: 270. Caridarctia is not closely related to Xanthopastis Hübner, [1821], so is here retained in the Arctiinae pending a detailed study by our colleagues V.O. Becker and M.R. Honey.

CARILEPHIA Schaus, 1924, Proc. U.S. natn. Mus. 65 (7): 24.

Type-species: Carilephia moninna Schaus, 1924, ibidem 65 (7): 24, by original designation. Philippines: Luzon, Mt Makiling.

CARIPODIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 92, 248.

Type-species: Caripodia chrysargyria Hampson, 1900, ibidem 2: 248, fig.162, by original designation. [Rhodesia]: Mashonaland, Salisbury.

‡CHARIPODIA; Grünberg, 1910, Arch. Naturgesch. 74 (2) 2: 228.

An incorrect subsequent spelling of *Caripodia* Hampson, 1900.

CAROIA Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1861.

Type-species: Bagisara bombycoides Walker, 1858, ibidem 15: 1792, by monotypy (of Bagisara Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1791). Brazil.

C. bombycoides is a junior subjective synonym of Phalaena licormas Cramer, 1775, Uitlandsche Kapellen (Papillons exot.) 1: 117, pl.74, fig.E.

Caroia was established in the Noctuidae as the objective replacement name for Bagisara Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1791; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 342.

BAGISARA Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1791.

Type-species: Bagisara bombycoides Walker, 1858, by monotypy.

A junior homonym of *Bagisara* Walker, 1858, *ibidem* **15**: 1770, – Lepid., Noctuidae. The objective replacement name is *Caroia* Walker, 1858.

‡BUGISARA; Neave, 1939, Nomencl. zool. 1: 498.

An incorrect subsequent spelling of *Bagisara* Walker, 1858, *List Specimens lepid. Insects Colln Br. Mus.* **15**: 1791, mistakenly attributed to Butler by Neave.

CARPENTERELLA Collenette, 1933, Entomologist's mon. Mag. 69: 258.

Type-species: Carpenterella nesiotica Collenette, 1933, ibidem 69: 259, by original designation. UGANDA: Lake Victoria, Sesse Islands, Bugalla Island.

CARPOSTALAGMA Karsch, 1894, Ent. Nachr. Berlin 20: 379.

Type-species: *Caryatis viridis* Plötz, 1880, *Stettin. ent. Ztg* 41: 80, by monotypy. [NIGERIA] West Africa: Victoria.

CARRIOLA Swinhoe, 1922, Ann. Mag. nat. Hist. (9) 10: 468.

Type-species: *Leucoma ecnomoda* Swinhoe, 1907, *ibidem* (7) **20**: 77, by original designation. JAVA: [Bogor] Buitenzorg.

‡CARSINOPODIA; misspelling. See Carcinopodia Hampson, 1900.

CARYATIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 171.

Type-species: *Phalaena phileta* Drury, 1782, *Illust. nat. Hist. exot. Insects* 3: index, 30, pl.22, fig.5, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 391. SIERRA LEONE.

ASOTOMORPHA Hulstaert, 1923, Revue zool. afr. 11: 410. Type-species: Phalaena phileta Drury, 1782, by original designation

Asotomorpha is a junior objective synonym of Caryatis Hübner, [1819].

CASAMA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 611.

Type-species: Casama indeterminata Walker, 1865, ibidem 32: 611, by monotypy. [INDIA]: South Hindostan.

C. indeterminata is a junior subjective synonym of Euproctis vilis Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 348.

Casama was established in the Bryophilidae, now Noctuidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 466.

BEERIA Hartig, 1963, in Hartig & Prola, Mitt. münch. ent. Ges. 53: 64.

Type-species: Ocneria flavipalpata Staudinger, 1895, Dt. ent. Z. Iris 8: 296, pl.5, fig.5, by monotypy. [JORDAN]: [R.] Jordan valley

O. flavipalpata is a junior subjective synonym of Spilosoma innotata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 674.

A junior homonym of *Beeria* Match, 1957, *Univ. Wash. Publs Biol.* 16 (2): 231, – Insecta, Coleoptera. There is no objective replacement name but *O. flavipalpata* is congeneric with *Casama indeterminata* Walker, 1865, the type-species of *Casama* Walker, 1865. The latter is therefore available for use as a subjective replacement name.

CASCERA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 460.

Type-species: Cascera muscosa Walker, 1865, ibidem 32: 461, by monotypy. [AUSTRALIA]: [Western Australia], Swan River.

Turner, 1903, *Proc. Linn. Soc. N.S.W.* **28**: 73, 74 recorded *C. muscosa* from Queensland and stated that the locality given by Walker was 'certainly erroneous'.

CASCHARA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 133.

Type-species: Caschara punctifera Walker, 1862, ibidem 6: 133, by monotypy. BORNEO: Sarawak.

‡CACHARA; Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 236.

An incorrect subsequent spelling of *Caschara* Walker, 1862. The error was incorrectly attributed to Roepke, 1943, by Kiriakoff.

CASSIDIA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 129.

Type-species: Cassidia obtusa Walker, 1862, ibidem 6: 130, by monotypy. Borneo: Sarawak.

CASTABALA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 270.

Type-species: *Castabala roseata* Walker, [1865] 1864, *ibidem* 31: 271, by monotypy. [INDIA]: Hindostan.

‡CASTABELA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 422.

An incorrect subsequent spelling of *Castabala* Walker, [1865].

‡CASTABELA; misspelling. See Castabala Walker, [1865].

CASTASTYGNE Kiriakoff, 1962, Revue Zool. Bot. afr. 66: 18.

Type-species: Antheua tristicolor Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 433, pl.71 row h, by original designation. [ZAMBIA] Nord-Rhodesia.

The date of publication of the part of Seitz containing page 433 is cited on that page as 30.v.1928 in the German edition; and as 26.xi.1928 in the English edition.

CASTRICA Schaus, 1896, Jl N.Y. ent. Soc. 4: 136. ARCT Type-species: Castrica oweni Schaus, 1896, ibidem 4: 136, by monotypy. VENEZUELA, COSTA RICA.

C. oweni is a junior subjective synonym of Sphinx phalaenoides Drury, 1773, Illust. nat. Hist. exot. Insects 2: 50, index, pl.28, fig.6.

CASTRONIA Jones, 1912, Trans. ent. Soc. Lond. 1912: 420.

Type-species: Castronia collaris Jones, 1912, ibidem 1912: 421, by original designation. Brazil: Parana, Castro.

C. collaris is a junior subjective synonym of Tessellota apostata Schaus, 1905, Proc. U.S. natn. Mus. 29: 227.

CASTULO Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 561.

Type-species: *Castulo plagiata* Walker, 1854, *ibidem* 2: 562, by monotypy. [AUSTRALIA] New Holland: and [Tasmania] Van Dieman's Land.

Castulo was established in the 'Lithosiidae'; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 491; and was transferred back to the 'Lithosiidae' by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903; 498.

CATAPHRACTES Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.99, fig.8.

Type-species: Cataphractes holdingii Felder, 1874, ibidem 2 (Abt.2): pl.99, fig.8, by monotypy. [SOUTH AFRICA]: [Cape of Good Hope] Cap., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

C. holdingii has been misspelt as ‡boldingii by some authors.

PYGETERA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 446.

Established, unnecessarily, as an objective replacement name for *Cataphractes* Felder, 1874, which is not preoccupied by *Cataphractus* Gronovius, 1763, – Mammalia.

‡PIGETERA; Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 78. An incorrect subsequent spelling of Pygetera Kirby, 1892.

CATARCTIA Holland, 1892, Psyche, Camb. 6: 537. NOTO Type-species: Arctia divisa Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 765, by original designation. SIERRA LEONE.

CATENINA Burmeister, 1883, An. Mus. publ. B. Aires 3 (1): 42.

Type-species: *Ecpantheria heterogena* Oberthür, 1881, *Etudes d'Entomologie* 6: 109, pl.15, fig.2, by monotypy. Brazil.

CATOCHRIA Herrich-Schäffer, [1855 July] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.67, fig.380.

Type-species: *Catochria catocaloides* Herrich-Schäffer, [1855] 1850-1858, *ibidem* 1 (1): wrapper, pl.67, fig.380, by monotypy. [SOUTH] AFRICA.

Catochria was also included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' published in 1855 December, Syst. Bearbeitung Schmett. Eur. 6: 91, and again in [1856], Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 11. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

CAULOCERA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 90, 529.

Type-species: *Chamaita crassicornis* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 122, by original designation. [BORNEO]: Sarawak.

CAVIRIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 824.

Type-species: Caviria substrigosa Walker, 1855, ibidem 4: 825, by monotypy. BRAZIL.

CEBYSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 486. GLYPHIPTERIGIDAE

Type-species: *Cebysa leucotelus* Walker, 1854, *ibidem* 2: 486, by monotypy. Holotype δ, [Australia]: Sydney.

Cebysa was established in the Lithosiidae, now Arctiidae Lithosiinae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 490; it was transferred to the Tineidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498; and was placed in the Glyphipterigidae by Fletcher, 1929, Mem. Dep. Agric. India (Ent.) 11: 42.

CECRITA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 992. NOTO

Type-species: *Cecrita guttivitta* Walker, 1855, *ibidem* 5: 992, by monotypy. [CANADA]: Nova Scotia.

CEIRA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 462.

Type-species: *Ceira metaphaea* Walker, 1865, *ibidem* 32: 462, by monotypy. [SRI LANKA] Ceylon.

CELAMODES Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxii, 557.

Type-species: *Manoba rufotincta* Rothschild, 1912, *Novit. zool.* **19**: 231, by original designation. New Guinea: [Irian Jaya], Mt Goliath.

CELEIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 463.

Type-species: Celeia plusiata Walker, 1865, ibidem 32: 463, by monotypy. [INDIA]: [Kanara] Canara.

CENTRONIA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 19.

Type-species: Centronia melanitis Hübner, 1818, ibidem 1: 19, pl.[17], figs 93, 94, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 402. BRAZIL.

CERAEOTRICHA Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 519.

Type-species: *Xenomigia cuneifera* Dognin, 1913, *Mêm. Soc. ent. Belg.* 22: 7, by original designation. Colombia: Mt Tolima.

The date of publication of the part of Seitz containing page 519 is cited on page 513 as 24.x.1925 in both the German and the English editions.

CERAMIDIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 412.

Type-species: *Euchromia fumipennis* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 241, by original designation. [Brazil]: Amazon River, [Tefé] Ega.

CERAMIDIODES Hampson, 1914, Cat. Lepid. Phalaene Br. Mus. (Suppl.) 1: xiv, 255.

Type-species: Ceramidia mathani Rothschild, 1912, Novit. zool. 19: 153, by original designation. [Brazil]: Amazonas, [Tefé] Teffé.

C. mathani is a junior subjective synonym of Ceramidia obscura Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: x, 40, pl.16, fig.5.

CERASANA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 123.

Type-species: *Cerasana anceps* Walker, 1862, *ibidem* (Zool.) 6: 123, by monotypy. Borneo: Sarawak.

Cerasana was established in the Arctiidae; it was placed in the Notodontidae by Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 612.

See also Billetia Joannis, 1900.

‡SERASANA; Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 45.

An incorrect subsequent spelling of Cerasana Walker, 1862.

CERATOPHORA Guenée, 1858. See Sorocaba Moore, 1882.

CERCOCLADIA Draudt, 1915, in Seitz, Gross-Schmett. Erde

Type-species: Cercocladia novicia Draudt, 1915, ibidem 6: 91, pl.15, row d, by monotypy. [BRAZIL]: Matto Grosso, Cuyaba.

The date of publication of the part of Seitz containing page 91 is cited on page 89 as 31.iii.1915 in both the German and the English editions.

CERCOPHORA Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl. 49, fig. 266; [1858], ibidem 1: 74.

Type-species: *Cercophora urophora* Herrich-Schäffer, [1858] 1850-1858, *ibidem* 1: 74 (but included and figured by Herrich-Schäffer, [1855] as '*Cercophora caudata* F.' a misidentification; see below), by monotypy. BRAZIL.

Herrich-Schäffer, [1855], cited a single included nominal species Zygaena caudata Fabricius, [1777], Genera Insect.: 277. In his text published later, Herrich-Schäffer, [1858], cited 'urophora Bd. (non caudata F.) – H-S. f.266' thus indicating that he had originally misidentified the specimen represented in figure 266 which in fact belonged to a hitherto unnamed species. Herrich-Schäffer attributed urophora to Boisduval, an incorrect authorship.

Cercophora is a genus which had a misidentified type-species and should be referred to the Commission under Article 70(a) of the Code.

Cercophora Herrich-Schäffer probably originated from Boisduval who attributed it to himself in Boisduval, 1870, Considérations Lépid. Guatemala: 80.

CERCOPIMORPHA Butler, 1876, J. Linn. Soc. (Zool.) 12: 424.

Type-species: Cercopimorpha homopteridea Butler, 1876, ibidem 12: 424, by original designation. [BRAZIL]: Para.

CERMA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 14.

Type-species: *Cerma cora* Hübner, 1818, *ibidem* 1: 14, pl.[11], figs 59, 60, by monotypy.

Further information in Nye, 1975, Generic Names Moths World 1: 111.

PACHYCERMA Van Duzee, 1897, Bull. Buffalo Soc. nat. Sci. 5: 126, (but attributed to Grote, an incorrect authorship).

Type-species: *Cerma cora* Hübner, 1818, by monotypy. A junior objective synonym of *Cerma* Hübner, 1818.

Pachycerma in the Noctuidae, Acronictinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

CERTILA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 448.

Type-species: *Certila flexuosa* Walker, 1865, *ibidem* **32**: 449, by monotypy. North America.

Certila was established in the Notodontidae; it was placed in the Noctuidae by Dyar, [1903] 1902, Bull. U.S. natn. Mus. 52: 99.

CERURA Schrank, 1802, Fauna Boica 2 (2): 155. NOTO Type-species: Phalaena vinula Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 499, by subsequent designation by Curtis, 1827, Br. Ent. 4: 193. Type-locality: not stated, [Europe].

‡ANDRIA Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) (Opinion 97): 19. Also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140.

Only included species: Phalaena vinula Linnaeus, 1758.

ANDRIA Hübner, 1822, Syst.-alphab. Verz.: 15, 16, 18, 20. Type-species: Phalaena vinula Linnaeus, 1758, by subsequent designation by Grote, 1897, Illte Wschr. Ent. 2: 389.

A junior objective synonym of Cerura Schrank, 1802.

DICRANURA Boisduval, 1828, Eur. Lepid. Index meth. 1: 54. Type-species: Phalaena vinula Linnaeus, 1758, by subsequent designation by Blanchard, 1840, in Castelnau, Hist. nat. Insectes 3: 494.

A junior homonym of *Dicranura R.L.*, 1817, *Jena. allg. Lit.-Ztg* 1 (36): 282, – Lepid., Notodontidae. The objective replacement name is *Cerura Schrank*, 1802.

Boisduval mistakenly attributed the authorship of *Dicranura* to Latreille, probably because Latreille, 1825, in Cuvier, Fanilles nat. Règne Anim.: 473, used the vernacular term Dicranoure.

CERURELLA Kiriakoff, 1962, Tijdschr. Ent. 105: 245. NOTO Type-species: Cerurella natalensis Kiriakoff, 1962, ibidem 105: 245, fig.8, by original designation. [SOUTH AFRICA]: Natal, Balgowan.

CERURIDIA Dyar, 1893, Ent. News 4: 35.

Typc-species: Ceruridia slossonii Dyar, 1893, ibidem 4: 35,

by original designation. TYPE-LOCALITY: not stated, [U.S.A.]:

[New Hampshire, Franconia], see below.

Packard had evidently intended describing a new genus and a new species as ‡'Ceruridia slossoniae'. Whilst Packard's descriptions were still in manuscript, however, Dyar saw the type-specimen from the Slosson collection and published a note expressing the opinion that 'Ceruridia slossonii Packard' was a form of Gluphisia avimacula Hudson, 1891, Ent. News 2: 155, which is its current status. Both the genus and species were thus made available by Dyar.

The type-species was subsequently redescribed from the same specimen by Packard, 1895, Mem. natn. Acad. Sci. 7: 97 as Gluphisia severa Edwards var. slossoniae and type-locality data

given.

CERURINA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 207.

Type-species: Cerura marshalli Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 456, by original designation. [Rhodesia]: Mashonaland, Salisbury.

CERURITES Kernbach, 1967, Ber. naturhist. Ges. Hannover 111: 107. FOSSIL NOTO

Type-species: Cerurites wagneri Kernbach, 1967, ibidem 111: 107, fig. 10, by monotypy. [Germany]: Willershausen.

CERVIPLUSIA Chou & Lu, 1974, Acta ent. sin. 17: 72, 77.

Type-species: Cerriplusia wukongensis Chou & Lu, 1974, ibidem 17: 73, 78, figs 10, 22, by original designation. Holotype &, CHINA: Shensi province, Wukung [sic], (Northwestern College of Agric., Shensi).

Cerviplusia in the Noctuidae, Plusiinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

1

CERYX Wallengren, 1863, Wien. ent. Monatschr. 7: 140.

Type-species: Naclia thyretiformis Wallengren, 1860, ibidem 4: 40, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 108. [SOUTH AFRICA]: interior of eastern Caffraria.

Ceryx was redescribed as new by Wallengren in 1865, K.

svenska Vetensk Akad. Handl. 5 (4): 13.

N. thyretiformis is a junior subjective synonym of Syntomis fulvescens Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 132.

CETOLA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1015.

NOCTUIDAE

Type-species: Cetola dentata Walker, 1855, ibidem 5: 1016,

by monotypy. NEPAL.

Cetola was established in the Notodontidae; it was placed in the Noctuidae by Hampson, 1894, Fauna Br. India (Moths) 2: 434.

CHADISRA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 81.

Type-species: *Chadisra bipars* Walker, 1862, *ibidem* (3) 1: 82, by monotypy. [INDIA]: Hindostan.

See also Metasomera Matsumura, 1925.

CHADISRELLA Kiriakoff, 1967, Tijdschr. Ent. 110: 57.

Type-species: *Chadisrella javensis* Kiriakoff, 1967, *ibidem* 110: 58, fig.22, by original designation. JAVA: [Djakarta] Batavia.

CHADISRINA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 636.

Type-species: Heterocampa variegata Moore, 1879, in Hew-

itson Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 61, by monotypy. INDIA: Darjeeling.

The date of publication of the part of Seitz containing page 636 is cited on page 633 as 21.ii.1930 in the German edition; and as 17.v.1930 in the English edition.

CHADISROIDES Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 35.

Type-species: Ochrostigma ussuriensis Püngler, 1912, in Seitz, Gross-Schmett. Erde 2: 305, pl.49 row b, by original designation. [U.S.S.R.]: Ussuri district.

O. ussuriensis is a subspecies of Stauropus bidentatus Wileman, 1911, Trans. ent. Soc. Lond. 1911; 287, pl.30, fig.9.

See also Wilemanus Nagano, 1916.

‡CHAEROTRICHE; misspelling. See Choerotricha Felder, 1874.

CHAETOGNATHA Felder, 1874. See Tifama Walker, 1855.

CHALEPA Kiriakoff, 1959, Ark. Zool. (2) 12: 332. NOTO Type-species: Chalepa binotata Kiriakoff, 1959, ibidem (2) 12: 333, text-figs 40, 40a, pl.1, fig.11, by original designation. [Burma]: Kambaiti.

CHALLA Moore, 1879, Proc. zool. Soc. Lond. 1879: 398.

Type-species: *Challa discalis* Moore, 1879, *ibidem* 1879: 399, pl.32, fig.7, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 359. HIMALAYAS: northwest.

C. discalis is a junior subjective synonym of Challa bimaculata Moore, 1879, ibidem 1879: 398.

CHAMAITA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 121.

Type-species: Chamaita trichopteroides Walker, 1862, ibidem 6: 121, by original designation. Borneo: Sarawak. See also Homopsyche Butler, 1882.

CHAMESTHEMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.103, fig.13.

PERI

Type-species: *Chamesthema dioptis* Felder, 1874, *ibidem* 2 (Abt.2): pl.103, fig.13, by monotypy. [Brazil]: Amazon River, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 3.

CHAONIA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 29.

Type-species: *Noctua roboris* Fabricius, [1777], *Genera Insect*.: 283, by subsequent designation by Duponchel, 1843, *in* d'Orbigny, *Dict. univl Hist. nat.* 3: 391. GERMANY.

N. roboris is a junior subjective synonym of Bombyx ruficornis Hufnagel, 1766, Berl. Mag. 2: 424.

Invalid type-species designation: Bombyx dodonaea [Denis & Schiffermüller], 1775, was designated by Boisduval, 1836,

Hist. nat. Insectes (Spec. gén. Lépid.) 1: 142.

In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different type-species for each generic name. In his 'Exposé de notre méthode', pages 155-690, no type-species designation was made for any of the genera that he himself used. Under the Code, Article 69(a)(iii), the type-species designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other

authors. The acceptance of Boisduval's type-species designation in the case of *Chaonia* would make no difference to the generic concept.

Chaonia was again proposed by Stephens, 1829 [June], Nom. Br. Insects: 39; and 1829 [July], Syst. Cat. Br Insects (2): 44

CHARADRA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 445.

Type-species: Charadra contigua Walker, 1865, ibidem 32: 446, by monotypy. [U.S.A.]: Georgia. Walker cited the type-locality as 'Georgia?' which is probably correct as the species is known to occur there.

Charadra was established in the Notodontidae; it was transferred to the Noctuidae by Grote & Robinson, 1868, Trans. Ani. ent. Soc. 2: 86.

CHARALA Moore, 1884, Trans. ent. Soc. Lond. 1884: 359.

Type-species: Stilpnotia sericea Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 45, by monotypy. INDIA: [N.], Masuri; and [W.Bengal], [Darjeeling] Darjiling.

CHARIDEA Dalman, 1816. See Marmax Rafinesque, 1815.

ZYGAENIDAE

‡CHARIPODIA; misspelling. See Caripodia Hampson, 1900.

‡CHARNIDA; misspelling. See Charnidas Walker, 1855.

CHARNIDAS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key, as ‡Charnida), 797.

Type-species: Charnidas litura Walker, 1855, ibidem 4: 797, by subsequent designation by Moore, [1883] 1882-3, Lepid. Ceylon 2: 82. [Nepal] Nepaul.

‡CHARNIDA; Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key).

An incorrect (of a multiple) original spelling of *Charnidas* Walker, 1855. ‡*Charnida* was used only in the key to genera and has not subsequently been applied to this taxon.

CHATFIELDIA Grote, 1895, Abh. naturw. Ver. Bremen 14: 48.

Type-species: Notodonta basitriens Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1000, by original designation. [U.S.A.]: New York, Trenton Falls.

CHATRACHARTA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 142.

Type-species: Chatracharta tortricoides Walker, 1862, ibidem 6: 142, by monotypy. Borneo: Sarawak.

Chatracharta was established in the Notodontidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 853; and has been placed in the Lymantriidae by Collenette in the BMNH collection.

CHAZARIA Moore, 1881, Proc. zool. Soc. Lond. 1881: 361.
NOCTUIDAE

Type-species: *Heliothis incarnata* Freyer, [1838] 1839, *Neuere Beitr. Schmett.* 3: 91, pl.256, fig.4, by monotypy. Syntypes [TURKEY]: [Istanbul] Konstantinopel.

Chazaria in the Noctuidae, Noctuinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

RHODOCLEPTRIA Hampson, 1903, Cat. Lepid. Phalaenae Br. Mus. 4: ix, 33.

Type-species: Heliothis incarnata Freyer, [1838], by original designation.

Rhodocleptria is a junior objective synonym of Chazaria Moore, 1881.

In Nye, 1975, Generic Names Moths World 1, the generic name Chazaria was omitted and Rhodocleptria was therefore incorrectly treated as though it could be used as a valid name.

CHELIOSEA Watson nom. n. for Heliocaes Turner, 1940.

Type-species: Spilosoma cosmeta Lower, 1907, Trans. Proc. R. Soc. S. Aust. 31: 170, by original designation (for Heliocaes Turner, 1940). AUSTRALIA: New South Wales, Broken Hill.

HELIOCAES Turner, 1940, Proc. R. Soc. Qd 51: 120.

Type-species: Spilosoma cosmeta Lower, 1907, by original designation.

A junior homonym of *Heliocaes* Bedel, 1906, *Bull. Soc. ent.* Fr. 1906: 92 – Insecta, Coleoptera. The objective replacement name is *Cheliosea* Watson nom. n.

CHELIS Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 256.

ARCT

Type-species: *Phalaena maculosa* [Denis & Schiffermüller], 1775, *Ankündung syst. Werkes Schmett. Wienergegend*: 54, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 263 (but cited as ‡*Cletis maculosa* Gern[ing], an incorrect subsequent spelling of *Chelis*, and an incorrect authorship). [AUSTRIA]: Vienna district.

‡CLETIS; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 263.

An incorrect subsequent spelling of Chelis Rambur, 1866.

CHELONIA Godart, [1823]. See Arctia Schrank, 1802.

CHETONE Boisduval, 1870, Considérations Lépid. Guatemala: 89. Peri

Type-species: *Phalaena catilina* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 126, 152, pl.79, figs E, F, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 185. SURINAM.

P. catilina is dated from the wrapper of the part and not from the title-page of the volume, 1779.

CHIBIDOKUGA Matsumura, 1933, Insecta matsum. 7: 113.

Type-species: Chibidokuga hypenodes Inoue, (in press), by original designation (but cited by Matsumura as 'Dasychira

nigra Hampson'; see below). JAPAN.

C. hypenodes is the name being proposed by Inoue for the taxon which Matsumura misidentified as 'Dasychira nigra Hamps.' and also cited an incorrect reference. Bryk, 1934, in Strand, Lepid. Cat. 62: 58 also cited an incorrect original reference. The correct reference is Selca nigra Hampson, 1898, J. Bombay nat. Hist. Soc. 11: 442, type-locality Sikkim, for a species differing from that in Japan. This is a case of a misidentified type-species which should be referred to the Commission under Article 70(a) of the Code.

CHICHIMECA Hogue, 1963, Contr. Sci. 64: 61. NOCTUIDAE Type-species: Eulithosia thoracica Edwards, 1884, Papilio 4: 44, by original designation. Holotype 9, [U.S.A.]: Arizona. Chichimeca in the Noctuidae, Acronictinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

See also Cyptonychia Druce, 1899.

CHILENA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1070.

LASIOCAMPIDAE
Type-species: Chilena similis Walker, 1855, ibidem 5: 1071,

by monotypy. INDIA: northern, NEPAL.

Chilena was established in the Notodontidae; it was placed in the Limacodidae by Moore, [1860] 1858-59, in Horsfield

& Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 413; and was placed in the Lasiocampidae by Moore, [1887] 1884-87, Lepid. Ceylon 3: 543.

CHIONAEMA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 100, 101; [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 20, 21.

Type-species: Phalaena puella Drury, 1773, *Illust. nat. Hist. exot. Insects* 2: index & 3, pl.2, fig.2, by monotypy (but attributed to Donovan, an incorrect authorship). [INDIA]: Madras.

Chionaema was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

LEPTOTHRIX Heylaerts, 1892, Annls Soc. ent. Belg. 36: 47. Type-species: Leptothrix tettigonioides Heylaerts, 1892, ibidem 36: 47, by monotypy. SUMATRA: Padang-pandjang.

A junior homonym of Leptothrix Menge, 1869, Schr. Ges. Danzig (Nov. Ser.) 2 (2) 10: 240, — Arachnida. There is no objective replacement but van Eecke, 1925, Zoöl. Meded. Leiden 8: 171, and 1927, ibidem 10: 102, transferred L. tettigonioides to Chionaema Herrich-Schäffer, 1855. The latter is therefore available for use as the subjective replacement name.

‡CHIONEMA; Swinhoe, 1901, Ann. Mag. nat. Hist. (7) 8: 124.

An incorrect subsequent spelling of *Chionaema* Herrich-Schäffer, 1855.

CHIONAEMOPSIS Cockerell & Le Veque, 1931, Am. Nat. 65: 354.

YPONOMEUTIDAE FOSSIL

Type-species: Chionaemopsis quadrifasciatus Cockerell & Le Veque, 1931, ibidem 65: 355, photo.1, text-fig.1, by monotypy. U.S.A.: Colorado, Parachute Creek, (Green River Eocene).

Chionaemopsis was compared with Chionaema Herrich-Schäffer, 1855, – Lepid., Arctiidae, by its authors but placed provisionally in the Oecophoridae. Forbes, 1931 October, Am. Nat. 65: 479-480 tentatively placed Chionaemopsis in the Yponomeutidae.

‡CHIONEMA; misspelling. See Chionaema Herrich-Schäffer, 1855.

CHIONOPHASMA Butler, 1886, Trans. ent. Soc. Lond. 1886: 384.

Type-species: *Chionophasma paradoxa* Butler, 1886, *ibidem* **1886**: 385, 441, pl.9, fig.2, by monotypy. Australia: [Queensland], Rockhampton.

CHIONOPHILA Guenée, 1865. See Parasemia Hübner, [1820].

CHIONOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 85, 359.

Type-species: Calligenia apicalis Zeller, 1874, Verh. zool.-bot. Ges. Wien 24 (Abh.): 424, pl.12, fig.1, by original designation. [Brazil]: Rio [de] Janeiro.

CHIRETOLPIS Watson nom. n. for Tricholepis Hampson, 1891.

Type-species: *Tricholepis erubescens* Hampson, 1891, *Illust. typical Specimens Lepid. Heterocera Colln Br. Mus.* 8: 44, pl.139, figs 3, 12, by monotypy (of *Tricholepis* Hampson, 1891). [INDIA]: Nilgiri Hills, plateau.

TRICHOLEPIS Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 3, 44.

Type-species: Tricholepis erubescens Hampson, 1891, by

monotypy

A junior homonym of *Tricholepis* Blanchard, [1851] 1850, *Cat. Colln ent. Mus. Hist. nat. Paris* 1: 155, – Insecta, Coleoptera. The objective replacement name is *Chiretolpis* Watson nom. n.

CHIRIPHE Walker, [1863] 1862, List Specimens lepid. Insects Colln Br. Mus. 26: 1691.

Type-species: *Chiriphe monogrammaria* Walker, [1863] 1862, *ibidem* **26**: 1692, by monotypy. [AUSTRALIA]: Sydney.

Chiriphe was established in the Geometridae; it was placed in the Arctiidae by Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 692, 732.

CHIROMACHLA Strand, 1909, Annls Soc. ent. Belg. 53: 326.

Type-species: *Nyctemera leuconoe* Hopffer, 1858, *Mber. K. preuss. Akad. Wiss.* **1857**: 422, by PRESENT DESIGNATION. MOZAMBIQUE.

Chiromachla was established as a subgenus of Deilemera Hübner, [1820].

CHLANIDOPHORA Berg, 1877, Bull. Soc. imp. Nat. Moscou 52 (3): 9. ARCT

Type-species: Chlanidophora patagiata Berg, 1877, ibidem 52 (3):11, by monotypy. [ARGENTINA]: Carmen de Patagones. Chlanidophora was redescribed as new by Berg, 1877, An. Soc. cient. argent. 4: 95.

CHLIARA Walker, [1858] 1857, List Specimens lepid. Insects Colln Br. Mus. 12: 880 (key), 938.

Type-species: *Phalaena cresus* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 72, pl.142, fig.C, by subsequent designation by Schaus, 1901, *Trans. ent. Soc. Lond.* 1901: 342 (but cited as ‡*croesus*, an incorrect subsequent spelling). [GUYANA]: Berbice.

Chliara was established in the Noctuidae; it was placed in the Notodontidae by Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 697.

P. cresus is dated from the wrapper of part 12, not from the title-page of volume 2 which is dated 1779.

CHLORHODA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xv, 6, 421.

Type-species: *Idalus rufoviridis* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 285, by original designation. [COLOMBIA]: Bogota.

CHLOROCALLIOPE Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 62. NOTO

Type-species: Stauropus rivata Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 466, by original designation. NIGERIA: Akassa.

CHLOROCERAMIS Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 237.

Type-species: *Fentonia viridinota* Hampson, 1896, *Fauna Br. India* (Moths) **4**: 459, by original designation. Bhutan.

CHLOROCHADISRA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 340.

Type-species: Chadisra viridipulverea Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 419, pl.70 row c, by original designation

nation. Type-locality: not stated, [Central African Empire]: [Ouam district, Bozoum]. The locality data cited is that on the type-specimen in the Zool. Mus., Berlin; it was provided by Dr H.J. Hannemann in a personal communication.

CHLOROCRISIA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 408.

Type-species: *Diacrisia irrorata* Rothschild, 1910, *Novit. zool.* 17: 119, by original designation. PERU: Huancabamba.

CHLOROGENIA Meyrick, 1889, Trans. ent. Soc. Lond. 1889: 461.

Type-species: *Chlorogenia cholerota* Meyrick, 1889, *ibidem* **1889**: 461, by monotypy. New Guinea: [Papua New Guinea], Port Moresby.

‡CHLOROMELES; misspelling. See Choromeles Meyrick, 1886.

CHLOROPSINUS Butler, 1876, J. Linn. Soc. (Zool.) 12: 409.

Type-species: Chloropsinus lanceolatus Butler, 1876, ibidem 12: 409, by original designation. [BRAZIL]: [São Paulo] St Paulo.

C. lanceolatus is a junior subjective synonym of Euchromia basalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 251.

See also *Pseudosphenoptera* Butler, 1876, and *Enioche* Druce, 1896.

CHLOROSTAUROPUS Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 135.

Type-species: Stauropus viridissimus Bethune-Baker, 1904, Novit. zool. 11: 378, pl.4, fig.1, by original designation. New Guinea: [Papua New Guinea], Dinawa, Ekeikei.

CHLOROSTOLA Hampson, 1898. See Euchlorostola Watson nom. n. CTEN

CHOEROTRICHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.98, figs 12-17.

LYMA

Type-species: Choerotricha glandulosa Felder, 1874, ibidem 2 (Abt.2): pl.98, fig.14, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 450, but cited as Gogana atrosquama Walker, a nominal species not originally included in Choerotricha. Kirby, however, placed atrosquama as the senior synonym of glandulosa, a nominal species originally included in Choerotricha. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. [PHILIPPINES] ins. philipp., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

C. glandulosa is a junior subjective synonym of Gogana atrosquama Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1921.

GOGANA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1920.

Type-species: Gogana atrosquama Walker, 1866, ibidem 35: 1921, by monotypy. Philippines.

A junior homonym of Gogana Walker, 1865, ibidem 34: 1219, — Lepid., Drepanidae. There is no objective replacement name but G. atrosquama is the senior subjective synonym of Choerotricha glandulosa Felder, 1874, the type-species of Choerotricha Felder, 1874; the latter is therefore available for use as a subjective replacement name.

‡GOGANE; Butler, 1878, Ann. Mag. nat. Hist. (5) 2: 459. An incorrect subsequent spelling of Gogana Walker, 1866.

‡CHAEROTRICHE; Butler, 1881, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 5: 51. An incorrect subsequent spelling of *Choerotricha* Felder, 1874.

CHOKAIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 195.

Type-species: *Pheosia pictibasis* Hampson, 1897, *J. Bombay nat. Hist. Soc.* 11: 282, by original designation. INDIA: Khasia [Hills].

‡CHORIA; misspelling. See Ghoria Moore, 1878. LITH

CHOROMELES Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 773, 785.

Type-species: Choromeles geographica Meyrick, 1886, ibidem (2) 1: 785, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 115. AUSTRALIA: Queensland.

‡CHLOROMELES; Neave, 1939, Nomencl. zool. 1: 708.
An incorrect subsequent spelling of Choromeles Meyrick, 1886.

CHROSTOSOMA Hübner, [1819] 1816, Verz. bekannter Schmett.: 126.

Type-species: *Sphinx evadnes* Stoll, 1781, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* **4**: 129, pl.357, fig.A, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* **1**: 144. Surinam.

S. evadnes is dated from the wrapper of the part and not from the title-page of the volume, 1782.

CHRYSAEGLIA Butler, 1877, Trans. ent. Soc. Lond. 1877: 356.

Type-species: *Lithosia magnifica* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 103, by original designation. [Borneo]: Sarawak.

CHRYSAEGLIODES Strand, 1912, Arch. Naturgesch. 78 (A) 7: 184.

Type-species: Chrysaegliodes noliformis Strand, 1912, ibidem 78 (A) 7: 185, by original designation. GABON: [Ogooué] Ogowe.

CHRYSALLACTIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 84, 334.

Type-species: *Chrysallactis aureorubra* Hampson, 1900, *ibidem* 2: 334, fig.234, by original designation. [New Guinea]: Louisiades, Rossel Island.

‡CHRYSAOR Hübner, [1806]. See Adscita Retzius, 1783. ZYGAENIDAE

‡CHRYSAOR Hübner, 1808. See Empyreuma Hübner, 1818. CTEN

CHRYSAOR Hübner, [1809]. See Belemnia Walker, 1854.

CHRYSASURA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxvi, 730.

Type-species: *Halone flavopunctata* Bethune-Baker, 1904, *Novit. zool.* 11: 421, pl.5, fig.28, by original designation. New GUINEA: [Papua New Guinea], Dinawa and Aroa River.

CHRYSOCALE Walker, 1854, List Specimens lepid. Insects
Colln Br. Mus. 1: 208.

Type-species: Euchromia ignita Walker, 1854, ibidem 1: 209, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 329. COLOMBIA: Santa Fé.

Chrysocale was established as a subgenus of Euchromia Hübner, [1819].

CHRYSOCHLOROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 85, 351.

Type-species: *Chrysochlorosia callistia* Hampson, 1900, *ibidem* **2**: 352, fig.260, by original designation. BOLIVIA: Chaco.

‡CHRYSOCNEMA; misspelling. See Chrysocneme Draudt, 1917.

CHRYSOCNEME Draudt, 1917, in Seitz, Gross-Schmett. Erde 6: 204.

Type-species: *Chrysocneme mirabilis* Draudt, 1917, *ibidem* 6: 205, pl.28, row g, by PRESENT DESIGNATION. COLOMBIA: Pacho.

The date of publication of the part of Seitz containing page 204 is stated on page 201 as 17.ix.1917 in both the German and the English editions.

‡CHRYSOCNEMA; Neave, 1939, Nomencl. zool. 1: 727.

An incorrect subsequent spelling of *Chrysocneme* Draudt, 1917.

CHRYSOCYMA Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 435.

Type-species: *Chrysocyma mesopotamia* Hampson, 1905, *ibidem* (7) **15**: 435, by original designation. [RHODESIA]: Mashonaland, Salisbury.

Chrysocyma was established in the 'Arctiadae, Lithosianae'; it was placed in the Lymantriidae by Collenette, 1935, Stylops 4: 241.

CHRYSOMESIA Hampson, 1903, Ann. Mag. nat. Hist. (7) 11: 350.

Type-species: *Chrysomesia barbicostata* Hampson, 1903, *ibidem* (7) 11: 350, by original designation. New Guinea: [Papua New Guinea], Port Moresby.

CHRYSOPSYCHE Butler, 1880, Ann. Mag. nat. Hist. (5) 5: 387. LASIOCAMPIDAE

Type-species: Choerotriche mirifica Butler, 1878, ibidem (5)

2: 458, by original designation.

Chrysopsyche was established in the Liparidae, now Lymantriidae; it was placed in the Bombycidae by Dewitz, 1881, Nova Acta Acad. Caesar. Leop. Carol. 42 (2): 76; and was transferred to the Lasiocampidae by Aurivillius, 1902, Ent. Tidskr. 23: 273.

CHRYSORABDIA Butler, 1877, Trans. ent. Soc. Lond. 1877: 357.

Type-species: Lithosia viridata Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 225, by original designation. INDIA: Darjeeling.

‡CHRYSORRHABDIA; Bertkau, 1879, Arch. Naturgesch. 45 (2): 124.

An incorrect subsequent spelling of *Chrysorabdia* Butler, 1877.

‡CHRYSORRHABDIA; misspelling. See Chrysorabdia Butler, 1877.

CHRYSOSCOTA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 81, 109.

Type-species: Chrysoscota auranticeps Hampson, 1900, ibidem 2: 110, pl.21, fig. 2, by original designation. New Guinea. [Papua New Guinea, D'Entrecasteaux Islands], Fergusson Island.

CHRYSOSTOLA Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.49, figs 263, 264.

Type-species: Chrysostola humilis Herrich-Schäffer, [1855

June] 1850-1858, *ibidem* 1 (1): wrapper, pl.49, fig.264, by PRESENT DESIGNATION. BRAZIL.

Invalid type-species: Chrysostola mellina Herrich-Schäffer, [1855 December] 1850-1858, ibidem 1 (1): wrapper, pl.74, fig.427, a nominal species not originally included in Chrysostola, and not linked with one of the originally included nominal species when cited by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 372. Further information on the dates of Herrich-Schäffer's Samml. neuer oder wenig bekannter aussereur. Schmett. is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

CHRYSOZANA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xx, 86, 553.

Type-species: Chrysozana croesus Hampson, 1900, ibidem 2: 554, fig. 406, by original designation. BOLIVIA: Songo.

CHURINGA Moore, 1878, Proc. zool. Soc. Lond. 1878: 9.

Type-species: *Churinga rufifrons* Moore, 1878, *ibidem* **1878**: 10, pl.1, fig.12, by original designation. INDIA: Darjeeling.

CIACA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 396.

Type-species: *Ciaca urapterides* Walker, 1865, *ibidem* 32: 397, by monotypy. Sumatra.

CICYNNA Walker, [1858] 1857, List Specimens lepid. Insects Colln Br. Mus. 13: 1079 (key), 1104.

Type-species: Cicynna sericea Walker, [1858] 1857, ibidem 13: 1105, by monotypy. [DOMINICAN REPUBLIC]: St Domingo.

C. sericea is a junior subjective synonym of Elymiotis noto-dontoides Walker, 1857, ibidem 11: 609.

Cicynna was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 272.

See also Elymiotis Walker, 1857.

CIFUNA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1158 (key), 1172.

Type-species: *Cifuna locuples* Walker, 1855, *ibidem* **5**: 1173, by monotypy. [BANGLADESH]: [Sylhet] Silhet.

Cifuna was established in the Drepanulidae, now Drepanidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 458.

CIMBINA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 444.

Type-species: Cimbina cucullodes Walker, 1865, ibidem 32:445, by monotypy. JAVA.

‡LIMBINA; Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 608.

An incorrect subsequent spelling of Cimbina Walker, 1865.

CIMOLA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 817.

Type-species: *Cimola opalina* Walker, 1855, *ibidem* 4: 817, by monotypy. [South Africa]: [Durban] Port Natal.

CINCIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 538.

Type-species: *Cincia conspersa* Walker, 1854, *ibidem* 2: 538, by monotypy. JAMAICA.

CINCIANA Strand, 1915, Arch. Naturgesch. 80 (A) 10: 96.

Type-species: Cinciana nubica Strand, 1915, ibidem 80 (A) 10: 96, by original designation. SUDAN: [Nubian Desert] Nubien.

CINCIANELLA Strand, 1928, Arch. Naturgesch. 92 (A) 8: 51.

Established, unnecessarily, as an objective replacement name for *Cinciana* Strand, 1915, which is not a junior homonym of ‡*Cinciana* Oppenheim, 1891, *Z. dt. geol. Ges.* 43: 462, as the latter is an incorrect subsequent spelling of *Cincinna* Hübner, 1810, *Epistolae*: 1, – Mollusca.

CINCIANELLA Strand, 1928. See Cinciana Strand, 1915.

CISPIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 857.

Type-species: Cispia punctifascia Walker, 1855, ibidem 4: 857, by subsequent designation by Moore, [1883] 1882-3, Lepid. Ceylon 2: 91. [BANGLADESH]: [Sylhet] Silhet.

‡CISPIDIA; Kozhanchikov, 1950, Fauna SSSR (N.S.) 42 (Lepid. 12): 308.

An incorrect subsequent spelling of Cispia Walker, 1855.

‡CISPIDIA; misspelling. See Cispia Walker, 1855. LYMA

CISSEPS Franclemont, 1936, Ent. News 47: 275. CTEN Type-species: Glaucopis fulvicollis Hübner, [1818], Samml. exot. Schmett.: pl.[164], by monotypy (of Scepsis Walker, 1854). Type-locality: not stated, [North America].

Cisseps was proposed as the objective replacement name for Scepsis Walker, 1854.

SCEPSIS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 285.

Type-species: Glaucopis fulvicollis Hübner, [1818], by monotypy.

Scepsis was established as a subgenus of Ctenucha Kirby, 1837.

A junior homonym of *Scepsis* Walker, 1850, in Saunders, *Insecta Saundersiana* 1: 71 – Insecta, Diptera. The objective replacement name is *Cisseps* Franclemont, 1936.

CISSURA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 488.

Type-species: Cissura decora Walker, 1854, ibidem 2: 489, by monotypy. Brazil.

See also Cratosia Felder, 1874.

CISTHENE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 533.

Type-species: Cisthene subjecta Walker, 1854, ibidem 2: 534, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 289. U.S.A.

CLADOCERA Rambur, 1858. See Episema Ochsenheimer, 1816. NOCTUIDAE

CLADOPHORA Geyer, 1832. See Gynaephora Hübner, [1819].

CLAMARA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1099.

Type-species: Clamara terminata Walker, 1855, ibidem 5: 1099, by monotypy. [Brazil]: Rio [de] Janeiro.

Clamara was established in the Notodontidae; it was placed in the Limacodidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 525.

CLAPHE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 993.

LASIOCAMPIDAE

Type-species: Claphe guttivena Walker, 1855, ibidem 5: 993,

by monotypy. [Brazil]: Rio [de] Janeiro.

Claphe was established in the Notodontidae; it

Claphe was established in the Notodontidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 848.

CLASTOGNATIIA Felder, 1874, in Felder & Rogenhofer,

Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.104, fig.8.

Type-species: Clastognatha integra Felder, 1874, ibidem 2 (Abt.2): pl.104, fig.8, by monotypy. [Brazil]: River Amazon, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

CLATHE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 994.

LASIOCAMPIDAE

Type-species: *Clathe arida* Walker, 1855, *ibidem* 5: 994, by monotypy. TYPE-LOCALITY: not known, [AUSTRALIA].

ABELLA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 272.

Type-species: Abella costalis Walker, 1862, ibidem (3) 1: 272, by monotypy. Type(s) 9, [Australia]: [Queensland],

Moreton Bay, (Type(s) not found).

A junior homonym of Abella Heyden, 1826, Isis Oken, Jena 1826: 611, — Arachnida. There is no objective replacement name but from the original description of Abella and A. costalis it is likely that A. costalis is a junior subjective synonym of C. arida Walker, 1855, the type-species of Clathe Walker; the latter is thus available for use as a subjective replacement name.

Abella Walker, 1862, was established in the Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 924, and since then has not been cited; it is here transferred to the Lasiocampidae as a junior subjective synonym of Clathe Walker, 1855.

CLEAPA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1036.

Type-species: Cleapa latifascia Walker, 1855, ibidem 5: 1037, by monotypy. [Indonesia] East Indies.

See also Marushachia Matsumura, 1934.

CLEMENDANA Schaus, 1929, Proc. ent. Soc. Wash. 31: 46.

Type-species: Clemendana pacifera Schaus, 1929, ibidem 31: 46, by original designation. BRAZIL: Santa Catarina.

CLEMENSIA Packard, 1864, Proc. ent. Soc. Philad. 3: 100.

Type-species: Clemensia albata Packard, 1864, ibidem 3: 101, by monotypy. U.S.A.: M[ain]e; Norway and Brunswick. See also Uxia Walker, 1866, and Repa Walker, 1866.

‡CLEOLOSIA; misspelling. See Aeolosia Hampson, 1900.

CLEPTOPHASIA Prout, 1918, Novit. zool. 25: 397 (key), 405.

Type-species: Oricia scissa Warren, 1909, ibidem 16: 72, by original designation. [BRAZIL]: Upper Amazons, Fonte Boa.

CLERCKIA Aurivillius, 1882, K. svenska VetenskAkad. Handl. (Nov. Ser.) 19 (5): 157, 158.

Type-species: *Phalaena fulvia* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 509, by monotypy. [INDONESIA] 'Indiis'.

See also Exotrocha Meyrick, 1886.

CLETHROGYNA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 281.

Type-species: Orgya splendida Rambur, [1837], Faune ent. Andalousie 2: pl.15, figs 3-6, by subsequent designation by Kirby, 1870, Zool. Rec. 6: 389. [Spain]: [Andalusia] Andalousie.

Clethrogyna is dated from the wrapper of livraison 2, and not from the title-page of the work, 1858.

‡CLETIS; misspelling. See Chelis Rambur, 1866. ARCT

CLETORNA Boisduval, 1870. See Hyalurga Hübner, [1819].

PERI

CLISOBARA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 269.

Type-species: Clisobara catocalina Walker, [1865] 1864, ibidem 31: 269, by monotypy. Australia: and [Queensland], Moreton Bay.

CLOESIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 90, 345.

Type-species: *Ichoria parthia* Druce, 1889, *Ann. Mag. nat. Hist.* (6) **4**: 83, by original designation. NICARAGUA: Chontales, PANAMA: Bugaba.

Cloesia was proposed as ‡Cloësia, but under the Code, Article

32(c)(i), the diaeresis must be deleted.

CLOSTERA Samouelle, 1819, Entomologist's useful Compendium: 247, but attributed to Hoffmansegg, an incorrect authorship.

Type-species: *Phalaena curtula* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 503, by subsequent designation by Curtis, 1838, *Br. Ent.* 15: 715. Type-Locality: not stated, [Europe].

Invalid type-species designation: *Bombyx reclusa* [Denis & Schiffermüller], 1775, was designated by Boisduval, 1836, *Hist*.

nat. Insectes (Spec. gén. Lépid.) 1: 142.

In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different type-species for each generic name. In his 'Exposé de notre méthode', pages 155-690, no type-species designation was made for any of the genera that he himself used. Under the Code, Article 69(a)(iii), the type-species designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other authors. The acceptance of Boisduval's designation in the case of Clostera would make no difference to the generic concept.

‡MELALOPHA Hübner, [1806], Tentamen determinationis

digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) (Opinion 97): 19. Also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140.

Only included species: Phalaena curtula Linnaeus, 1758.

MELALOPHA Hübner, 1822, Syst.-alphab. Verz.: 14, 16, 19, 20

Type-species: *Phalaena curtula* Linnaeus, 1758, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 609.

A junior objective synonym of Clostera Samouelle, 1819.

THOLODA Gistl, 1848, Naturg. Thierreichs: xi.

Tholoda was established as an objective replacement name for Clostera Samouelle, 1819, but was attributed by Gistl to Hoffmansegg, an incorrect authorship.

CLOSTERELLUS Fletcher nom. n. for Closteroides Kiriakoff, 1977.

Type-species: Closteroides dorsalis Kiriakoff, 1977, Mitt. münch. ent. Ges. 66: 34, fig.4, by original designation (for Closteroides Kiriakoff, 1977). Sumatra: northern, Deli, Dolok Merangir.

Prof. Dr S.G. Kiriakoff was informed of the homonymy and has suggested this replacement name.

nas suggested this replacement name

CLOSTEROIDES Kiriakoff, 1977, Mitt. münch. ent. Ges. 66: 33.

Type-species: Closteroides dorsalis Kiriakoff, 1977, by original designation.

A junior homonym of Closteroides Tomlin, 1929, Proc.

malac. Soc. Lond. 18: 258, - Mollusca. The objective replacement name is Closterellus Fletcher nom. n.

CLOSTEROIDES Kiriakoff, 1977. See Closterellus Fletcher nom. n. NOTO

CLOSTRA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 273.
NOTO

Type-species: *Clostra tenebriplaga* Walker, 1862, *ibidem* (3) 1: 274, by monotypy. [INDIA]: Hindostan.

CLUACA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 268.

Type-species: Cluaca rubricosta Walker, [1865] 1864, ibidem 31: 268, by monotypy. [AUSTRALIA]: Tasmania.

C. rubricosta is a junior subjective synonym of Castulo doubledayi Newman, 1857, Trans. ent. Soc. Lond. (2) 4: 55.

Cluaca was established in the Callimorphidae, now Arctiidae, Arctiinae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 491; and was transferred to the 'Lithosiidae' by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498.

CLYSTEA Watson nom. n. for Scytale Felder, 1874. CTEN Type-species: Scytale platyzona Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.24, by monotypy (of Scytale Felder, 1874). [Colombia]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 9.

SCYTALE Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.24.

Type-species: Scytale platyzona Felder, 1874, by monotypy. A junior homonym of Scytale Latreille, in Sonnini & Latreille, in Buffon, [1803] An X, Hist. Nat. Reptiles 3: 158, – Reptilia. The objective replacement name is Clystea Watson nom. n.

Scytale was first used by Gronovius, 1763, Zoophylacium Gronovianum (1): 19, and later by Meuschen, 1778, Mus. Gronovianum: 12, – Reptilia. Both these works were rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1954, Opin. Decl. int. Commn zool. Nom. 5 (Opinions 260 and 261).

CNETHOCAMPA Stephens, 1828. See Thaumetopoea Hübner, [1820].

CNETHODONTA Staudinger, 1887, in Romanoff, Mém. Lépid. 3: 214, 215.

Type-species: Cnethodonta grisescens Staudinger, 1887, ibidem 3: 214, pl.12, fig.11, by monotypy. [U.S.S.R.]: Vladivostok, Askold [Island], Ussuri, Suifun.

‡CNETOCAMPA; misspelling. See Thaumetopoea Hübner, [1820].

COBANILLA Moore, [1883] 1882-3, Lepid. Ceylon 2: 120.

Type-species: Cobanilla marginata Moore, [1883] 1882-3, ibidem 2: 121, pl.124, fig.4, by monotypy. [SRI LANKA] Ceylon. Cobanilla was established in the Drepanulidae, now Drepanidae; it was placed in the Lymantriidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 392.

COBORISA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 915.

Type-species: *Coborisa fenestrata* Walker, 1855, *ibidem* 4: 915, by monotypy. MEXICO.

COCASTRA Boisduval, 1870. See Composia Hübner, [1820].

COCYTIA Boisduval, 1828, Essai Monogr. Zygénides: 22.

Type-species: *Cocytia durvillii* Boisduval, 1828, *ibidem*: 22, pl.1, fig.1, by monotypy. NEW GUINEA.

RHOPTROPHALAENA Hampson, 1918, Novit. zool. 25: 366. Proposed, unnecessarily, as an objective replacement name for Cocytia Boisduval, 1828, which Hampson considered to be a homonym of Cocytius Hübner, [1819] 1816, Verz. bekannter Schmett.: 140, – Lepid., Sphingidae.

‡ROPTROPHALAENA; Neave, 1940, Nomencl. zool. 4: 83. An incorrect subsequent spelling of Rhoptrophalaena Hampson, 1918.

CODALITHIA Holloway, 1979, Series Ent. 15: 427.

Type-species: *Codalithia quincuncialis* Holloway, 1979, *ibidem* 15: 428, text-fig.103, pl.78, fig.8, by original designation. Holotype & New Caledonia: Pic du Pin, (BMNH).

Codalithia in the Noctuidae, Acontiinae is cited here as it was not included in Nye, 1975, Generic Names Moths World

CODIOSOMA Kirby, 1892. Sce Kodiosoma Stretch, 1872.

CODITA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 40.

Type-species: Codita decipiens Kiriakoff, 1962, ibidem 114: 41, fig.52, by original designation. [ZAIRE]: Sankaru, Lusambo.

COELODASYS Packard, 1864, Proc. ent. Soc. Philad. 3: 363.

Type-species: *Phalaena unicornis* Smith, 1797, *in* Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 171, pl.86, by PRESENT DESIGNATION. [U.S.A.]: Georgia.

COENOCHROMIA Hübner, [1819]. See Amata Fabricius, 1807.

COENOSTEGIA Mabille, 1891, Annls Soc. ent. Fr. (6) 10 (Bull. ent.): cxlvi.

Type-species: Coenostegia flavens Mabille, 1891, ibidem (6) 10 (Bull. ent.): cxlvii, by subsequent designation by Kiriakoff, 1969, Faune Madagascar 29: 187. MADAGASCAR: [Menabé], cited on the lectotype label.

‡CAENOSTEGIA; Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 23.

An incorrect subsequent spelling of *Coenostegia* Mabille, 1891.

COFIMPACIA Holloway, 1979, Series Ent. 15: 528.

Type-species: Cofimpacia luteata Holloway, 1979, ibidem 15: 529, text-fig.149, pl.87, fig.14, by original designation. Holotype &, New Caledonia: Grand Lac, (BMNH).

Cofimpacia in the Noctuidae, Hypeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

COLAX Hübner, [1819] 1816, Verz. bekannter Schmett.: 141.

Type-species: *Phalaena apulus* Cramer, 1776, *Uitlandsche Kapellen (Papillons exot.)* 1: 139, pl.88, fig.E, by monotypy. Surinam.

P. apulus is dated from the wrapper of part 8, not from the title-page which is dated 1779.

COLETA Roepke, 1949, Trans. R. ent. Soc. Lond. 100: 53.

Type-species: *Phalaena coleta* Stoll, 1781, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* **4**: 153, pl.368, fig.H, by monotypy. [Indonesia]: [Ambon I.] Amboina.

Coleta was established as a subgenus of Nyctemera Hübner,

[1820].

P. coleta is dated from the wrapper of the part and not from the title-page of the volume, 1782.

COLLARTISA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 81. THYRET

Type-species: *Collartisa collartorum* Kiriakoff, 1953, *ibidem* **26**: 81, pl.2, fig.11, pl.7, figs 94, 95, by original designation. [ZAIRE] Congo Belge: Tshuapa, Bokuma.

COLLENETTEMA Griveaud, 1977, Faune Madagascar 43: 259.

Type-species: Cypra crocipes Boisduval, 1833, Nouv. Annls Mus. Hist. nat. Paris 2: 235, by original designation. MADAGASCAR.

COLLETRIA Felder & Rogenhofer, 1875, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.139, fig.7 (but attributed to Zeller, an incorrect authorship).

Type-species: Colletria pyrrhocrocis Felder & Rogenhofer, 1875, ibidem 2 (Abt.2): pl.139, fig.7 (but attributed to Zeller, an incorrect authorship), by monotypy. [Colombia]: Bogota.

COLLETRIA Nolken & Zeller, 1876, Horae Soc. ent. Ross. 12: 76.

Type-species: Colletria pyrrhocrocis Felder & Rogenhofer, 1875, by monotypy.

A junior objective synonym and a junior homonym of *Colletria* Felder & Rogenhofer, 1875, – Lepid., Arctiidae. The objective replacement name is *Colletria* Felder & Rogenhofer, 1875.

COLLETRIA Nolken & Zeller, 1876. See Colletria Felder & Rogenhofer, 1875.

COLLITA Moore, 1878, Proc. zool. Soc. Lond. 1878: 16.

Type-species: *Bombyx griseola* Hübner, [1803] 1796, *Samml. eur. Schmett.* 3: pl.23, fig.97, by original designation. Europe.

COLLOCALIODES Kiriakoff, 1957, Bull. Annls Soc. r. ent. Belg. 93: 145.

Type-species: *Metarctia dracoena* Kiriakoff, 1953, *Annls Mus. r. Congo Belge* (8vo Sci. zool.) **26**: 53, pl.4, fig.45, pl.7, fig.85, by original designation. [ZAIRE] Congo Belge: Sankuru, Lusambo.

Collocaliodes was established as a subgenus of Metarctia Walker, 1855.

COLLYTA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 452.

Type-species: Collyta lanceolata Walker, 1865, ibidem 32: 452, by monotypy. [Australia]: [Queensland], Moreton Bay. C. lanceolata is a junior subjective synonym of Destolmia lineata Walker, 1855, ibidem 5: 992.

See also Destolmia Walker, 1855.

COLNETTIA Holloway, 1979, Series Ent. 15: 490.

Type-species: Colnettia brinoni Holloway, 1979, ibidem 15: 490, text-figs 126, 127, 131, pl.85, figs 11, 12, by original designation. Holotype &, New Caledonia: Plaine des Lacs, (BMNH).

Colnettia in the Noctuidae, Ophiderinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

AGAN

COMACHARA Franclemont, 1939, Ent. News 50: 216. LITH Type-species: Comachara cadburyi Franclemont, 1939, ibidem 50: 216, fig., by original designation. U.S.A.: Pennsylvania, Homebrook, Lower Merion Township.

Comachara was established in the Noctuidae; it was transferred to the Arctiidae, Lithosiinae, by McDunnough, 1943,

Can. Ent. 75: 62.

COMACLA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 276.

Type-species: Comacla murina Walker, [1865] 1864, ibidem

31: 276, by monotypy. [U.S.A.]: E. Florida.

C. murina is a junior subjective synonym of Pagara simplex Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1679.

See also Pagara Walker, 1856, and Vanessodes Grote & Robinson, 1870.

COMANA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 495.

Type-species: Comana collaris Walker, 1865, ibidem 32: 496,

by monotypy. AUSTRALIA: northern.

Comana was established in the Limacodidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 442; it was transferred from the Liparidae but not placed elsewhere by Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 440. Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 1, stated that it was best placed in the Limacodidae.

COMARCHIS Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 693, 739.

Type-species: Comarchis isophragma Meyrick, 1886, ibidem (2) 1: 739, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 499. AUSTRALIA: Tasmania, Launceston.

COMIDAVA Walker, [1863] 1862, List Specimens lepid. Insects Colln Br. Mus. 26: 1695.

Type-species: Comidava obliquilinea Walker, [1863] 1862, ibidem 26: 1695, by subsequent designation by Fletcher, 1979, in Nye, Generic Names Moths World 3: 49. VENEZUELA.

C. obliquilinea is a junior subjective synonym of Hemiceras cadmia Guenée, 1852, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 6: 383, pl.13, fig.2.

Comidava was established in the Geometridae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 333.

See also Hemiceras Guenée, 1852.

COMMONIA Kiriakoff, 1968, in Wytsman, Genera Insect.

Type-species: *Destolmia atalopa* Turner, 1931, *Proc. Linn. Soc. N.S.W.* **65**: 329, by monotypy. Australia: Western Australia, Denmark.

COMPOSIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 179.

Type-species: *Bombyx credula* Fabricius, 1775, *Syst. Ent.*: 584, by monotypy. AMERICA: [tropical].

COCASTRA Boisduval, 1870, Considérations Lépid. Guate-mala: 88.

Type-species: *Bombyx credula* Fabricius, 1775, by original designation.

A junior objective synonym of Composia Hübner, [1820].

COMPSOCHROMIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 71.

Type-species: *Pseudapiconoma compsa* Jordan, 1904, *Novit. zool.* 11: 441, by original designation. ANGOLA: Pungo Andongo.

Compsochromia was established as a subgenus of Balacra Walker, 1856.

COMPSOPRIUM Blanchard, 1852, in Gay, Hist. Chile (Zool.) 7: 67.

Type-species: Compsoprium vittigerum Blanchard, 1852, ibidem 7: 67, by monotypy. CHILE: Coquimbo.

‡CONSOPRIUM; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1644.

An incorrect subsequent spelling of *Compsoprium* Blanchard, 1852.

CONGRUIA Dyar, 1908, Proc. ent. Soc. Wash. 9: 45. NOTO Type-species: Congruia congrua Dyar, 1908, ibidem 9: 45, by original designation (in accordance with the Code, Article 68(a)(i)). Brazil: Rio [de] Janeiro.

CONIGEPHYRA Collenette, 1934, Ann. Mag. nat. Hist. (10) 13: 218.

Type-species: Laelia unipunctata Möschler, 1887, Abh. senckenb. naturforsch. Ges. 15: 74, by original designation. [Ghana] Goldküste: Aburi.

CONILEPIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 83, 219.

Type-species: *Oenistis nigricosta* Leech, 1889, *Proc. zool. Soc. Lond.* **1888**: 598, pl.30, fig.11, by original designation. Type-locality: not stated, [Japan].

CONIORTODES Collenette, 1955. See Melgona Nye nom.

CONOSIA Hampson, 1900. See Euconosia Watson nom. n. LITH

‡CONSOPRIUM; misspelling. See Compsoprium Blanchard, 1852.

CONTREBIA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 9: 69 (key), 134.

Type-species: Contrebia extrema Walker, 1856, ibidem 9: 134, by monotypy. HONDURAS.

Contrebia was established in the Noctuidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 596.

COPAENA Herrich-Schäffer, [1855] 1850-1858, Sanuml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.[49], fig.260.

Type-species: Copaena scapularis Herrich-Schäffer, [1855] 1850-1858, ibidem 1(1): wrapper, pl.[49], fig.260, by monotypy. BRAZIL.

C. scapularis is a junior subjective synonym of Zygaena eriphia Fabricius, [1777], Genera Insect.: 276.

Copaena Herrich-Schäffer probably originated from Boisduval who attributed it to himself in Boisduval, 1870, Considérations Lépid. Guatemala: 81.

See also Antichloris Hübner, 1818, and Illipula Walker, 1854.

CORANIA Schaus, 1939, Ann. Carnegie Mus. 27: 325. NOTO Type-species: Corania pedrana Schaus, 1939, ibidem 27: 326, pl.31, fig.9, by original designation. ARGENTINA: northern, prov. Oran.

CORCURA Walker, 1862, J. Proc. Linn. Soc. 6 (Zool.): 110.

Type-species: *Corcura torta* Walker, 1862, *ibidem* 6 (Zool.): 110, by monotypy. Borneo: Sarawak.

C. torta is a junior subjective synonym of Teulisna chiloides Walker, 1862, J. Proc. Linn. Soc. 6 (Zool.): 109.

See also Ityca Walker, 1866, and Diastrophia Felder, 1874.

COREMATURA Butler, 1876, J. Linn. Soc. (Zool.) 12: 403.

Type-species: Glaucopis chrysogastra Perty, [1833] 1830-1834, in Martius, Delectus Anim. articulatorum Bras.: 157, by original designation. BRAZIL: equatorial.

COREODONTA Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 32, 35.

Type-species: Coreodonta coreana Matsumura, 1924, ibidem 9: 32, by original designation. KOREA: Shakuoji.

COREURA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 142.

Type-species: Coreura euchromoides Walker, [1865] 1864, ibidem 31: 142, by monotypy. [COLOMBIA]: Bogota and New Grenada.

CORINELLA Kiriakoff, 1967, Tijdschr. Ent. 110: 54. NOTO Type-species: Allodonta vittata Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 643, pl.80 row d, by original designation. SUMATRA: Mt Korintji.

CORMA Moore, 1881. See Ambadra Moore, 1883. NOTO

CORREBIA Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.49, fig.265.

Type-species: *Correbia ceramboides* Herrich-Schäffer, [1855] 1850-1858, *ibidem* 1(1): wrapper, pl.49, fig.265, by monotypy. Brazil.

C. ceramboides is a junior subjective synonym of Euchromia lycoides Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 256.

See also Pionia Walker, 1854.

CORREBIDIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xx, 31, 517.

Type-species: *Pionia terminalis* Walker, 1856, *List Specimens lepid. Insects Colln Br. Mus.* 7: 1633, by original designation. BRAZIL.

CORTICOMIS van Eecke, 1924, Nova Guinea 15 (Zool.) (1): 46.

ANTHELIDAE

Type-species: *Corticomis eupterotioides* van Eecke, 1924, *ibidem* 15 (Zool.) (1): 46, pl.1, figs 6, 6a, by original designation. New Guinea: [Irian Jaya], Doormantop.

Corticomis was established for two new nominal species; only Corticomis eupterotioides was followed by the formula 'Nov. gen. spec. nov.', thus satisfying the conditions required by Article 68(a)(i) of the Code for such a formula to be interpreted as an original designation.

Corticomis was established in the Lymantriidae; it is here transferred to the Anthelidae.

CORYDALLA Walker, 1856. See Agyrta Hübner, [1820].
CTEN

CORYMBIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 33: 765.

Type-species: Corymbia smerinthoides Walker, 1865, ibidem 33: 765, by subsequent designation by Berio, 1966, Annali Mus. civ. Stor. nat. Giacomo Doria 76: 54. [GUYANA]: Demerara, [BRAZIL]: Vila Nova.

Corymbia was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 341.

CORYPHYALA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 98. Nomenclaturally available but without included nominal species until, [1856], Samml. neuer oder

wenig bekannter aussereur. Schmett. 1 (1): 18, wrapper, pl.93, fig.532.

Type-species: Coryphyala orbigera Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): 18, wrapper, pl.93, fig.532; [1858], ibidem: 70, by subsequent monotypy. BRAZIL.

Coryphyala was established in a key in the upper part of page 98; in the lower part of the page Herrich-Schäffer included in Coryphyala his then unnamed fig.532, and two nominal species which he doubtfully included. Under the Code, Article 67(h) and 69(a), there were no nominal species eligible to be fixed as type-species when Coryphyala was established.

Coryphyala was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

COSCINIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 169.

Type-species: *Phalaena cribrum* Linnaeus, 1761, *Fauna Suecica* (Edn 2): 302, by subsequent designation by Hampson, 1901, *Cat. Lepid. Phalaenae Br. Mus.* 3: 203, 204, but cited as *P. cribraria* Linnaeus, 1758, a nominal species not originally included in *Coscinia*. Hampson, on page 204, placed *cribraria* as the senior synonym of *cribrum*, a nominal species originally included in *Coscinia*. Under the *Code*, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. Sweden: southern.

P. cribrum is a junior subjective synonym of P. cribraria Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 507.

Incorrect type-species: *Phalaena striata* Linnaeus, 1758, a nominal species not originally included in *Coscinia*, and not linked in synonymy with one of the originally included nominal species when designated by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 343.

COSCODACA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 245.

Type-species: *Spatalia acharista* West, 1932, *Novit. zool.* 37: 210, by original designation. Philippines: Luzon I., subprov. Benguet, Klondyke.

COSMETHIS Hübner, [1820] 1816, Verz. bekannter Schmett.: 179. GEOMETRIDAE

Type-species: *Phalaena barbara* Stoll, 1781, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 151, pl.368, fig.B, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 498. [INDONESIA]: [Moluccas], Ambon Island.

P. barbara is dated from the wrapper of the part, not from the title-page of the volume, 1782.

Cosmethis was established in the 'Phalaenae'; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, ibidem 1: 498; it was placed in the Boarmiidae, now Geometridae Ennominae, by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 312.

COSMOSOMA Hübner, [1823], Samml. exot. Schmett. 2: pl.[156].

Type-species: Cosmosoma omphale Hübner, [1823], ibidem 2: pl.[156], by monotypy. Type-locality: not stated, [Central America].

C. omphale is a junior subjective synonym of Sphinx auge Linnaeus, 1767, Syst. Nat. (Edn 12): 807.

‡AUGE Oken, 1815, Okens Lehrbuch Naturg. 3: 751.

Included in a work rejected for nomenclatural purposes by Int. Commn zool. Nom., 1956, *Opin. Decl. int. Commn zool. Nom.* 14 (Opinion 417): 3.

One of the several included species is *Sphinx auge* Linnaeus, 1767, *Syst. Nat.* (Edn 12): 807, which is currently placed in

Cosmosoma Hübner, [1823].

COSSA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 232.

Type-species: Cossa basigera Walker, [1865] 1864, ibidem 31: 233, by monotypy. INDIA: S. Hindostan.

COSTALIMAIDA Orfila, 1958, Physis, B. Aires 21 (60): 68.

Type-species: Cisthene triplaga Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 434, by original designation. Paraguay: Sapucay.

COSTARCHA Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 5, 53.

Type-species: Costarcha indistincta Hampson, 1891, ibidem 8: 53, pl.140, fig.22, by monotypy. INDIA: Nilgiri Hills, plateau.

COTHOCIDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 499.

Type-species: Cothocida nigrifera Walker, 1865, ibidem 32: 499, by monotypy. North America.

‡COTOBARA; misspelling. See Cottobara Walker, 1862.

COTTOBARA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 95.

Type-species: Cottobara concinna Walker, 1862, ibidem (3)

1: 95, by monotypy. [BRAZIL]: Amazon region.

Cottobara was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 274.

‡COTOBARA; Gaede, 1934, in Strand, Lepid. Cat. 59: 215. An incorrect subsequent spelling of Cottobara Walker, 1862.

COUTHA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 234.

Type-species: Coutha semiclusa Walker, [1865] 1864, ibidem 31: 234, by monotypy. [INDONESIA]: [Moluccas, Seram I.] Ceram.

COXEYA Schaus, 1937, Proc. U.S. natn. Mus. 84: 567. NOTO Type-species: Coxeya sinistra Schaus, 1937, ibidem 84: 567, by oroginal designation. ECUADOR: Dos Puentes.

COZISTRA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 342.

Type-species: Cozistra subnudata Walker, 1865, ibidem 32: 342, by monotypy. [INDONESIA]: [Moluccas], [Batjan I.] Batchian.

Cozistra was established in the Liparidae, now Lymantriidae; it was transferred to the Boarmiidae, now Geometridae Ennominae, by Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 442.

COZOLA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 390.

Type-species: *Cozola leucospila* Walker, 1865, *ibidem* 32: 390, by original designation. [SULAWESI] Celebes.

CRAGIA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 53.

Type-species: *Ilema distigmata* Hampson, 1901, *Ann. Mag. nat. Hist.* (7) 8: 181, by original designation. [NIGERIA]: Old Calabar, [KENYA]: Nairobi Plains, Kikuyu.

CRAMBIDIA Packard, 1864, Proc. ent. Soc. Philad. 3: 99.

Type-species: Crambidia pallida Packard, 1864, ibidem 3: 99, by monotypy. U.S.A.: Mass[achusetts]; and M[ain]e, Brunswick.

CRAMBOMETRA Prout, 1915, Novit. zool. 22: 376. NOTO Type-species: Crambometra derelicta Prout, 1915, ibidem 22: 376, by original designation. SOUTH AFRICA: [Natal], Tongaat.

Crambometra was established in the Geometridae; it was placed in the Notodontidae by Janse, 1920, Ann. Transv. Mus. 7: 218.

CRAMBOMORPHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, figs 9, 10.

Type-species: Crambomorpha umbrifera Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.9, by subsequent designation (for Macrocrambus Kirby, 1892) by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 337. [COLOMBIA]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

MACROCRAMBUS Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 337.

Established unnecessarily as an objective replacement name for *Crambomorpha* Felder, 1874, which is not preoccupied by *Crambomorphus* M'Lachlan, [1867], *J. Linn. Soc.* (Zool.) 9: 243, – Insecta, Neuroptera.

CRATOPLASTIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.8. ARCT Type-species: Cratoplastis diluta Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.8, by monotypy. [BRAZIL]: Amazon R., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

CRATOSIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.8.

Type-species: *Cratosia parallela* Felder, 1874, *ibidem* 2 (Abt.2): pl.106, fig.8, by monotypy. Brazil: Cuyaba, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 3.

C. parallela is a junior subjective synonym of Cissura decora Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 489.

See also Cissura Walker, 1854.

‡CREAGA; misspelling. See Creagra Wallengren, 1865.

CREAGRA Wallengren, 1865, K. svenska VetenskAkad. Handl. (N.F.) 5 (4): 38.

Type-species: Creagra aliena Wallengren, 1865, ibidem (N.F.) 5 (4): 38, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 460, but cited as Liparis dealbata Herrich-Schäffer, [1854], a nominal species not originally included in Creagra. Kirby placed dealbata as the senior synonym of aliena, a nominal species originally included in Creagra. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as type-species. [SOUTH AFRICA]: Caffraria, eastern.

C. aliena is a junior subjective synonym of Orgyia liturata Guérin-Ménville, [1844] 1829-1838, Iconographie Règne ani-

mal de G. Cuvier (Insectes) 1: 511.

‡CREAGA; Janse, 1917, Check-List S. Afr. Lepid. Heterocera: 68.

An incorrect subsequent spelling of *Creagra* Wallengren, 1865.

CREATONOTOS Hübner, [1819] 1816, Verz. bekannter Schmett.: 170.

Type-species: *Phalaena interrupta* Linnaeus, 1767, *Syst. Nat.* (Edn 12) 1 (2): 840, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 241. Type-locality: not stated, [India and China to Australia].

P. interrupta is a junior subjective synonym of Phalaena gangis Linnaeus, 1763, Amoenitates Acad. 6: 410.

CREATONOTUS Agassiz, 1847, Nomencl. zool. (Index univl.): 103.

An unjustified emendation of Creatonotos Hübner, [1819].

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Creatonotus* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

CREATONOTUS Agassiz, 1847. See Creatonotos Hübner, [1819].

CREMNOPHORA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xvi, 7, 453. AGARISTIDAE

Type-species: Apina angasii Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 757, by original designation. AUSTRALIA: South.

Apina angasii Walker is a junior subjective synonym of Agarista angasii Angas, 1847, South Aust. illust.: pl.37, fig.24.

Cremnophora was established in the Arctiidae but is currently placed in the Agaristidae following Franclemont, 1973, (in litt.), Common, 1974 (in litt.) and Nye, 1975, Generic Names Moths World 1: 139.

CRESERA Schaus, 1894, Proc. zool. Soc. Lond. 1894: 232.

Type-species: Cresera annulata Schaus, 1894, ibidem 1894: 232, by monotypy. [Brazil]: Rio [de] Janeiro.

CRESTONICA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 60.

Type-species: *Desmeocraera circulosa* Gaede, 1928, *in* Seitz, *Gross-Schmett. Erde* 14: 414, pl.69 row g, by original designation. [ZAIRE] Congo: [upper] Lowa [valley, near Masisi].

CRINO Hübner, 1821. See Tarsolepis Butler, 1872. NOTO

CRINODES Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 91.

Type-species: *Phalaena bellatrix* Stoll, 1780, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* **4**: 32, pl.305, fig.F, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* **1**: 616. SURINAM.

Crinodes was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

ASTYLIS Boisduval, 1872, Annls Soc. ent. Belg. 15: 94.

Type-species: *Phalaena bellatrix* Stoll, 1780, by PRESENT DESIGNATION.

A junior objective synonym of *Crinodes* Herrich-Schäffer, 1855.

Astylis was established in the Adelocephalidae, now Saturniidae, Ceratocampinae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 616.

CRINOLA Leech, 1890. See Halseyella Nye nom. n. LYMA

CRINOPHORA Kaye, 1918, Ann. Mag. nat. Hist. (9) 1: 325.

Type-species: *Crinophora bicellulata* Kaye, 1918, *ibidem* (9) 1: 326, text-fig., by original designation. Philippines: Cebu.

CRISTULOSIA Toulgoët, 1958, Mém. Inst. scient. Madagas-car (E) 9: 199.

Type-species: Cristulosia bilunata Toulgoët, 1958, ibidem 9: 199, pl.2, fig.15, by original designation. MADAGASCAR: E., Périnet, Analamazaotra forest.

CRITOBULUS Kiriakoff, 1963. See Simesia Kiriakoff, 1955.

CROCODETA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxviii, 802.

Type-species: *Nudaria variegata* Rothschild, 1913, *Novit. zool.* **20**: 220, by original designation. New Guinea: [Papua New Guinea], N.E., Kumusi River.

CROCOMELA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 405.

Type-species: *Darna colorata* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 185, by monotypy (of *Darna* Walker, [1865]). [COLOMBIA] New Grenada: Bogota.

Crocomela was established as the objective replacement name for Darna Walker, [1865].

DARNA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 184.

Type-species: *Darna colorata* Walker, [1865], by monotypy. A junior homonym of *Darna* Walker, 1862, *J. Proc. Linn. Soc.* 6 (Zool.): 174, – Lepid., Limacodidae. The objective replacement name is *Crocomela* Kirby, 1892.

CROCOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xix, 470.

Type-species: *Crocosia phaeocraspis* Hampson, 1914, *ibidem* 1: 470, fig.91, by original designation. [GHANA] Gold Coast: Bibianaha.

CROCOTA Hübner, [1823] 1816, Verz. bekannter Schmett.: 295.

Type-species: Geometra tinctaria Hübner, [1799] 1796, Samml. eur. Schmett. 5: pl.23, fig.121, by subsequent designation by Grote, 1902, Allg. Z. Ent. 7: 472. EUROPE.

Crocota has been used as the original generic name for species now placed in the Arctiidae.

CROPERA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 825.

Type-species: *Cropera testacea* Walker, 1855, *ibidem* 4: 826, by monotypy. [SOUTH AFRICA]: [Durban] Port Natal.

CROPEROIDES Berio, 1940, Memorie Soc. ent. ital. 19: 125.

Type-species: *Croperoides negrottoi* Berio, 1940, *ibidem* 19: 125, by original designation. [ETHIOPIA]: [Negelli] Neghelli, Borana.

CROREMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 811.

Type-species: *Crorema mentiens* Walker, 1855, *ibidem* **4**: 811, by monotypy. Congo.

CROREMOPSIS Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 136.

Type-species: Croremopsis innocens Hering, 1926, ibidem 14: 136, pl.28, row b, by original designation. MADAGASCAR: Diego-Suarez.

The date of publication of the part of Seitz containing page 136 is cited on page 129 as 1.xi.1926 in both the German and the English editions.

‡CRUCIA; misspelling. See Grucia Dyar, 1916.

CTENANDRA Felder, 1861, Sher. Akad. Wiss. Wien 43 (Abt.1): 38.

Type-species: *Ctenandra naclioides* Felder, 1861, *ibidem* 43 (Abt.1): 39, by monotypy. [INDONESIA]: [Ambon I.] Amboina.

CTENANE Swinhoe, 1905, Ann. Mag. nat. Hist. (7) 15: 497.

Type-species: Ctenane labuana Swinhoe, 1905, ibidem (7) 15: 497, by monotypy. Borneo: [Sabah], Labuan.

CTENIA Le Peletier, 1825. See Spiris Hübner, [1819].

CTENOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 82, 130.

Type-species: *Pusiola psectriphora* Distant, 1899, *Ann. Mag. nat. Hist.* (7) **3**: 462, by original designation. South Africa: Transvaal, Pretoria.

CTENUCHA Kirby, 1837, in Richardson, Fauna boreal am. 4: 305.

Type-species: Ctenucha latreillana Kirby, 1837, ibidem 4:

305, by monotypy. Canada: Nova Scotia.

C. latreillana is a junior subjective synonym of Sphinx virginica Charpentier, 1830, in Esper, Ausl. Schmett. 2 (Suppl.): pl.2, figs 3, 4.

CTENUCHIDIA Grote, 1865, Proc. ent. Soc. Philad. 5: 227.

Type-species: Ctenucha virgo Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.54, fig.301, by original designation. Antill[ES] [including CUBA].

CTIANOPHA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 259 (key), 272.

Type-species: *Tiauspa argyria* Butler, 1879, *ibidem* 1879: 36, by original designation. [BRAZIL]: Amazon[a]s, Rio Jutahi.

CUAHTEMOCA Hogue, 1963, Contr. Sci. 64: 70.

Type-species: *Chalcopasta chalcocraspedon* Dyar, 1913, *Proc. U.S. natn. Mus.* 44: 297, by original designation. Holotype 9, MEXICO: Zacualpan, (USNM, Washington).

Cuahtemoca in the Noctuidae, Acronictinae is cited here as it was not included in Nye, 1975, Generic Names Moths World

1.

CUBIRA Watson nom. n. for Ira Neumoegen, 1890. CTEN Type-species: Ira gundlachiana Neumoegen, 1890, Entomologica am. 6: 64, by monotypy (of Ira Neumoegen, 1890). CUBA: S.E., Plantation Jagey.

IRA Neumoegen, 1890, Entomologica am. 6: 63.

Type-species: Ira gundlachiana Neumoegen, 1890, by

monotypy.

A junior homonym of *Ira* Walker, 1866, *List Specimens lepid. Insects Colln Br. Mus.* 35: 1562, – Lepid., Geometridae. The objective replacement name is *Cubira* Watson nom. n.

CUROBA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 272.

Type-species: *Curoba fasciata* Walker, [1865] 1864, *ibidem* 31: 272, by monotypy. TYPE-LOCALITY: not stated, [S.E. ASIA mainland].

C. fasciata is a junior subjective synonym of Phalaena sangarida Stoll, 1782, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 181, pl.381, fig.D.

CURUZZA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 152 (key), 171.

Type-species: *Pydna frugalis* Leech, 1898, *Trans. ent. Soc. Lond.* **1898**: 302, by original designation. CHINA: western, Putsu-fong, Moupin.

Curuzza was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 43.

CUTUZA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 153 (key), 168.

Type-species: Ceira straminea Moore, 1877, Ann. Mag. nat.

Hist. (4) 20: 91, by original designation. JAPAN.

Cutuza was established as a subgenus of Bireta Walker, 1856; it was placed as a junior subjective synonym of Torigea Matsumura, 1934, by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 40.

CYANA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 528.

Type-species: *Cyana detrita* Walker, 1854, *ibidem* **2**: 529, by monotypy. [BANGLADESH]: [Sylhet] Silhet.

MACRONOLA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 299.

Established, unnecessarily, as an objective replacement name for *Cyana* Walker, 1854, which is not preoccupied by *Cyane* Felder, 1861, *Nova Acta Acad. Caesar. Leop. Carol.* 28 (3): 22, – Lepid., Nymphalidae.

CYANARCTIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 6, 206.

Type-species: *Eupyra dama* Druce, 1894, *Ann. Mag. nat. Hist.* (6) 13: 170, by original designation. MEXICO: Orizaba.

CYANOHYPSA Giacomelli, 1911, An. Soc. cient. argent. 72:

Type-species: Cyanohypsa stefanellii Giacomelli, 1911, ibidem 72: 31, text-fig., by original designation. ARGENTINA: La Rioja.

CYANOPEPLA Clemens, 1861, Proc. Acad. nat. Sci. Philad. 1860: 545.

Type-species: Cyanopepla cruenta Clemens, 1861, ibidem 1860: 545, by monotypy. Mexico: near Jalapa.

C. cruenta is a junior subjective synonym of Glaucopis bella Guérin-Méneville, [1844], Icon. Règne Anim. Cuvier 3 (Insectes): 502.

AUTOMOLIS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 213.

Type-species: Glaucopis bella Guérin-Méneville, 1829-1838 [1844], Icon. Règne Anim. Cuvier 3 (Insectes): 502, by PRES-ENT DESIGNATION. MEXICO: Orizaba.

A junior homonym of *Automolis* Hübner, [1819] 1816, *Verz. bekannter Schmett.*: 170, — Lepid., Thyretidae. There is no objective replacement name, but *G. bella* is the senior subjective synonym of *Cyanopepla cruenta* Clemens, 1861, the type-species of *Cyanopepla* Clemens, 1861. The latter is therefore available for use as a subjective replacement name.

Automolis Walker was established as a subgenus of Euchromia Hübner, [1819].

CYANOTRICHA Prout, 1918, Novit. zool. 25: 397 (key), 426, (but attributed to Warren, an incorrect authorship).

Type-species: Sangala necyria Felder & Rogenhofer, 1875, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.133, fig.19, by original designation. PERU: Pebas.

CYBOSIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 167.

Type-species: *Noctua eborina* [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 68, by monotypy. [AUSTRIA]: Vienna district.

N. eborina is a junior subjective synonym of Phalaena mesomella Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 535.

See also Ecteina Rambur, 1866.

CYCETHRA Bethune-Baker, 1904. See Pseudodreata Bethune-Baker, 1904. ANTHELIDAE

CYCLOMILTA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 87, 512.

Type-species: *Miltochrista melanolepia* Dudgeon, 1899, *J. Bombay nat. Hist. Soc.* 13: 85, pl.2, fig.14, by monotypy (but incorrectly attributed to Hampson who redescribed this species as new in 1900, *ibidem* 2: 512). [INDIA]: Sikkim.

CYCLOSIELLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xx, 86, 548.

Type-species: *Hemonia dulcicula* Swinhoe, 1890, *Trans. ent. Soc. Lond.* **1890**: 183, by original designation. BURMA: Rangoon.

CYCLOSODES Hampson, 1901, Ann. Mag. nat. Hist. (7) 8: 183.

Type-species: Cyclosodes flavicostata Hampson, 1901, ibidem (7) 8: 184, by monotypy. Borneo: [Sarawak], Kuching.

CYCLOSTICTA Schaus, 1899, Jl N.Y. ent. Soc. 7: 217. LITH Type-species: Cyclosticta discata Schaus, 1899, ibidem 7: 217, by monotypy. [BRAZIL]: Rio [de] Janeiro.

CYCLOSTICTA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 87, 188.

Type-species: Cyclosticta discata Schaus, 1899, by original designation.

A junior homonym of *Cyclosticta* Schaus, 1899. This generic name probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal and generic description by Hampson, 1900. The objective replacement name is *Cyclosticta* Schaus, 1899.

CYCLOSTICTA Hampson, 1900. See Cyclosticta Schaus, 1899.

CYCNIA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 7.

Type-species: Cycnia tenera Hübner, 1818, ibidem 1: 7, pl.[1], figs 3, 4, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 237. [U.S.A.]: Georgia [then part of Florida].

See also Tanada Walker, 1856.

‡HYPERCOMPE Hübner, 1808, Erste Zuträge Samml. exot. Schmett.: [3], 7.

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1966, Bull. zool. Nom. 23 (Opinion 789): 213. Placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1862.

Only included species: ‡Hypercompe tenera Hübner, 1808, ibidem: [3], 7, later established as Cycnia tenera Hübner, 1818.

CYDOSIA Duncan [& Westwood], 1841, in Jardine, Naturalist's Library (Edn 1) 33 (Ent. 7): 193.

NOCTUIDAE

Type-species: *Phalaena nobilitella* Cramer, 1779, *Uitlandsche Kapellen (Papillons exot.)* 3: 128, pl.264, fig.G, by monotypy. [West Indies]: Curação [I.].

Cydosia has been used as the original generic name for species now in the Arctiidae.

CYLLENE Walker, 1854. See Lyclene Moore, 1858.

CYMAROA Hampson, 1905, Ann. S. Afr. Mus. 3: 410. ARCT Type-species: Cymaroa leptopepla Hampson, 1905, ibidem 3: 411, by original designation. [LESOTHO] Basutoland, SOUTH AFRICA: Cape Colony, Grahamstown.

Cymaroa was established in the Lymantri[i]dae; it is here transferred to the Arctiidae, Arctiinae, on the advice of, and with acknowledgements to Dr L. Vari, Transvaal Museum, South Africa.

CYMBALOPHORA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 231.

Type-species: *Phalaena pudica* Esper, 1785, *Die Schmett.* 3: 177, pl.33, fig.1, by monotypy. [France]: [Savoie mountains] savoyischen Gebürgen.

P. pudica is dated from the wrapper of Heft 20 and not from the title-page of the volume, 1782.

TYMPANOPHORA Laboulbène, 1865, Annls Soc. ent. Fr. (4) 4: 704

Type-species: *Phalaena pudica* Esper, 1785, by monotypy. A junior homonym of *Tympanophora* White, 1841, *in* Grey, *J. Exped. Austral.* 2: 468, – Insecta, Orthoptera. The objective replacement name is *Cymbalophora* Rambur, 1866.

CYME Felder, 1861, Sher. Akad. Wiss. Wien 43 (Abt.1): 36.

Type-species: Cyme reticulata Felder, 1861, ibidem 43 (Abt.1): 36, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 300. [INDONESIA]: [Ambon I.] Amboina.

CYMELLA Felder, 1874. See Myclela Watson nom. n. LITH

CYMOPSIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.23.

Type-species: Cymopsis fulviceps Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.23, by monotypy. [BRAZIL]: River Amazon, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

CYNOSARGA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 385.

Type-species: Cynosarga ornata Walker, 1865, ibidem 32: 386, by monotypy. [Australia]: [Queensland], Moreton Bay.

Cynosarga was established in the Liparidae, now Lymantridae; it was placed in the Limacodidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 528, and by Hering, 1931, in Seitz, Gross-Schmett. Erde 10: 720; it was placed in the Notodontidae, Cnethocampinae, now Thaumetopoeidae by Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 373; and its placing in this family was confirmed by Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 34.

CYPHOCAMPA Harris, 1869, in Scudder, Occ. Pap. Boston Soc. nat. Hist. 1: 138.

AGARISTIDAE

Type-species: *Bombyx grata* Fabricius, 1793, *Ent. Syst.* 3 (1): 457, by monotypy. Type(s) [U.S.A.]: Georgia.

Cyphocampa in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PARATHISANOTIA Kiriakoff, 1976, Tierreich 99: 76.

Type-species: Bombyx grata Fabricius, 1793, by original designation.

Parathisanotia is a junior objective synonym of Cyphocampa Harris, 1869.

Parathisanotia is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

CYPHOPORA Warren, 1900, Novit. zool. 7: 125. PERI Type-species: Cyphopora dubia Warren, 1900, ibidem 7: 126, by original designation. ECUADOR: Paramba.

Cyphopora Warren was established in the Geometridae; it was transferred to the Arctiidae, Pericopinae, by Fletcher, 1979, in Nye, Generic Names Moths World 3: 56.

CYPRA Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 201. GEOMETRIDAE

Type-species: Cypra delicatula Boisduval, 1832, ibidem 1:

201, pl.3, fig.1, by monotypy. New Guinea.

Cypra was not placed in a family when it was established; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 437; and was transferred to the Boarmiidae, now Geometridae Ennominae, by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497.

Druce, 1899, Ann. Mag. nat. Hist. (7) 3: 469 established 'Cypra (?) Forbesi, sp.n.' in the 'Liparidae' and this usage of Cypra has incorrectly been cited as an available name by Neave,

1939, Nomencl. zool. 1: 942.

CYPTONYCHIA Druce, 1899, Ann. Mag. nat. Hist. (7) 3: 467.

Type-species: Cyptonychia flaviceps Druce, 1899, ibidem (7) 3: 467, by monotypy. Type(s) 8, MEXICO, (UM, Oxford).

C. flaviceps is a junior subjective synonym of Eulithosia thoracica Edwards, 1884, Papilio 4: 44, the type-species of

Chichimeca Hogue, 1963.

Cyptonychia was established in the Lithosiidae, now Arctiidae, Lithosiinae; the type of Cyptonychia flaviceps Druce, 1899, has been examined and this nominal species and its genus is here transferred to the Noctuidae, Acronictinae, and placed near Cirrhophanus Grote, 1872.

See also Chichimeca Hogue, 1963.

CYPTONYCHIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 92, 253.

Type-species: Cyptonychia flaviceps Druce, 1899, by original

designation.

A junior homonym of *Cyptonychia* Druce, 1899. This generic name probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal and generic description by Hampson, 1900. The objective replacement name is *Cyptonychia* Druce, 1899.

CYPTONYCHIA Hampson, 1900. See Cyptonychia Druce, 1899.

CYRRHESTA Walker, 1857, List Specimens lepid. Insects Colln Br. Mus. 11: 611 (key), 633.

Type-species: *Phalaena nyseus* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 119, pl.75, fig.E, by monotypy. SURINAM.

P. nyseus is dated from the wrapper of part 7, not from the

title-page of volume 1 which is dated 1779.

Cyrrhesta was established in the Noctuidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 617.

CYTORUS Grote, 1865, Proc. ent. Soc. Philad. 5: 232. ARCT Type-species: Cytorus latus Grote, 1865, ibidem 5: 232, pl.4, fig.1, by monotypy. CUBA.

DACTYLORHYNCHA Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 470.

Type-species: Charnides pallida Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 56, pl.140, fig.10, by original designation. INDIA: southern, Nilgiri Hills.

Dactylorhyncha was established in the Lymantriidae; the holotype of C. pallida and one other specimen, both in BMNH, were transferred by C.L. Collenette to the Limacodidae, but no published reference to the transfer has been found.

DACTYLORHYNCHIDES Strand, 1920, Arch. Naturgesch. 84 (A) 12: 185.

Type-species: *Dactylorhynchides limacodiformis* Strand, 1920, *ibidem* 84 (A) 12: 185, by monotypy.

Dactylorhynchides was established in the Lymantriidae; it was transferred to the Limacodidae by Collenette, 1931, Ann. Mag. nat. Hist. (10) 7: 512.

DAHANA Grote, 1875, Can. Ent. 7: 175. CTEN Type-species: Dahana atripennis Grote, 1875, ibidem 7: 175, by monotypy. U.S.A.: Florida, Enterprise.

‡DEHANA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420

An incorrect subsequent spelling of Dahana Grote, 1875.

DAMALIS Hübner, [1819]. See Psephea Billberg, 1820.

DAMATA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1044.

Type-species: *Damata longipennis* Walker, 1855, *ibidem* 5: 1044, by monotypy. HIMALAYAS.

DAMATOIDES Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19: 13.

Type-species: *Damatoides baibarana* Matsumura, 1927, *ibidem* 19: 13, pl.1, fig.3, by original designation. TAIWAN: Horisha, Baibara.

D. baibarana is a junior subjective synonym of Damata microsticta Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 157.

See also Macrohoplitis Roepke, 1943.

DAMIAS Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 259.

Type-species: *Damias elegans* Boisduval, 1832, *ibidem* 1: 259, fig., by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 397. New Guinea: [Papua New Guinea] Papous et la Nouvelle Guinée.

Damias was established in the Castnii, now Castniidae; it was placed in the Callidulidae by Pagenstecher, 1887, Jb. nassau. Ver. Naturk. 40: 219; and was placed in the Arctiidae Lithosiinae by Strand, 1922, in Wagner, Lepid. Cat. 26: 667.

Damias has been used incorrectly by Kiriakoff, 1976, Tierreich 97: 79, as a valid name in the Agaristidae. Kiriakoff based his usage on '1854 Damias Walker, List Lepidopt. Het. Brit. Mus., 1: 14. – Species typica: Damias varia Walker (by monotypy)'. Walker had, however, correctly attributed Damias to Boisduval and used this generic name for four species, not one, the first being D. elegans.

See also Caprimima Hampson, 1900.

DANAKA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 71.

PYRALIDAE

Type-species: *Danaka pyraliformis* Moore, 1879, *ibidem* (1): 71, pl.3, fig.10, by monotypy. INDIA: Darjeeling.

Danaka was established in the Notodontidae; it was placed in the Pyralidae by Hampson, 1896, Fauna Br. India (Moths) 4: 169.

‡DANIELA Hartig, 1963. See Parocneria Dyar, 1897.

DANIELITA Kiriakoff, 1970. See Pterotes Berg, 1901.

DANIMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1052.

Type-species: *Bombyx banksiae* Lewin, 1805, *Prodromus Ent.*, *nat. Hist. lepid. Insects N.S.W.*: 10, pl.9, by monotypy. [AUSTRALIA]: New South Wales.

DAPHAENISCA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 69.

Type-species: *Pseudapiconoma daphaena* Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 150, fig.62, by original designation. [NIGERIA]: R. Niger, Asaba.

Daphaenisca was established as a subgenus of Balacra Walker, 1856.

DAPHNE Neumoegen, 1894. See Didaphne Berg, 1899.

DAPLASA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 51.

Type-species: *Daplasa irrorata* Moore, 1879, *ibidem* (1): 52, pl.2, fig.17, by monotypy. INDIA: [W. Bengal], [Darjeeling] Darjiling.

DARABITTA Walker, 1859, J. Proc. Linn. Soc. (Zool.) 3: 187.

Type-species: *Darabitta strigicosta* Walker, 1859, *ibidem* (Zool.) 3: 187, by monotypy. SINGAPORE.

DARANTASIA Walker, 1859, J. Proc. Linn. Soc. (Zool.) 3: 186.

Type-species: *Darantasia cuneiplena* Walker, 1859, *ibidem* (Zool.) 3: 186, by monotypy. SINGAPORE.

DARANTASIELLA Roepke, 1943, Natuurh. Maandbl. 32: 78.

Type-species: *Darantasiella javanica* Roepke, 1943, *ibidem* 32: 78, fig.6, by monotypy. Java: E., Djunggo-Ardjuno.

DARANTOIDES Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xx, 92, 565.

Type-species: *Darantoides rubroflava* Hampson, 1900, *ibidem* 2: 566, pl.35, fig.13, by original designation. New Guinea: [Papua New Guinea], Milne Bay.

DARGEOCHAETA Laporte, 1973, Bull. Mus. natn. Hist. nat. Paris (3) Zool. (126): 1492.

Type-species: *Dargeochaeta bangeni* Laporte, 1973, *ibidem* (3) **Zool**. (126): 1492, fig. 7, by original designation. Holotype &, CAMEROUN: massif du Manengouba, piste de Bangen par Mwakoumel, (MNHN, Paris).

DARITIS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 618.

Type-species: *Eyprepia thetis* Klug, 1836, *Neue Schmett*. (1): 5, pl.4, figs 1, 2 (but cited as ‡*Euprepia*, an incorrect subsequent spelling), by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 186. MEXICO.

E. thetis is a junior subjective synonym of Eucharia marianne Geyer, [1835], in Hübner, Samml. exot. Schmett. 3:

pl.[46].

DORIMENA Boisduval, 1870, Considérations Lépid. Guatemala: 98.

Type-species: Eyprepia thetis Klug, 1836 (but cited as ‡Euprepia, an incorrect subsequent spelling), by subsequent designation by Bryk, 1931, in Strand, Lepid. Cat. 45: 5.

A junior objective synonym of Daritis Walker, 1855.

‡DORIMENE; Bryk, 1931, in Strand, Lepid. Cat. 45: 5.

An incorrect subsequent spelling of *Dorimena* Boisduval, 1870.

DARNA Walker, [1865]. See Crocomela Kirby, 1892. PERI

DASIPPIA Draudt, 1932, in Seitz, Gross-Schmett. Erde 6: 988.

Type-species: Dasippia didugana Draudt, 1932, ibidem 6:

988, pl.152 row b, by original designation. BRAZIL: southern, Sa[nta] Cruz.

The date of publication of the part of Seitz containing page 988 is cited on page 985 as 5.x.1932 in both the German and the English editions.

DASORGYIA Staudinger, 1881, Stettin. ent. Ztg 42: 406.

Type-species: *Dasychira pumila* Staudinger, 1881, *ibidem* **42**: 405, by monotypy. [U.S.S.R.]: [Kazakhstan], [Zaysan] Saisan.

Dasorgyia was proposed at the end of the original description of D. pumila in case it should require a new genus.

‡DASUNAGA; misspelling. See Nagidusa Walker, 1862.

DASYARCTIA Gaede, 1923, Ent. Rdsch. 40: 28.

Type-species: *Dasyarctia grisea* Gaede, 1923, *ibidem* **40**: 29, by original designation. [SOUTH AFRICA]: [Cape Province] Kapland, Hopefield.

Dasyarctia was established in the Arctiinae; it was placed in the Lymantriidae by Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 164; it is here retained in the Arctiinae.

DASYCAMPA Janse, 1915. See Hemerophanes Collenette, 1953.

‡DASYCHIRA Hübner, [1806]. See Elkneria Börner, 1932.

DASYCHIRA Hübner, [1809], Samml. exot. Schmett. 1: pl.[178].

Type-species: Dasychira tephra Hübner, [1809], ibidem 1: pl.[178], by monotypy. Type-locality: not stated. [U.S.A.].

DASYCHIRANA Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 550.

Type-species: *Dasychirana obliqualinea* Bethune-Baker, 1911, *ibidem* (8) 7: 551, by original designation. [ANGOLA]: Malange.

‡DASYCHIRINA; Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 411.

An incorrect subsequent spelling of *Dasychirana* Bethune-Baker, 1911.

DASYCHIRELLA Kiriakoff, 1970, Ann. Transv. Mus. 26 (9): 193.

Type-species: *Dasychirella cinderella* Kiriakoff, 1970, *ibidem* **26** (9): 194, fig. 14, by original designation. South Africa: Natal National Park, Mont-aux-Sources.

‡DASYCHIRINA; misspelling. See Dasychirana Bethune-Baker, 1911.

DASYCHIRINULA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 187. LASIOCAMPIDAE

Type-species: *Dasychirinula chrysogramma* Hering, 1926, *ibidem* 14: 187, pl.28, rows c, d, by monotypy. [TANZANIA]: Kilimatinde, [Kenya]: Kibwezi (not Kilawesi as in the English edition).

The date of publication of the part of Seitz containing page 187 is cited on page 185 as 11.xi.1926 in both the German and the English editions.

Dasychirinula was established in the Lymantriidae; the female syntype from Kibwezi and one other specimen, both in BMNH, were transferred by C.L. Collenette to the Lasiocampidae, but no published reference to the transfer has been found.

DASYCHIROIDES Bethune-Baker, 1904, Novit. zool. 11: 405.

Type-species: *Dasychiroides obsoleta* Bethune-Baker, 1904, *ibidem* 11: 406, pl.6, fig.43, by original designation. New Guinea: [Papua New Guinea], Owen Stanley Range, Dinawa.

DASYCHOPROCTIS Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 151.

Type-species: *Dasychoproctis dubiosa* Hering, 1926, *ibidem* 14: 151, pl.22, row f, by monotypy. MADAGASCAR: Betsileo.

The date of publication of the part of Seitz containing page 151 is cited on page 145 as 4.xi.1926 in both the German and the English editions.

DASYLAELIA Aurivillius, 1925, Ergebn. zweiten dt. zent.-Afr. Exped. 1910-1911 (Zool.) 1 (18): 1284.

Type-species: *Dasylaelia strigata* Aurivillius, 1925, *ibidem* 1: 1285, fig.68, by monotypy. [CAMEROUN] Kamerun: S., Bundji.

DASYLOPHIA Packard, 1864, Proc. ent. Soc. Philad. 3: 362.

Type-species: *Phalaena anguina* Smith, 1797, in Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 167, pl.84, by subsequent designation by Grote, 1895, *Abh. naturw. Ver. Bremen* 14: 48. [U.S.A.]: Georgia.

DASYSPHINX Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.25. CTEN

Type-species: Dasysphinx mucescens Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.25, by monotypy. [COLOMBIA]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

DATANA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1060.

Type-species: *Phalaena ministra* Drury, 1773, *Illust. nat. Hist. exot. Insects* 2: 25, Index [1], pl.14, fig.3, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 613. [U.S.A.]: New York.

EUMETOPONA Fitch, 1856, Trans. N.Y. St. agric. Soc. 15: 467, 473.

Type-species: *Phalaena ministra* Drury, 1773, by monotypy. A junior objective synonym of *Datana* Walker, 1855.

Eumetopona was also published by Fitch, 1856, Rep. noxious, beneficial and other Insects, State N.Y. 2: 235. Neither Dr D. Ferguson of the United States National Museum nor Dr J. Franclemont of Cornell University, Ithaca, U.S.A., has been able to establish the precise dates of publication of the two Fitch works cited above. The sequence adopted in this catalogue follows that used by Franclemont, 1973, in Dominick et al., Moths Am. N. of Mexico 20 (1): 33, 38.

DAULOPAECTES Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) **66**: 68.

Type-species: Stauropus trichosa Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 471, by original designation. [GHANA] Gold Coast: Kumasi.

DECIMIA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1717.

Type-species: *Decimia bicolora* Walker, 1856, *ibidem* 7: 1718, by monotypy. [South Africa]: [Durban] Port Natal.

Decimia was established in the Arctiidae; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 30.

DEDIAMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1074.

Type-species: *Dediama antica* Walker, 1855, *ibidem* 5: 1074, by monotypy. Type-locality: not known, [Cameroun].

Dediama was established in the Notodontidae; it was placed in the Lymantriidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 464.

See also Bathmochtha Karsch, 1895.

‡DEHANA; misspelling. See Dahana Grote, 1875. CTEN

DEILEMERA Hübner, [1820] 1816, Verz. bekannter Schmett.: 179.

Type-species: *Phalaena evergista* Stoll, 1781, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 155, pl.369, fig.E, by monotypy. [INDONESIA]: [Ambon 1.] Amboina.

P. evergista is dated from the wrapper of the part and not from the title-page of the volume, 1782.

DILEMERA Agassiz, 1847, Nomencl. zool. (Index univl.): 118, 124.

An unjustified emendation of Deilemera Hübner, [1820]

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Dilemera* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

DEINARCHIA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 52.

Type-species: *Dasychira apateloides* Holland, 1893, *Psyche, Camb.* 6: 434, pl.18, fig.1, by original designation. [GABON]: [Ogooué] Ogové R. valley.

DEINHUGIA Laporte, 1974, Bull. Soc. ent. Fr. 79: 211.
NOCTUIDAE

Type-species: *Deinhugia nigra* Laporte, 1974, *ibidem* 79: 212, fig.12, by original designation. Holotype & CAMEROUN: [W.], Massif du Kala, Mont Nkolbiyong, (MNHN, Paris).

Deinhugia in the Noctuidae, Ophiderinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡DEIOPEA; misspelling. See Deiopeia Curtis, 1827. ARCT

DEIOPEIA Curtis, 1827, Br. Ent. 4: 169.

Type-species: Phalaena pulchella Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 534, by original designation. EUROPE: southern, [ALGERIA/ MOROCCO/ TUNISIA] Mauritania.

DEIOPEIA Stephens, 1829 February, Illust. Br. Ent. (Haustellata) 2: 89, 92.

Type-species: Phalaena pulchella Linnaeus, 1758, by

monotypy.

A junior homonym of *Deiopeia* Curtis, 1827. This generic name originated from Stephens but was described and made nomenclaturally available by Curtis prior to its proposal and generic description by Stephens, 1829. The objective replacement name is *Deiopeia* Curtis, 1827.

Deiopeia was proposed again by Stephens, 1829 [June], Nom. Br. Insects: 40; and 1829 [July], Syst. Cat. Br. Insects (2): 60.

‡DIOPEA; Burmeister, 1878, Descr. phys. Répub. Argent. 5 (1): 431.

An incorrect subsequent spelling of Deiopeia Curtis, 1827.

‡DEIOPEJA; Mabille, 1879, Naturaliste 1 (3): 21.

An incorrect subsequent spelling of Deiopeia Curtis, 1827.

‡DEJOPEIA; Mabille, 1879, Bull. Soc. zool. Fr. 3: 88.
An incorrect subsequent spelling of Deiopeia Curtis, 1827.

‡DEIOPEA; Mabille, 1880, Annls Soc. ent. Fr. (5) 9: 305.
An incorrect subsequent spelling of Deiopeia Curtis, 1827.

DEIOPEIA Stephens, 1829. See Deiopeia Curtis, 1827.

‡DEIOPEJA; misspelling. See Deiopeia Curtis, 1827.

‡DEJOPEIA; misspelling. See Deiopeia Curtis, 1827.

DELOPLOTELA Hulstaert, 1924, Ann. Mag. nat. Hist. (9) 13: 132.

Type-species: Deloplotela minuta Hulstaert, 1924, ibidem (9) 13: 132, by original designation. [INDONESIA]: [Tanimbar Is] Tenimbar, Olilit.

DELPHYRE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 537. CTEN

Type-species: Delphyre hebes Walker, 1854, ibidem 2: 537, by monotypy. HONDURAS.

See also Nodoza Schaus, 1896.

DEMOLIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus.

Type-species: Demolis albicostata Hampson, 1901, ibidem 3: 31, fig.24, by original designation. BRAZIL: Rio Grande do

DENDROPHLEPS Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 491.

Type-species: Dendrophleps semihyalina Hampson, [1893] 1892, ibidem 1: 491, fig. 326, by original designation. INDIA: [Assam], Khasi Hills.

DENSITAS Marumo, 1920. See Epizaranga Matsumura, 1920.

DERIDDERA Kiriakoff, 1955, Revue Zool. Bot. afr. 52: 334.

Type-species: Deriddera margarethae Kiriakoff, 1955, ibidem 52: 335, fig.10, pl.2, fig.10, by original designation. [ZAIRE] Congo Belge: Sankaru, Katako-Kombe.

DEROCA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key), 822. DREPANIDAE

Type-species: Deroca hyalina Walker, 1855, ibidem 4: 823,

by monotypy. Himalaya Mountains.

Deroca was established in the Liparidae, now Lymantriidae; it was placed in the 'Drepanulidae' by Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 344.

DERRHIS Wallengren, 1885. See Nudaria Haworth, [1809].

DESMEOCRAERA Wallengren, 1865, K. svenska Vetensk-Akad. Handl. (N.F.) 5 (4): 52.

Type-species: Stauropus interpellatrix Wallengren, 1860, Wien. ent. Monatschr. 4: 164, by monotypy. [South Africa]: Caffraria.

DESMEOCRAERA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.94, fig.9.

Type-species: Desmeocraera nugatrix Felder, 1874, ibidem 2 (Abt.2): pl.94, fig.9, by monotypy. [South Africa] Afr. mer.: Knysna, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

A junior homonym of Desmeocraera Wallengren, 1865, K. svenska VetenskAkad. Handl. (N.F.) 5 (4): 52. There is no objective replacement name, but D. nugatrix is believed to be congeneric with Stauropus interpellatrix Wallengren, 1860, the type-species of Desmeocraera Wallengren, 1865; the latter is thus available for use as a subjective replacement name.

DESMEOCRAERA Felder, 1874. See Desmeocraera Wallengren, 1865.

DESMEOCRAERULA Strand, 1912, Arch. Naturgesch. 78 (A) 6: 168.

Type-species: Desmeocraerula inconspicuana Strand, 1912,

ibidem 78 (A) 6: 169, by original designation. EQUATORIAL GUINEA: Rio Muni.

DESMIDOCNEMIS Möschler, 1872, Stettin. ent. Ztg 33: 346.

Type-species: Desmidocnemis prittwitzi Möschler, 1872, ibidem 33: 348, by monotypy. [French Guianal: Cavenne.

Desmidocnemis was attributed to Hampson (an incorrect authorship) by Zerny, 1912, in Wagner, Lepid. Cat. 7: 106; and by Neave, 1939, Nomencl. zool. 2: 49. Some confusion has resulted from Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 429, who stated 'Page 363. Desmidocnemis rename Hyalomis, n. n., the former being a synonym of HEREA and Prittwitzi = xanthogaster' Druce, 1898. Hampson was not, however, proposing Hyalomis as a replacement name for Desmidocnemis but was establishing a new genus for species placed incorrectly in Desmidocnemis Möschler, in Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 363.

DESMOLOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.99, fig.22. Type-species: Desmoloma styracis Felder, 1874, ibidem 2 (Abt.2): pl.99, fig.22, by monotypy. VENEZUELA: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

DESMOTRICHA Hampson, 1911. See Epidesma Hübner, [1819].

DESTOLMIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 991.

Type-species: Destolmia lineata Walker, 1855, ibidem 5: 992, by monotypy. [AUSTRALIA]: [New South Wales], Sydney, [Western Australia], Swan River.

See also Collyta Walker, 1865.

DEUA Walker, 1864, List Specimens lepid. Insects Colln Br. Mus. 30: 1013.

Type-species: Deua imbutana Walker, 1864, ibidem 30: 1013, by monotypy. [Brazil]: [R. Amazon], [Tefé] Ega.

DIABAENA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.103, fig.1. Type-species: Diabaena cincticollis Felder, 1874, ibidem 2 (Abt.2): pl.103, fig.1, by monotypy. [Colombia]: Bogota, cited

by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

DIACONISIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxvii, 778.

Type-species: Melanaema ochraceorufa Rothschild, 1913, Novit. zool. 20: 203, by original designation. New Guinea: [Irian Jaya], Snow Mts, near R. Oetakwa.

DIACRISIA Hübner, [1819] 1816, Verz. bekannter Schmett.:

Type-species: Phalaena russula Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 510, by monotypy. EUROPE.

P. russula is a junior subjective synonym of Phalaena sannio Linnaeus, 1758, ibidem 1: 506.

EUTHEMONIA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 55 (key), 68.

Type-species: Phalaena russula Linnaeus, 1758, by monotypy.

A junior objective synonym of Diacrisia Hübner, [1819].

Euthemonia was proposed again by Stephens, 1829 [June], Nom. Br. Insects: 40; and 1829 [July], Syst. Cat. Br. Insects (2): 54.

ELPIS Dyar, 1893, Ent. News 4: 36.

Type-species: Antarctia rubra Neumoegen, 1881, Papilio 1: 79, by original designation. U.S.A.: Oregon, Mount Hood.

A junior homonym of Elpis Mulsant, 1850, Annls Soc. Agric. Lyon [2] 2: 449, - Insecta, Coleoptera. There is no objective

replacement name but the type-species of *Elpis* Dyar is currently considered to be congeneric with the type-species of *Diacrisia* Hübner, [1819]. The latter is therefore available for use as a subjective replacement name.

DIADESMOLA Hulstaert, 1924, Annls Soc. ent. Belg. 64: 87.

Type-species: *Diadesmola bicolor* Hulstaert, 1924, *ibidem* 64: 88, by original designation. New Guinea: [Irian Jaya], [upper R. Merauke] région de la Haute Merauke, Toerai.

DIAKONOFFIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 20, 22. THYRET

Type-species: Apisa kivensis Dufrane, 1945, Bull. Annls Soc. ent. Belg. 81: 128, by original designation. [ZAIRE]: Kivu, Kamituga.

DIALEPHTIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.14.

Type-species: *Dialephtis salvini* Felder, 1874, *ibidem* 2 (Abt.2): pl.105, fig.14, by monotypy. GUATEMALA: cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 4.

BELCIANA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 156.

Type-species: Belciana strigularis Walker, [1865] 1864,

ibidem 31: 157, by monotypy. [COLOMBIA]: Bogota.

A junior homonym of *Belciana* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 182, – Lepid., Noctuidae. There is no objective replacement name, but *B. strigularis* is a junior subjective synonym of *Dialephtis semimarginata* Dognin, 1902, *Annls Soc. ent. Belg.* 46: 342, currently treated as congeneric with *Dialephtis salvini* Felder, 1874, the type-species of *Dialephtis* Felder, 1874; the latter is thus available for use as a subjective replacement name.

Neave, 1939, Nomencl. zool. 1: 412, suggested incorrectly that Nalca Walker, 1866, was the objective replacement name for Belciana Walker, [1865]; Nalca was, however, proposed unnecessarily as an objective replacement name for Belciana

Walker, 1862.

‡DIALEPLETIS; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 405.

An incorrect subsequent spelling of Dialephtis Felder, 1874.

‡DIALEPLETIS; misspelling. See Dialephtis Felder, 1874.

DIALEUCIAS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: ix, 3, 97.

Type-species: *Dialeucias pallidistriata* Hampson, 1901, *ibidem* 3: 98, fig.73, by original designation. [BRAZIL]: [Amazonas], Olivenças.

DIAPHORA Stephens, 1827, in Anonymous, Retrospective Review (2) 1: 244.

Type-species: *Phalaena mendica* Clerck, 1759, *Icon. Insect. rariorum* 1: pl.3, fig.5, by monotypy. Type-Locality: not stated, [Europe].

Diaphora was proposed three more times by Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 77; Stephens, 1829 [June], Nom. Br. Insects: 40; and Stephens, 1829 [July], Syst. Cat. Br. Insects (2): 57.

DIARHABDOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 85, 517.

Type-species: Eustixis minima Butler, 1878, Trans. ent. Soc. Lond. 1878: 55, by subsequent designation by Collins, 1962, Ann. Mag. nat. Hist. (13) 5: 125. [BRAZIL]: [Amazonas], Rio Jutahi.

DIASTEMA Herrich-Schäffer, [1855]. See Antheua Walker, 1855.

DIASTEMA Aurivillius, 1904. See Antheua Walker, 1855.

DIASTEMINA Gaede, 1928. See Antheua Walker, 1855.

DIASTROPHIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.13. LITH

Type-species: *Diastrophia dasypyga* Felder, 1874, *ibidem* 2 (Abt.2): pl.106, fig.13, by monotypy. [SULAWESI]: Celebes, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 4.

D. dasypyga Felder is a junior subjective synonym of Teulisna chiloides Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 109. See also Corcura Walker, 1862, and Ityca Walker, 1866.

DIAXANTHIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xix, 28, 436.

Type-species: *Automolis lucinia* Druce, 1884, *Biologia cent.-am.* (Zool.) Lcpid.-Heterocera 1: 75, fig., by original designation. Costa Rica: Irazu.

DICALLOMERA Butler, 1881, Trans. ent. Soc. Lond. 1881: 12.

Type-species: *Phalaena fascelina* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 503, by subsequent designation by Collenette, 1935, *Stylops* 4: 243. Type-locality: not stated, [Europe].

DICENTRIA Herrich-Schäffer, 1855 December, Syst. Bearbeitung Schmett. Eur. 6: 91.

Type-species: *Notodonta centralis* Herrich-Schäffer, [1855 June], *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.67, fig.383, by monotypy. [Brazil]: Rio [de Janeiro].

Dicentria was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

DICERATOSTAURA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 23.

Type-species: *Eurystaura fuscata* Talbot, 1929, *Bull. Hill Mus. Witley* 3: 65, pl.2, fig.11, by original designation. São Tomé Island, W. Africa.

DICERATUCHA Swinhoe, 1904, Ann. Mag. nat. Hist. (7) 14: 133.

Type-species: *Oenone xenopis* Lower, 1902, *Trans. Proc. R. Soc. S. Aust.* **26**: 277, by original desognation. Australia: New South Wales, Broken Hill.

Diceratucha was established in the Geometridae; it was transferred to the Thaumetopoeidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

DICHROSTOPTERA Hulstaert, 1924, Annls Soc. ent. Belg. 64: 86.

Type-species: *Dichrostoptera basilinea* Hulstaert, 1924, *ibidem* 64: 87, by original designation. New Guinea: [Irian Jaya], [R. Digul] Digoel.

DICRANOURA Berthold, 1827. See Dicranura R.L., 1817.

NOTO

DICRANURA R.L., 1817, Jena. allg. Lit.-Ztg 1 (36): 282.

Type-species: Noctua ulmi [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 66, by monotypy (but attributed to Borkhausen, an incorrect authorship). [AUSTRIA]: Vienna district.

EXAERETA Hübner, [1820] 1816, Verz. bekannter Schmett.: 200.

Type-species: *Noctua ulmi* [Denis & Schiffermüller], 1775, by monotypy.

A junior objective synonym of Dicranura R.L., 1817.

‡EXERETA; Hübner, [1826] 1816, Verz. bekannter Schmett. (Anz.): 38.

An incorrect subsequent spelling of Exaereta Hübner, [1820].

DICRANOURA Berthold, 1827, in Latreille, Nat. Familien Thierreichs: 479.

Type-species: *Noctua ulmi* [Denis & Schiffermüller], 1775, by monotypy.

A junior objective synonym of Dicranura R.L., 1817.

UROPUS Rambur, 1832, Annls Soc. ent. Fr. 1 (3): 278.

Type-species: *Noctua ulmi* [Denis & Schiffermüller], 1775, by monotypy.

A junior objective synonym of *Dicranura R.L.*, 1817.

‡EXCERATA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 428.

An incorrect subsequent spelling of Exaereta Hübner, [1820].

DICRANURA Boisduval, 1828. See Cerura Schrank, 1802.

DICRANUROPSIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.100, fig.7.

Type-species: *Dicranuropsis vilis* Felder, 1874, *ibidem* 2 (Abt.2): pl.100, fig.7, by monotypy. [SOUTH AFRICA] Afr. m.: Knysna, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 4.

‡DICTA; misspelling. See Diota Wallengren, 1865. ARCT

DICTENUS Butler, 1897, Proc. zool. Soc. Lond. 1896: 846.

Type-species: *Dictenus inconstans* Butler, 1897, *ibidem* **1896**: 846, pl.42, fig.5, by monotypy. [MALAWI]: Nyika, Mt Kasangu.

DIDAPHNE Berg, 1899, Comun. Mus. nac. B. Aires 1: 78.

Type-species: *Daphne cyanomela* Neumoegen, 1894, *Can. Ent.* **26**: 334, by monotypy (of *Daphne* Neumoegen, 1894). CUBA.

Didaphne was proposed as the objective replacement name for Daphne Neumoegen, 1894.

DAPHNE Neumoegen, 1894, Can. Ent. 26: 334.

Type-species: *Daphne cyanomela* Neumoegen, 1894, by monotypy.

A junior homonym of *Daphne* Müller, 1776, *Zoologiae Danicae Prodromus*: xxvii, 199, – Crustacea. The objective replacement name is *Didaphne* Berg, 1899.

DIDASYS Grote, 1875, Can. Ent. 7: 174. CTEN Type-species: Didasys belae Grote, 1875, ibidem 7: 175, by monotypy. U.S.A.: Florida, Cedar Keys.

DIDUGA Moore, [1877] 1884-7, Lepid. Ceylon 3: 535. LITH Type-species: Diduga costata Moore, [1887] 1884-7, ibidem 3: 535, pl.211, figs 8, 9, by monotypy. [SRI LANKA] Ceylon: Dickoya.

D. costata is a junior subjective synonym of Pitane flavicostata Snellen, 1878, Tijdschr. Ent. 22: 92.

DIDUGUA Druce, 1891, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 483.

Type-species: *Didugua argentilinea* Druce, 1891, *ibidem* 1: 483, pl.40, fig.13, by monotypy. GUATEMALA: [Guatemala] City, San Geronimo.

DIDYMONYX Watson nom. n. for Diplonyx Hampson, 1900.

Type-species: *Diplonyx infumata* Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* **2**: 92, 249, fig.163, by original designation (for *Diplonyx* Hampson, 1900). [SOUTH AFRICA]: [Cape Province] Cape Colony, Annshaw.

DIPLONYX Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 92, 249.

Type-species: *Diplonyx infumata* Hampson, 1900, by original designation.

A junior homonym of *Diplonyx* Saussure, 1887, *Societas ent*. **2**: 3, – Insecta, Hymenoptera. The objective replacement name is *Didymonyx* Watson nom. n.

DIGAMA Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 297. AGAN Type-species: Digama hearseyana Moore, [1860] 1858-9,

ibidem 2: 298, pl.7A, figs 3, 3a, by monotypy. INDIA: N., [central], [Deccan] Dukhun, [S.], [Kanara] Canara, [SRI LANKA] Ceylon.

DILEMERA Agassiz, 1847. See Deilemera Hübner, [1820].

DILOBA Boisduval, [1837] 1834, Icones hist. Lépid. nouv. ou peu connus 2º 189.

Type-species: *Phalaena caeruleocephala* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 504, by monotypy. Type-LOCALITY: not stated. [EUROPE].

P. caeruleocephala has been placed in the Notodontidae, the Arctiidae, the Thyatiridae, and in the Noctuidae subfamily Plusiinae; Boursin, 1953, Bull. mens. Soc. linn. Lyon 22: 125, placed it in the Noctuidae subfamily Pantheiinae. Kiriakoff, 1970, NachrBl. bayer. Ent. 19: 101, considered that, on the basis of tympanal characters, D. caeruleocephala did not belong to the Noctuidae but should be placed in a separate family, the Dilobidae. This family-group name had already been established as Arctiidae subfamily Dilobinae by Aurivillius, 1889, Nordens Fjärilar: 79, 95, and placed on the Official List of Family-group Names in Zoology with Name No.198.

Fletcher, 1972, in Kloet & Hincks, Check List Br. Insects (Edn 2) 2: 70 retained Diloba in the Notodontidae.

The opinion of Kiriakoff, 1970, has been followed in this Catalogue.

Diloba Boisduval, 1840, was considered by the International Commission on Zoological Nomenclature, 1957, Opin. Decl. int. Commn zool. Nom. 17 (Opinion 494): 267, and placed on the Official List of Generic Names in Zoology with Name No.1233. The original reference cited above was overlooked by the Commission in Opinion 494, where Boisduval, 1840, Genera Index meth. eur. Lepid.: 88, was cited. However, both Diloba Boisduval, [1837] and Diloba Boisduval, 1840, were based on the same species.

‡HETEROMORPHA Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) (Opinion 97): 19. Also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140. Placed on the Official Index of Rejected and Invalid Generic Names in Zoology with Name No.1061.

HETEROMORPHA Hübner, 1822, Syst.-alphab. Verz.: 15, 18.

Type-species: Phalaena caeruleocephala Linnaeus, 1758, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 585.

Heteromorpha Hübner, 1822, was suppressed for the purposes of the Law of Priority but not for those of the Law of Homonymy in 1957, Opin. Decl. int. Commn zool. Nom. 17 (Opinion 494): 267. It was placed on the Official Index of Rejected and Invalid Generic Names in Zoology with Name No.1062.

DINARA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1699.

Type-species: *Dinara lineolata* Walker, 1856, *ibidem* 7: 1700, by monotypy. [INDIA]: Hindostan.

D. lineolata is a junior subjective synonym of Anticyra combusta Walker, 1855, ibidem 5: 1092.

ANTICYRA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1091.

Type-species: Anticyra combusta Walker, 1855, ibidem 5:

1092, by monotypy. PHILIPPINES.

A junior homonym of Anticyra Curtis, 1834, Lond. Edinb. Phil. Mag. (3) 4 (21): 216, — Insecta, Trichoptera. There is no objective replacement name, but A. combusta is a senior subjective synonym of Dinara lineolata Walker, 1856, the typespecies of Dinara Walker, 1856; the latter is thus available for use as a subjective replacement name.

Lectotype designated by Swinhoe, 1892, Cat. east. and Aust.

Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 303.

DINIA Walker, 1854, List Specimens lepid. Insects Colln Br.
Mus. 1: 189.

Type-species: Sphinx eagrus Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 10, pl.198, fig.C, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 338 (but cited by Hampson as ‡aeagrus, an incorrect subsequent spelling and included by Walker as Sphinx auge Linnaeus, a misidentification; see below). SURINAM.

Hampson linked eagrus in synonymy with the originally included species name Sphinx auge Linnaeus, 1767, Syst. Nat. (Edn 12): 807, but considered that auge Linnaeus sensu Walker, 1854, ibidem 1: 189 was a misidentification of Sphinx eagrus Cramer. This is a case of a misidentified type-species which should be referred to the Commission under Article 70(a) of the Code.

S. eagrus is dated from the wrapper of the part and not from

the title-page of the volume, 1782.

Dinia was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; and was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 338.

‡DINODONTA; misspelling. See Dinotodonta Holland, NOTO

DINOTODONTA Holland, 1893, Psyche, Camb. 6: 551.

Type-species: *Dinotodonta longa* Holland, 1893, *ibidem* 6: 552, fig., by original designation. [GABON]: [Ogooué] Ogové R. valley.

Dinotodonta was established in the Noctuidae; it was placed in the Notodontidae by Strand, 1912, Fauna exot. 2: 27.

‡DINODONTA; Neave, 1939, Nomencl. zool. 2: 96.

An incorrect subsequent spelling of *Dinotodonta* Holland, 1893.

DIODORINA Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 313.

Type-species: *Diodorina insueta* Kiriakoff, 1965, *ibidem* 101: 313, fig.3, by monotypy. Cameroun: Lolodorf.

DIONYCHOPUS Herrich-Schäffer, 1855. See Rhodogastria Hübner, [1819]. ARCT

DIONYCHOSCELIS Aurivillius, 1922, Ark. Zool. 14 (5): 23.

ARCT

Type-species: *Dionychoscelis venata* Aurivillius, 1922, *ibidem* 14 (5): 23, fig., by monotypy. ANGOLA.

‡DIOPEA; misspelling. See Deiopeia Curtis, 1827. ARCT

DIOPEITHES Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 58.

Type-species: *Diopeithes barnesi* Kiriakoff, 1958, *ibidem* **66**: 59, pl.3, fig.11, pl.5, fig.12, pl.9, fig.35, by original designation. [ZAIRE]: Kivu, middle Lowa valley, near Walikale.

DIOPTIS Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 9.

Type-species: *Dioptis cyma* Hübner, 1818, *ibidem* 1: 9, pl.[3], figs 17, 18, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 413. BRAZIL.

EPILAIS Boisduval, 1870, Considérations Lépid. Guatemala: 78.

Type-species: Dioptis cyma Hübner, 1818, by original designation.

A junior homonym of *Epilais* Kaup, 1829, *Skizz. Europ. Thierw.*: 145, – Aves. The objective replacement name is *Dioptis* Hübner, 1818.

‡PHELODES Boisduval, 1870, Considérations Lépid. Guatemala: 78.

Published as a junior synonym of *Epilais* Boisduval, 1870, and not subsequently used as an available name under the *Code*, Article 11(d).

DIOSPAGE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 212. CTEN

Type-species: *Sphinx rhebus* Cramer, 1779, *Uitlandsche Kapellen (Papillons exot.)* 3: 72, pl.234, fig.F, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 169. [SOUTH AMERICA]: [tropical]. The type-locality indications given in the original description, 'Indes Orientales' and 'Côte d'Afrique', are erroneous.

S. rhebus is dated from the wrapper of the part and not from

the title-page of the volume, 1782.

Diospage was established as a subgenus of Euchromia Hübner, [1819], in the 'Zygaenides'; it was placed in the Arctiidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 169; and is here transferred to the Ctenuchidae.

ENTOMIS Herrich-Schäffer, 1855, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.73, fig.424.

Type-species: Sphinx rhebus Cramer, 1779, by monotypy. A junior objective synonym of Diospage Walker, 1854.

Entomis is dated from the wrapper of Hefte 20-21, and not from the title-page of the volume, 1850-1858.

DIOTA Wallengren, 1865, K. svenska VetenskAkad. Handl. 5 (4): 46.

Type-species: Euchelia rostrata Wallengren, 1860, Wien. ent. Monatschr. 4: 161, by monotypy. [South Africa]: Caffraria.

‡DICTA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 426.

An incorrect subsequent spelling of Diota Wallengren, 1865.

‡DIPAENA; misspelling. See Dipaenae Walker, 1854.

DIPAENAE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 257.

Type-species: Euchromia ferruginosa Walker, 1854, ibidem

1: 259, by PRESENT DESIGNATION. Type-locality: not

known, [South America]: [tropical].

The type-species designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 173, of Anycles rhodura Butler, 1876, J. Linn. Soc. (Zool.) 12: 425 is invalid as he linked this name not with an originally included name, but with an unnamed Walker 'var.' (in the Lithosiinae) of an originally included name Euchromia acharon Fabricius, [1777], Genera Insect.: 556 (Zygaenidae). The present designation of ferruginosa Walker maintains the long accepted concept of Dipaenae because Walker's 'var.' (subsequently named by Butler, 1876, ibidem 12: 425, as *rhodura*) is currently considered to be conspecific with ferruginosa Walker.

Dipaenae was established as a subgenus of Euchromia Hüb-

ner, [1819].

‡DIPAENA; Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 105.

An incorrect subsequent spelling of Dipaenae Walker, 1854.

DIPHTERAMOMA Berio, 1961, Memorie Soc. ent. ital. 40: 114. NOCTUIDAE

Type-species: Noctua orion Esper, 1787, Die Schmett. 4: pl.118, figs 4-7; 1791 ibidem 4: 283 by original designation.

Syntypes [NETHERLANDS]: Amsterdam.

N. orion is dated from the wrapper of the Heft and not from the title-page, 1786, of the volume. N. orion is a junior subjective synonym of Phalaena alpium Osbeck, 1778, the typespecies of Moma Hübner, [1820].

Diphteramoma in the Noctuidae, Acronictinae is cited here as it was not included in Nye, 1975, Generic Names Moths

World 1.

DIPHTHERASPIS Meyrick, 1892, in Lucas, Proc. R. Soc. Od 8: 74.

Type-species: Diphtheraspis modicus Lucas, 1892, ibidem 8: 74, by monotypy. Australia: Brisbane.

DIPLONYCHODES Kiriakoff, 1954, Revue Zool. Bot. afr.

Type-species: Diplonychodes ruficeps Kiriakoff, 1954, ibidem 50: 173, pl.1, fig.3, pl.2, fig.5, by original designation. [ZAIRE]: [Lubumbasha] Elisabethville.

DIPLONYX Hampson, 1900. See Didymonyx Watson nom.

DIPTEROMORPHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.28.

Type-species: Dipteromorpha adusta Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.28, by monotypy. Mexico: Cuernavaca, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls. 75-107: 4.

DIPTILON Prittwitz, 1870, Stettin. ent. Ztg 31: 349. Type-species: Diptilon telamonophorum Prittwitz, 1870, ibidem 31: 350, pl.2, fig.2, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 148. [Brazil]: Rio [de Janeiro].

DIPTILUM Zerny, 1912, in Wagner, Lepid. Cat. 7: 106. An unjustified emendation of Diptilon Prittwitz, 1870.

DIPTILUM Zerny, 1912. See Diptilon Prittwitz, 1870.

DISAULOTA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus.

Type-species: Areva leptalina Druce, 1885, Biologia cent.am. (Zool.) Lepid.-Heterocera 1: 133, pl.13, fig.5, by original designation. Costa Rica: Cache.

DISCONEURA Bryk, 1953, Ark. Zool. (2) 5: 208. ARCT Type-species: Disconeura tristriata Bryk, 1953, ibidem (2) 5: 208, by original designation. PERU: eastern.

DISCOPHLEBIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2.): pl.96, fig.8.

Type-species: Discophlebia catocalina Felder, 1874, ibidem 2 (Abt.2): pl.107, fig.8, by monotypy. AUSTRALIA: [South Australia], Adelaide, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

Discophlebia, when established, was not placed in a family; it was placed in the Zeuzeridae, now Cossidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 878; it was placed in the Notodontidae by Turner, 1903, Proc. Linn. Soc. N.S.W. and was transferred to the Thaumetopoeidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

DISOIDEMATA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 87, 188.

Type-species: Disoidemata osmophora Hampson, 1900, ibidem 2: 188, fig. 106, by original designation. Mexico: Vera Cruz

DISPARCTIA Toulgoët, 1978, Nouv. Revue Ent. 8: 228.

Type-species: Spilarctia vittata Druce, 1898, Ann. Mag. nat. Hist. (7) 1:211, by original designation. [CAMEROUN]: Mongoma-Lobah.

DISPARIA Nagano, 1916, Bull. Nawa ent. Lab. 1: 3. NOTO Type-species: Fentonia sordida Wileman, 1911, Trans. ent. Soc. Lond. 1911: 286, pl.30, fig.8, by original designation. JAPAN: Yoshino, Yamato.

F. sordida is a junior subjective synonym of Fentonia variegata Wileman, 1910, Entomologist 43: 290.

See also Epifentonia Kiriakoff, 1960.

DISPAROIDES Nakamura, 1976, Ent. Rev. Japan 29: 49, 50.

Type-species: Disparia seriatopunctata Matsumura, 1931, 6000 Illust. Insects Japan-Empire: 627, fig., by original designation. TAIWAN.

Disparoides was established as a subgenus of Pseudofentonia Strand, 1912.

DISPHRAGIS Hübner, [1820] 1816, Verz. bekannter Schmett.: 194.

Type-species: Phalaena tharis Stoll, 1780, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 38, pl.308, fig.D, by subsequent designation by Grote, 1902, Allg. Z. Ent. 7: 397. SURINAM.

P. tharis is dated from the wrapper of part 26, not from the title-page of volume 4 which is dated 1782.

DISRACHA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 46.

Type-species: Chadisra persimilis Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 463, by original designation. [South AFRICA]: Natal, Durban.

DITAXIS Boisduval, 1870. See Hyalurga Hübner, [1819].

DIXANAENE Dyar, 1914, Proc. U.S. natn. Mus. 47: 172.

Type-species: Dixanaene lepidocaena Dyar, 1914, ibidem 47: 172, by original designation. [TOBAGO] Tobaga I., PANAMA: La Chorrera and [Portobelo] Porto Bello.

DIXOPHLEBIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 397.

Type-species: Pseudomya quadristrigata Walker, [1865]

1864, List Specimens lepid. Insects Colln Br. Mus. 31: 82, by original designation. [BRAZIL]: [Tefé] Ega.

DOA Neumoegen & Dyar, 1894, Jl N.Y. ent. Soc. 2: 171.

Type-species: *Doa dora* Neumoegen & Dyar, 1894, *ibidem* 2: 171, by original designation. MEXICO: Guadalajara.

Doa was established in the Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Dyar, [1903] 1902, Bull. U.S. natn. Mus. 52: 261; it was transferred to the Pericopidae, now Arctiidae Pericopinae, by Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 563; it was transferred to the Lymantriidae Pterothysaninae, now Pterothysanidae, by Bryk, 1934, in Strand, Lepid. Cat. 62: 356; but was retained in the Pericopidae, now Arctiidae Pericopinae, by McDunnough, 1938, Mem. sth. Calif. Acad. Sci. 1: 132.

DOCYNA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1084.

Type-species: *Docyna triplaga* Walker, 1855, *ibidem* 5: 1084, by monotypy. [BRAZIL]: Rio de Janeiro.

DODIA Dyar, 1901, Jl N.Y. ent. Soc. 9: 85.

Type-species: *Dodia albertae* Dyar, 1901, *ibidem* 9: 85, by original designation. Canada: Alberta, Calgary.

DOGNINA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 261 (key), 301.

Type-species: Lirimiris veltini Dognin, 1890, Naturaliste 12: 128, by original designation. [ECUADOR]: Loja.

DOHERTYA Hampson, 1894, Fauna Br. India (Moths) 2: vi, 40, 63.

Type-species: *Dohertya cymatophoroides* Hampson, 1894, *ibidem* 2: 63, fig.31, by original designation. [BURMA]: E. Pegu.

DOKUGA Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19: 37.

Type-species: *Dokuga purpurescens* Matsumura, 1927, *ibidem* 19: 37, by original designation (but cited as ‡*purpurascens*, an incorrect (of a multiple) original spelling). [Taiwan] Formosa: Horisha.

D. purpurescens is a junior subjective synonym of Pida postalba Wileman, 1910, Entomologist 43: 287.

DOLGOMA Moore, 1878, Proc. zool. Soc. Lond. 1878: 20.

Type-species: *Lithosia reticulata* Moore, 1866, *ibidem* **1865**: 798, by original designation. INDIA: Darjeeling.

DOLICHE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 529.

Type-species: *Doliche gelida* Walker, 1854, *ibidem* 2: 529, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 300. [BANGLADESH]: [Sylhet] Silhet.

DOLICHESIA Schaus, 1911, Ann. Mag. nat. Hist. (8) 7: 362.

Type-species: *Dolichesia falsimonia* Schaus, 1911, *ibidem* (8) 7: 362, by monotypy. Costa Rica: Sixola.

DOLOPHROSYNE Prout, 1918, *Novit. zool.* **25**: 396 (key), 403.

Type-species: *Dolophrosyne mirax* Prout, 1918, *ibidem* 25: 403, by original designation. PERU: eastern, Huancabamba, near Cerro de Pasco.

DONDERA Moore, 1877, Ann. Mag. nat. Hist. (4) 20: 344.

Type-species: *Dondera alba* Moore, 1877, *ibidem* (4) **20**: 344, by monotypy. [SRI LANKA] Ceylon.

DORACIS Boisduval, 1870, Considérations Lépid. Guatemala: 96.

Type-species: *Doracis coracina* Boisduval, 1870, *ibidem*: 96, by monotypy. HONDURAS, MEXICO.

D. coracina is a junior subjective synonym of Ardonea morio Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 482.

See also Ardonea Walker, 1854.

DORIMENA Boisduval, 1870. See Daritis Walker, 1855.
PERI

‡DORIMENE; misspelling. See Daritis Walker, 1855.

DORSTIANA Laporte, 1974, Bull. mens. Soc. linn. Lyon 43: 248. NOCTUIDAE

Type-species: *Dorstiana aethiopica* Laporte, 1974, *ibidem* 43: 248, figs 8, 40, by monotypy. Holotype ô, ETHIOPIA: marais de Dinsho, (MNHN, Paris).

Dorstiana in the Noctuidae, Hadeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

DOTHA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 259. LITH Type-species: Dotha ctenuchoides Walker, [1865] 1864,

ibidem 31: 259, by monotypy. SUMATRA.

DOTTIA Schaus, 1911, Ann. Mag. nat. Hist. (8) 7: 621.

Type-species: *Dottia effecta* Schaus, 1911, *ibidem* (8) 7: 621, by monotypy. Costa Rica: Banana River.

DRAGONERA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 151 (key), 209.

Type-species: *Bireta nana* Swinhoe, 1889, *Proc. zool. Soc. Lond.* **1889**: 407, by original designation. INDIA: North Kanara.

Dragonera was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 87.

DRAPETIDES Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 364.

Type-species: *Lophopteryx uniformis* Swinhoe, 1907, *Ann. Mag. nat. Hist.* (7) **19**: 208, by original designation. [RHODESIA]: Mashonaland.

DRASTOMA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 258 (key), 264.

Type-species: *Tifama dardania* Druce, 1895, *Ann. Mag. nat. Hist.* (6) **15**: 49, by original designation. Costa Rica.

DRUCEA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 130.

Type-species: Mastigocera oedippus Boisduval, 1870, Considérations Lépid. Guatemala: 81, by original designation. GUATEMALA.

Drucea was cited incorrectly, by Neave, 1939, Nomencl. zool. 2: 160, as Kirby's replacement name for Mastigocera Harris, 1839, Am. J. Sci. 36: 315, — Lepid., Ctenuchidae, but Kirby, 1892, proposed Drucea for the then unnamed concept Mastigocera Harris sensu Boisduval, 1870, Considerations Lépid. Guatemala: 81, and was not replacing Mastigocera Harris, 1839.

DRUGERA Schaus, 1901, *Trans. ent. Soc. Lond.* 1901: 260 (key), 293.

Type-species: *Edema morona* Druce, 1898, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 2: 455, pl.90, fig.14, by original

designation. Costa Rica: Candelaria Mts, Panama: Volcan de Chiriqui, Obispo.

DRYMONIA Hübner, [1819] 1816, Verz. bekannter Schmett.: NOTO

Type-species: *Bombyx dodonaea* [Denis & Schiffermüller], 1775, *Ankündung syst. Werkes Schmett. Wienergegend*: 49, by subsequent designation by Curtis, 1839, *Br. Ent.* 16: 755. [AUSTRIA]: Vienna district.

DRYNOBIA Duponchel, [1845] 1844, *Cat. méth. Lépid. Eur.* (2): 92.

Type-species: *Phalaena velitaris* Hufnagel, 1766, *Berlin. Mag.* **3** (4): 394 (but included by Duponchel as 'velitaris Esper', an incorrect authorship), by PRESENT DESIGNATION. GERMANY: Berlin.

See also Ochrostigma Hübner, [1819].

DUBIANACLIA Griveaud, 1964, Faune Madagascar 17: 107.

Type-species: Syntomis butleri Mabille, 1882, Naturaliste 2: 134, by original designation. MADAGASCAR.

DUDUSA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 446.

Type-species: *Dudusa nobilis* Walker, 1865, *ibidem* **32**: 447, by monotypy. China: northern.

DUDUSOIDES Matsumura, 1929, Insecta matsum. **4**: 79 (key), 80.

Type-species: *Dudusa sphingiformis* Moore, 1872, *Proc. zool. Soc. Lond.* **1872**: 577, pl.34, fig.1, by original designation. [INDIA]: Sikkim.

DUDUSOPSIS Matsumura, 1929, Insecta matsum. 4: 79 (key), 81.

Type-species: *Dudusa fumosa* Matsumura, 1925, *Zool. Mag. Tokyo* 37: 391, fig., by original designation. TAIWAN.

DUFAYELLA Capuse, 1972, Ent. Z., Frankf.a.M. 82: 89.

Type-species: Sarrothripus asiatica Krulikovski, 1904, Revue russe Ent. 4: 91, by original designation. Holotype ô, [U.S.S.R.]: [Turkmenistan], [Ashkhabad] Aschabad.

S. asiatica was established as 'Sarrothripus revayana Sc., var. asiatica nova.'

Dufayella was established as a subgenus of Nycteola Hübner, 1822, Syst.-alphab. Verz.: 60, 66.

Dufayella in the Noctuidae, Sarrothripinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

DUFRANEELLA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 14. THYRET

Type-species: *Metarctia grisescens* Dufrane, 1945, *Bull. Annls Soc. ent. Belg.* 81: 131, by original designation. [ZAIRE]: Kivu, Kamituga.

Dufraneella was established as a subgenus of Apisa Walker, 1855.

DUGONIA Schaus, 1928, Proc. U.S. natn. Mus. 73 (19): 43.

Type-species: *Dugonia eliera* Schaus, 1928, *ibidem* **73** (19): 43, by original designation. French Guiana.

DUKINFIELDIA Schaus, 1894, Proc. zool. Soc. Lond. 1894: NOTO

Type-species: *Dukinfieldia suprema* Schaus, 1894, *ibidem* 1894: 234, by monotypy. [BRAZIL]: Parana, Castro.

Dukinfieldia was established in the Eupterotidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 320.

DULICHIA Walker, 1855. See Knappetra Nye nom. n.

DUNAMA Schaus, 1912, Ann. Mag. nat. Hist. (8) 9: 52.
NOTO

Type-species: *Dunama angulinea* Schaus, 1912, *ibidem* (8) 9: 52, by original designation. Costa Rica: Guapiles.

DURA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 56. LYMA

Type-species: *Dura alba* Moore, 1879, *in* Hewitson & Moore, *ibidem* (1): 56, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 482. INDIA: [W. Bengal], [Darjeeling] Darjiling.

DUSPONERA Dognin, 1914, Hétérocères nouv. Am. Sud 8: NOCTUIDAE

Type-species: *Dusponera semifalcata* Dognin, 1914, *ibidem* 8: 48, by monotypy. Holotype 9, [ECUADOR]: environs de Loja, Equateur.

Dusponera Dognin, 1914, in the Noctuidae, Hypeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

Dusponera probably originated from Schaus but was used and made nomenclaturally available by Dognin in 1914, prior to its proposal and generic description by Schaus in 1916, for a similar concept but having a different type-species, Dusponera fannia Schaus, 1916. Dusponera Schaus, 1916, (as cited in Generic Names Moths World 1: 167) is thus a junior homonym. There is no objective replacement name but D. fannia is congeneric with D. semifalcata, the type-species of Dusponera Dognin, 1914. The latter is therefore available for use as a subjective replacement name.

DYASIA Schaus, 1905, Proc. U.S. natn. Mus. 29: 231. NOTO Type-species: Dyasia viviana Schaus, 1905, ibidem 29: 232, by monotypy. French Guiana: Maroni River.

DYASMA Collenette, 1961, Lambillionea 60: 103. LYMA Type-species: Dasychira thaumatopoeides Schultze, 1934, Dt. ent. Z. Iris 48: 116, fig. 10, by original designation. [CENTRAL AFRICAN EMPIRE]: Ubangi, Mongoumba.

DYCLADIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.20. CTEN Type-species: Dycladia correbioides Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.20, by monotypy. [COLOMBIA]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

DYLOMIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.13. NOTO Type-species: Dylomia tortricina Felder, 1874, ibidem 2 (Abt.2): pl.97, fig.13, by subsequent designation by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 327. [BRAZIL]: River Amazon, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to

pls 75-107: 4.

DYPHLEBIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.32.

Type-species: *Dyphlebia trimenii* Felder, 1874, *ibidem* 2 (Abt.2): pl.106, fig.32, by subsequent designation by Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* 2: 104, but cited as *Lithosia pandula* Boisduval, 1847, a nominal species not originally included in *Dyphlebia*. Hampson, however, placed *pandula* as the senior synonym of *trimenii*, a nominal species originally included in *Dyphlebia*. Under the *Code*, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. [SOUTH AFRICA]: Caffraria, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 4.

D. trimenii is a junior subjective synonym of Lithosia pandula Boisduval, 1847, in Delegorgue, Voyage Afr. australe 2: 597. See also Lepista Wallengren, 1863.

DYPNA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 151, 166; 1962 October 20, Bonn. zool. Beitr. 13: 222.

Type-species: *Bireta triangularis* Kiriakoff, 1962 [July 31], *ibidem* **98** (10): 166; 1962 October 20, *ibidem* **13**: 222, abb.1, fig.4, pl.1, fig.3 (as *Dypna triangularis* sp. nov.), by original designation. China: [Chekiang, west Tien-mu-Shan]. Additional data cited by Kiriakoff, 1962 October 20, *ibidem* **13**: 222.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr. to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper B. triangularis was described as a new species in the combination Dypna triangularis and Dypna was proposed as a new genus. As the dates show, the Belgian paper appeared first.

In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

Dypna was established as a subgenus of Bireta Walker, 1856; it was treated as a genus by Kiriakoff, 1962 October 20, ibidem 13: 222 and again in 1967, in Wytsman, Genera Insect. 217 (B): 39.

DYSAUXES Hübner, [1819] 1816, Verz. bekannter Schmett.: 171.

Type-species: *Phalaena ancilla* Linnaeus, 1767, *Syst. Nat.* (Edn 12) 1 (2): 835, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 107. GERMANY.

NACLIA Boisduval, 1840, Genera Index eur. Lepid.: 60.

Type-species: *Phalaena ancilla* Linnaeus, 1767, by subsequent designation by Blanchard, 1840, *in* Castelnau, *Hist. nat. Insectes* 3: 490.

A junior objective synonym of Dysauxes Hübner, [1819].

DYSSCHEMA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 31.

Type-species: *Dysschema hypoxantha* Hübner, 1818, *ibidem* 1: 31, pl.[34], figs 191, 192, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 183. SURINAM.

TAXILA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 764.

Type-species: Chelonia crucifera Perty, [1833], in Martius, Delectus Anim. articulatorum Bras.: 160, fig., by monotypy. BRAZIL: Minarum.

C. crucifera is a junior subjective synonym of Eucharia sacrifica Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 21, pl. [82], figs 473, 474.

A junior homonym of *Taxila* Doubleday, 1847, *Synonymic List Br. Lepid*. 2: 2, – Lepid., Riodinidae. There is no objective replacement name but the type-species of *Taxila* Walker is currently considered to be congeneric with the type-species of *Dysschema* Hübner, 1818. The latter is therefore available for use as a subjective replacement name.

EALA Collenette, 1937, Revue Zool. Bot. afr. 29: 422. LYMA Type-species: Eala hemiluta Collenette, 1937, ibidem 29: 422, pl.6, fig.4, by original designation. [ZAIRE] Belgian Congo: [Equateur], Eala.

ECDEMUS Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.73, fig.420.

Type-species: Ecdemus hypoleucus Herrich-Schäffer, [1855]

1850-1858, *ibidem* 1(1): wrapper, pl.73, fig.420, by monotypy. [Brazil]: [Para], Caméta.

ECHEMIS Boisduval, 1870, Considérations Lépid. Guatemala: 99.

Type-species: *Phaegoptera elegantissima* Herrich-Schäffer, [1856] 1850-1858, *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.92, fig.528 (but cited as ‡*Phegoptera*, an incorrect subsequent spelling), by PRESENT DESIGNATION. MEXICO.

P. elegantissima is a junior subjective synonym of Anaxita decorata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 748.

See also Anaxita Walker, 1855.

ECHETA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 96, 97; [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 16, 17. ARCT

Type-species: Creatonotos divisus Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.52, fig.282, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 197. Type-locality: not stated, [South America]: [tropical].

Echeta was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

ECHLIDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 386.

Type-species: *Echlida subjecta* Walker, 1865, *ibidem* 32: 387, by monotypy. [INDIA]: South Hindostan.

E. subjecta is a junior subjective synonym of Thyrgorina spilosomata Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 318.

See also Thyrgorina Walker, [1865].

ECHONEURA Butler, 1876, J. Linn. Soc. (Zool.) 12: 383.

Type-species: *Euchromia intricata* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 287, by original designation. BRAZIL: Rio [de] Janeiro.

ECNOMODES Turner, 1903, Proc. Linn. Soc. N.S.W. 28: 46 (key), 69.

Type-species: *Chlenias sagittaria* Lucas, 1900, *Proc. R. Soc. Qd* **15**: 148, by original designation. AUSTRALIA: Queensland, Brisbane.

ECPANTHERIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 183.

Type-species: *Bombyx ocularia* Fabricius, 1775, *Syst. Ent.*: 564, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 216. AMERICA: [tropical] 'meridionali'.

ECREGMA Walker, [1858] 1857, List Specimens lepid. Insects Colln Br. Mus. 12: 941 (key), 966.

Type-species: *Ecregma tranducta* Walker, [1858] 1857, *ibidem* 12: 967, by monotypy. VENEZUELA.

Ecregma was established in the Noctuidae; it was placed

in the Notodontidae by Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 333.

ECTEINA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 196, 215.

Type-species: *Phalaena mesomella* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 535, by monotypy. TYPE-LOCALITY: not stated, [EUROPE].

See also Cybosia Hübner, [1819].

‡ECTHEMA; misspelling. See Esthema Hübner, [1820].

‡ECTYPA; misspelling. See Ectypia Clemens, 1861. ARCT

ECTYPIA Clemens, 1861, Proc. Acad. nat. Sci. Philad. 1860: 529.

Type-species: *Ectypia bivittata* Clemens, 1861, *ibidem* **1860**: 530, by monotypy. [U.S.A.]: Texas.

‡ECTYPA; Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 316.

An incorrect subsequent spelling of Ectypia Clemens, 1861.

EDEMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1028.

Type-species: *Phalaena albifrons* Smith, 1797, in Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 159, pl.80, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 573. [U.S.A.]: Georgia.

See also Symmerista Hübner, [1821].

EGONOCIA Marumo, 1920, J. Coll. Agric. imp. Univ. Tokyo 6: 281 (key), 333.

Type-species: Somera cyanea Leech, 1889, Proc. zool. Soc. Lond. 1888: 642, pl.32, fig.5, by original designation. Type-LOCALITY: not stated, [JAPAN]: [Gifu]; data taken from a male syntype in BMNH.

EGONOCIADES Kiriakoff, 1964, Bonn. zool. Beitr. 14: 269.

Type-species: *Egonociades basistriata* Kiriakoff, 1964, *ibideni* 14: 269, pl.4, fig.3, by original designation. China: Chekiang, west Tien-mu-Shan.

EGURIA Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 37.

Type-species: *Ptilodontis ornata* Oberthür, 1884, *Etudes d'Entomologie* 10: 15, pl.2, fig.5, by original designation. [U.S.S.R.]: Sidemi.

EILEMA Hübner, [1819] 1816, Verz. bekannter Schmett.: 165.

Type-species: *Bombyx caniola* Hübner, [1808] 1796, *Samml. eur. Schmett.* 3: pl.51, fig.220, by subsequent designation by Moore, 1878, *Proc. zool. Soc. Lond.* 1878: 17. EUROPE.

SYSTROPHA Hübner, [1819] 1816, Verz. bekannter Schmett.: 166.

Type-species: *Bombyx aureola* Hübner, [1803] 1796, *Samml. eur. Schmett.* 3: pl.24, fig.98, by subsequent designation by Moore, 1878, *Proc. zool. Soc. Lond.* 1878: 18. EUROPE.

A junior homonym of Systropha Illiger, 1806, Magazin Insektenk. (Illiger) 5: 145, – Insecta, Hymenoptera. There is no objective replacement name but the type-species of Systropha Hübner is currently considered to be congeneric with the type-species of Eilema Hübner, [1819]. The latter is therefore available for use as a subjective replacement name.

HILEMA Agassiz, 1847, Nomencl. zool. (Index univl.): 135, 182.

An unjustified emendation of Eilema Hübner, [1819].

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Hilema* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

ILEMA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 83, 130.

An unjustified emendation of Eilema Hübner, [1819].

A junior homonym of *Ilema* Moore, [1860] 1858-9, *Cat. lepid. Insects Mus. nat. Hist. East-India House* 2: 341, – Lepid., Lymantriidae. The objective replacement name is *Eilema* Hübner, [1819].

ELA Walker, 1862. See Urocoma Herrich-Schäffer, [1858].

ELAPHRODES Bethune-Baker, 1909, Ann. Mag. nat. Hist. (8) 3: 423.

Type-species: *Elaphrodes nephocrossa* Bethune-Baker, 1909, *ibidem* (8) 3: 423, by original designation. [ZAIRE]: Ituri Forest, between Makala and Beni.

ELASMIA Möschler, 1886, Abh. senckenb. naturforsch. Ges. 14 (3): 36.

Type-species: *Elasmia lignosa* Möschler, 1886, *ibidem* **14** (3): 36, pl.-fig.30, by monotypy. Jamaica.

Elasmia Möschler, 1886, is not preoccupied by ‡Elasmia Hübner, 1822, an incorrect (of a multiple) original spelling of Elasmion Hübner, 1822, Syst.-alphab. Verz.: 62, 66-79, – Lepid., Adelidae.

ELEALE Walker, 1862. See Tamseale Nye, 1975.

ELIGMODONTA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 181.

Type-species: *Phalaena ziczac* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 504, by original designation. Type-Locality: not stated, [EUROPE].

ELKNERIA Börner, 1932, in Brohmer, Fauna Dtl. (Edn 4): 393.

Type-species: *Phalaena pudibunda* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 503, by monotypy. Type-locality: not stated, [EUROPE].

‡DASYCHIRA Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) (Opinion 97): 19. Also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140.

Only included species: Phalaena pudibunda Linnaeus, 1758.

ELLIDA Grote, 1876, Can. Ent. 8: 125. NOTO Type-species: Ellida gelida Grote, 1876, ibidem 8: 125, by

monotypy. CANADA.

E. gelida is a junior subjective synonym of Cymatophora caniplaga Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 9: 18.

ELORIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 813.

Type-species: Eloria intacta Walker, 1855, ibidem 4: 814, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 430, but cited as 'E. Spectra, Hübn.', a nominal species not originally included in Eloria. Kirby, however, placed spectra as the senior synonym of intacta, a nominal species originally included in Eloria. Under the Code, Article 69(a)(iv), this constitutes a type-species designation. [Brazil]: Parà.

ELPIS Dyar, 1893. See Diacrisia Hübner, [1819]. ARCT

ELSA Kiriakoff, 1953. See Elsita Kiriakoff, 1954. THYRET

ELSITA Kiriakoff, 1954, Bull. Annls Soc. r. ent. Belg. 90: 29.

THYRET

Type-species: Anace invaria Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1720, by original designation (for Elsa Kiriakoff, 1953). WEST AFRICA.

Elsita was established as the objective replacement name for Elsa Kiriakoff, 1953.

ELSA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 23.

Type-species: Anace invaria Walker, 1856, by original designation.

A junior homonym of *Elsa* Honrath, 1892, *Berl. ent. Z.* 36: 436, – Lepid., Nymphalidae. The objective replacement name is *Elsita* Kiriakoff, 1954.

ELYMIOTIS Walker, 1857, List Specimens lepid. Insects Colln Br. Mus. 11: 609.

Type-species: *Elymiotis notodontoides* Walker, 1857, *ibidem* 11: 609, by monotypy. Type-locality: not known, [Brazil].

From a multiple original spelling in which ‡Elymistis was published in the key on page 494 and Elymiotis was published in the generic and type-species headings on page 609, we have adopted the spelling in current use, namely Elymiotis, as the correct spelling.

Elymiotis was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 272.

See also Cicynna Walker, [1858].

‡ELYMISTIS; Walker, 1857, List Specimens lepid. Insects Colln Br. Mus. 11: 494.

An incorrect (of a multiple) original spelling of *Elymiotis* Walker, 1857.

‡ELYMISTIS; misspelling. See Elymiotis Walker, 1857.

ELYSIUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 713. ARCT

Type-species: *Elysius conspersus* Walker, 1855, *ibidem* 3: 714, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 219. [Brazil]: Para.

EMINARIA Bethune-Baker, 1908, Ann. Mag. nat. Hist. (8) 2: 262.

Type-species: *Eminaria nigropunctata* Bethune-Baker, 1908, *ibidem* (8) 2: 262, by original designation. [UGANDA]: Açholi [District], [Patiko] Patigo.

E. nigropunctata is a junior subjective synonym of Spilosoma mhondana Bartel, 1903, Dt. ent. Z. Iris 16: 179.

EMPODISMA Kiriakoff, 1964. See Boscawenia Fletcher nom. n. NOTO

EMPUSA Hübner, [1819]. See Idalus Walker, 1855. ARCT

EMPYREUMA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 12.

Type-species: Sphinx pugione Linnaeus, 1767, Syst. Nat. (Edn 12) 1 (2): 807, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 422. [WEST INDIES]: [Virgin Islands, St Thomas Island] insula S. Thomae.

‡CHRYSAOR Hübner, 1808, Erste Zuträge Samml. exot. Schmett.: 4.

Included in a work rejected for nomenclatural purposes by Int. Commn zool. Nom., 1966, Bull. zool. Nom. 23 (Opinion 789): 213. Placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No. 1833.

Generic Names in Zoology: Name No. 1833.
Only included species ‡Chrysaor erythropterus Hübner, 1808, ibidem: 4, a concept previously named as Sphinx pugione Linnaeus, 1767, Syst. Nat. (Edn 12) 1: 807.

EMURENA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 52.

Type-species: *Emurena fernandezi* Watson, 1975, *ibidem* 25: 54, fig. Guyana: Potaro.

EMYDIA Boisduval, 1828. See Spiris Hübner, [1819].

ENDERA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 228.

Type-species: *Euchromia vulcanus* Walker, 1854, *ibidem* 1: 228, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 124. MEXICO.

Endera was established as a subgenus of Euchromia Hübner, [1819].

ENDROSA Hübner, [1819] 1816, Verz. bekannter Schmett.: 167.

Type-species: *Noctua roscida* [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 68, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 415. [Austria]: Vienna district.

‡ENERYTHRA; misspelling. See Euerythra Harvey, 1876.

ENIOCHE Druce, 1896, Ann. Mag. nat. Hist. (6) 18: 34.

Type-species: *Enioche pitthea* Druce, 1896, *ibidem* (6) **18**: 34, by original designation. PERU.

E. pitthea is a junior subjective synonym of Euchromia basalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 251.

See also *Pseudosphenoptera* Butler, 1876, and *Chloropsinus* Butler, 1876.

ENOME Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 883.

Type-species: *Enome ampla* Walker, 1855, *ibidem* 4: 883, by monotypy. [Sri Lanka] Ceylon, India: North.

ENOMOTARCHA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 54.

Type-species: *Somera chloana* Holland, 1893, *Psyche, Camb.* 6: 476, pl.17, fig.19, by original designation. [GABON]: [Ogooué] Ogové R. valley.

ENOPE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 208.

Type-species: *Euchromia hirsuta* Walker, 1854, *ibidem* 1: 208, by monotypy. Type-locality: not known, [AMERICA]: [tropical].

Enope was established as a subgenus of Euchromia Hübner, [1819].

E. hirsuta is a junior subjective synonym of Trichela tolumnensis Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.13, fig.53.

See also Trichela Herrich-Schäffer, [1853].

ENTOMIS Herrich-Schäffer, 1855. See Diospage Walker, 1854.

EOHEMERA Aurivillius, 1904, Ark. Zool. 2 (4): 42. ARCT Type-species: Nyctemera fulleri Druce, 1883, Entomologist's mon. Mag. 20: 157, by original designation. [CAMEROUN] Cameroon.

EOPIRGA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 156.

Type-species: Eopirga candida Hering, 1926, ibidem 14: 156,

pl.26, row h, by original designation. MADAGASCAR: Antanambé.

The date of publication of the part of Seitz containing page 156 is cited on page 153 as 5.xi.1926 in both the German and the English editions.

EPANAPHE Aurivillius, 1904, Ark. Zool. 2 (4): 8. THAU Type-species: Anaphe moloneyi Druce, 1888, Proc. zool. Soc. Lond. 1887: 673, pl.55, fig.5, by subsequent designation by Janse, 1920, Ann. Transv. Mus. 7: 234. GAMBIA.

Epanaphe was established in the Notodontidae; it was placed in the Thaumetopoeidae by Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 398.

EPANYCLES Butler, 1876, J. Linn. Soc. (Zool.) 12: 425.

Type-species: *Euchromia imperialis* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 241, by original designation. Brazil: Santarem.

EPATOLMIS Butler, 1877, Trans. ent. Soc. Lond. 1877: 348.

Type-species: Atolmis japonica Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 223, by original designation. JAPAN.

A. japonica is a junior subjective synonym of Phalaena caesarea Goeze, 1781, Ent. Beytrage 3 (3): 63.

‡EPATOSIS; misspelling. See Epitoxis Wallengren, 1863.

EPECTAPTERA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xxi, 31, 531.

Type-species: Aclytia laudabilis Druce, 1896, Ann. Mag. nat. Hist. (6) 18: 35, by original designation. BOLIVIA.

EPECTOPTERA Zerny, 1912, in Wagner, Lepid. Cat. 7: 151. An unjustified emendation of Epectaptera Hampson, 1898.

EPECTOPTERA Zerny, 1912. See Epectaptera Hampson, 1898.

EPEIROMULONA Field, 1952, Proc. U.S. natn. Mus. 102: 455.

Type-species: *Epeiromulona lephina* Field, 1952, *ibidem* **102**: 459, fig., by original designation. Panama: Colon Province, [Portobelo] Porto Bello.

EPEUPROCTIS Matsumura, 1933, Insecta matsum. 7: 122.

Type-species: *Euproctis tamahonis* Matsumura, 1927, *J. Coll. Agric. Hokkaido imp. Univ.* 19: 43, by original designation. [Talwan] Formosa: near Niitaka, Tamaho.

‡EPHALTIAS; misspelling. See Ephialtias Hübner, [1819].

EPHESTRIS Hübner, [1820] 1816, Verz. bekannter Schmett.: 178.

Type-species: *Hipocrita melaxantha* Hübner, [1809] 1806, *Samml. exot. Schmett.* 1: pl.[188] (but cited as ‡*Hypocrita*, an incorrect subsequent spelling), by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 187. Type-LOCALITY: not stated, [Brazil].

LAMA Walker, 1869, Characters undescr. Lepid. Heterocera: 7.

Type-species: *Hipocrita melaxantha* Hübner, [1809], by original designation.

A junior homonym of *Lama* Cuvier, 1800, *Lecons Anat. Comparée* 1: table [1], – Mammalia. The objective replacement name is *Ephestris* Hübner, [1820].

EPHIALTIAS Hübner, [1819] 1816, Verz. bekannter Schmett.: 176.

Type-species: *Hypocrita abrupta* Hübner, [1809] 1806, *Samml. exot. Schmett.* 1: pl.[184], by subsequent designation by Grote, 1866, *Proc. ent. Soc. Philad.* 6: 175. Type-locality: not stated, [Brazil]: [north-east].

‡EPHALTIAS; Hübner, 1823, Zuträge Samml. exot. Schmett. 2: 15.

An incorrect subsequent spelling of *Ephialtias* Hübner, [1819].

RETILA Boisduval, 1870, Considérations Lépid. Guatemala: 94

Type-species: *Hypocrita abrupta* Hübner, [1809], by subsequent designation by Prout, 1918, *Novit. zool.* **25**: 420.

A junior objective synonym of Ephialtias Hübner, [1819].

EPIBALACRA Kiriakoff, 1957, Bull. Annls Soc. r. ent. Belg. 93: 149.

Type-species: *Metarctia preussi* Aurivillius, 1904, *Ark. Zool.* **2** (4): 31, by original designation. [Cameroun]: [Mt Cameroun] Kamerunberg, Buea.

Epibalacra was established as a subgenus of Balacra Walker, 1856.

EPICALLIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 182.

Type-species: *Phalaena villica* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 501, by monotypy. Type-locality: not stated, [Europe].

EPICERURA Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 557.

Type-species: *Epicerura tanda* Bethune-Baker, 1911, *ibidem* (8) 7: 557, by original designation. ANGOLA: N'Dalla Tando. *E. tanda* is a junior subjective synonym of *Stauropus sten*-

iptera Hampson, 1910, ibidem (8) 5: 471.

EPICERURINA Kiriakoff, 1969, Faune Madagascar 29: 87.

Type-species: *Epicerurina grisea* Kiriakoff, 1969, *ibidem* **29**: 87, text-figs 55, 56, pl.3, fig.29, by original designation. MADAGASCAR: southern, Tuléar province, Androy plateau, Antanimora.

EPICOMA Hübner, [1819] 1816, Verz. bekannter Schmett.: 160.

Type-species: *Epicoma tristis* Hübner, [1819] 1816, *ibidem*: 160 (a junior secondary homonym, see below), by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 441 (but designated as *contristis* Hübner, the objective replacement name, see below). [AUSTRALIA] Neuholland.

Hübner, [1819], Verz. bekannter Schmett.: 160, named as Epicoma tristis the unnamed species he had figured in [1819], Zuträge Samml. exot. Schmett. 2: pl.[38], figs 217, 218. In the relevant text not published until 1823, Zuträge Samml. exot. Schmett. 2: 9, Hübner named as Epicoma contristis the species he had figured on pl.[38], figs 217, 218. Thus Epicoma contristis Hübner, 1823, is a junior objective synonym of Epicoma tristis Hübner, [1819].

Epicoma tristis Hübner, [1819], is, however, a junior secondary homonym of Bombyx tristis Donovan, 1805 January 1, General Illust. Ent., An Epitome nat. Hist. Insects New Holland, New Zealand ... Pacific Ocean: pl.[34] (dated from the plate); a junior secondary homonym and a junior subjective synonym of Bombyx tristis Lewin, 1805 [December 31], Prodromus Ent., Nat. Hist. lepid. Insects New South Wales: 9, pl.8 (dated according to the Code, Article 21(b)(ii). Both Bombyx tristis Donovan and Bombyx tristis Lewin were included in Epicoma by Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 47, 48. The objective replacement name, Epicoma con-

tristis Hübner, 1823, is now the valid name for the type-species

of Epicoma Hübner, [1819].

Under the *Code*, Article 59(b)(i), a junior secondary homonym rejected before 1961 is permanently rejected and cannot be restored unless the use of the replacement name is contrary to existing usage.

EPICOMANA Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 42.

Type-species: *Epicoma asbolina* Turner, 1902, *Trans. Proc. R. Soc. S. Aust.* **26**: 183, by original designation. AUSTRALIA: Queensland, Townsville.

EPICORIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 33: 852.

Type-species: *Epicoria canosparsa* Walker, 1865, *ibidem* 33: 853, by subsequent designation by Berio, 1966, *Annali Mus. civ. Stor. nat. Giacomo Doria* 76: 57. [COLOMBIA]: Bogota.

E. canosparsa is a junior subjective synonym of Hemiceras

sparsipennis Walker, 1857, ibidem 12: 972.

Epicoria was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 333.

EPICOSMOLOPHA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 281.

Type-species: *Epicosmolopha dorsoflavida* Kiriakoff, 1964, *ibidem* 14: 281, abb.4, fig.40, pl.4, fig.43, by original designation. China: south Shensi, Tapaishan im Tsinling.

EPICRISIAS Dyar, 1912, Proc. U.S. natn. Mus. 42: 55.

Type-species: *Epicrisias eschara* Dyar, 1912, *ibidem* 42: 55, by original designation. MEXICO: Zacualpan.

EPIDESMA Hübner, [1819] 1816, Verz. bekannter Schmett.: 176.

Type-species: Phalaena ursula Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 107, pl.345, fig.E, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 180. SURINAM.

P. ursula is dated from the wrapper of the part and not from

the title-page of the volume, 1782.

TRICHODESMA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xix, 31, 451.

Type-species: *Phalaena ursula* Stoll, 1781, by original designation.

Trichodesma is a junior objective synonym of Epidesma Hübner, [1819].

A junior homonym of *Trichodesma* Leconte, 1861, *Classification Coleoptera N. Am.*: 204, – Coleoptera. The objective replacement name is *Epidesma* Hübner, [1819].

DESMOTRICHA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 396.

Type-species: *Phalaena ursula* Stoll, 1781, by original designation (for *Trichodesma* Hampson, 1898).

Desmotricha was proposed unnecessarily as the objective replacement name for *Trichodesma* Hampson, 1898.

EPIDONTA Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 555.

Type-species: *Epidonta eroki* Bethune-Baker, 1911, *ibidem* (8) 7: 556, by original designation. [Kenya]: Doenyo Erok [2°29'S 36°45'E].

EPIFENTONIA Kiriakoff, 1960, Bull. Inst. r. Sci. nat. Belg. **36** (17): 3.

Type-species: Fentonia variegata Wileman, 1910, Entomologist 43: 290, by original designation. TAIWAN: Kanshirei.

See also Disparia Nagano, 1916.

EPILACYDES Butler, 1875, Cistula ent. 2: 27. ARCT Type-species: Epilacydes simulans Butler, 1875, ibidem 2: 27, by original designation. WEST AFRICA.

EPILAIS Boisduval, 1870. See Dioptis Hübner, 1818. DIOP

‡EPIMEDIA; misspelling. See Epimydia Staudinger, 1892.

EPIMETULA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 156.

Type-species: *Desmeocraera albipuncta* Gaede, 1928, in Seitz, *Gross-Schmett. Erde* 14: 413, pl.69 row f, by original designation (for *Metula* Kiriakoff, 1958). [ZAIRE]: [upper] Lowa [valley, near Masisi].

Epimetula was established as the objective replacement name for Metula Kiriakoff, 1958.

METULA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 68.

Type-species: *Desmeocraera albipuncta* Gaede, 1928, by original designation.

A junior homonym of *Metula* Adams H. & Adams A., [1853] 1858, *Genera rec. Moll.* 1: 84, – Mollusca. The objective replacement name is *Epimetula* Kiriakoff, 1964.

EPIMOLIS Dyar, 1913, Proc. U.S. natn. Mus. 44: 286. ARCT Type-species: Epimolis zatrephica Dyar, 1913, ibidem 44: 286, by original designation. MEXICO: Misantla.

EPIMYDIA Staudinger, 1892, Dt. ent. Z. Iris 5: 346. ARCT Type-species: Epimydia dialampa Staudinger, 1892, ibidem 5: 346, pl.3, fig.2, by original designation. [U.S.S.R. or Mongolia]: Kentei Mts.

‡EPIMEDIA; Rothschild, 1910, Novit. zool. 17: 79.

An incorrect subsequent spelling of *Epimydia* Staudinger, 1892.

EPINOTODONTA Matsumura, 1919, Zool. Mag. Tokyo 31: 76.

Type-species: *Epinotodonta fumosa* Matsumura, 1919, *ibidem* 31: 76, by monotypy. JAPAN: [Nagano pref.], Yatsugatake.

Epinotodonta was again proposed by Matsumura, 1920, ibidem 32: 146 and 1921, Thousand Insects Japan (Additam.) 4: 796.

See also Yazawaia Marumo, 1920.

EPIPHALERA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 418. NOTO

Type-species: *Epiphalera uniformis* Gaede, 1928, *ibidem* 14: 418, pl.70, row b, by monotypy. [South Africa]: Transvaal.

E. uniformis is a junior subjective synonym of Stauropus pergrisea Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 472.

The date of publication of the part of Seitz containing page 418 is cited on page 417 as 29.iii.1928 in both the German and the English editions.

EPIPHALERA Gaede, 1930. See Epiphalerina Gaede, 1934.

EPIPHALERINA Gaede, 1934, in Strand, Lepid. Cat. 59: 198.

Type-species: *Phalera ochropis* Hampson, 1910, *J. Bombay nat. Hist. Soc.* 20: 90, pl.F, fig.8, by original designation (for *Epiphalera* Gaede, 1930). [SRI LANKA] Ceylon: Haputale.

Épiphalerina was established as the objective replacement name for *Epiphalera* Gaede, 1930.

EPIPHALERA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 614

Type-species: *Phalera ochropis* Hampson, 1910, by original designation.

A junior homonym of *Epiphalera* Gaede, 1928, *in* Seitz, *Gross-Schmett. Erde* 14: 418, – Lepid., Notodontidae. The objective replacement name is *Epiphalerina* Gaede, 1934.

The date of publication of the part of Seitz containing page 614 is cited on page 609 as 10.iii.1930 in both the German and the English editions.

EPIPTILODONTIS Kiriakoff, 1964, Bonn. zool. Beitr. 14: 256.

Type-species: *Epiptilodontis pterostomina* Kiriakoff, 1964, *ibidem* 14: 257, abb.2, fig.12, pl.1, fig.11, by original designation. China: Hunan, Hoengshan.

EPIPYROPS Bowring, 1852, in Westwood, Trans. ent. Soc. Lond. (2) 2 (Proc.): 5. EPIPYROPIDAE

Type-species: *Epipyrops anomala* Bowring, 1852, *ibidem* (2) **2** (Proc.): 5, by monotypy. Type-locality: not stated, [Hong Kong].

Epipyrops anomala was very briefly but sufficiently characterised for the names to be regarded under the Code as nomenclaturally available. Twenty-four years later the same genus and species were both much more fully described as new by Westwood, see below.

Epipyrops when established was not placed in a family; it was placed in the 'Arctiidarum', now Arctiidae, by Westwood, 1876, in Bowring & Westwood, Trans. ent. Soc. Lond. 1876: 522; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 490; it was transferred to the Limacodidae by Sharp, 1899, in Harmer & Shipley, Cambridge nat. Hist. 6: 404; and was placed in the Epipyropidae by Dyar, [1903] 1902, Bull. U.S. natn. Mus. 52: 359.

EPIPYROPS Westwood, 1876, in Bowring & Westwood, Trans. ent. Soc. Lond. 1876: 522.

Type-species: Epipyrops anomala Westwood, 1876, ibidem

1876: 522, pl.7, by monotypy. Hong Kong.

A junior homonym of *Epipyrops* Bowring, 1852, *Trans. ent.* Soc. Lond. (2) 2 (Proc.): 5, — Lepid., Epipyropidae. There is no objective replacement name but *E. anomala* Westwood, 1876, and *E. anomala* Bowring, 1852, although not objective synonyms based on the same type-specimen are certainly conspecific. *Epipyrops* Bowring, 1852, is therefore available for use as a subjective replacement name.

EPIPYROPS Westwood, 1876. See Epipyrops Bowring, 1852.

EPISCEA Warren, 1901, Novit. zool. 8: 440. PERI Type-species: Episcea extravagans Warren, 1901, ibidem 8: 441, by original designation. [BRAZIL]: Rio [de] Janeiro and Petropolis.

EPISCEPSIS Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: xi, 49. CTEN

Type-species: *Episcepsis venata* Butler, 1877, *ibidem* 1: xi, 49, pl.16, fig.7, by original designation. [BRAZIL]: [Amazonas], Rio Jutahi.

EPISCHAUSIA Kiriakoff, 1976, Tierreich 98: 75.

Type-species: Aegocera dispar Rothschild, 1896, in Rothschild Jordan, Novit. zool. 3: 43, pls 27, 28, by original designation. Syntypes 18, 39, [Kenya]: [Wasin 1.] Wassein, (BMNH).

Epischausia in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

EPISEMA Ochsenheimer, 1816, Schmett. Eur. 4: 65.

Type-species: *Bombyx glaucina* Esper, 1789, *Die Schmett*. 3 (Suppl.) Abschnitt 3: 11, pl.81, figs 4, 5, by subsequent

designation by the International Commission on Zoological Nomenclature, 1957, *Opin. Decl. int. Commn zool. Nom.* 17 (Opinion 494): 267. GERMANY: Frankfurt am Main.

Further information in Nye, 1975, Generic Names Moths World 1: 181.

CLADOCERA Rambur, 1858, Cat. syst. Lépid. Andalousie (1): pl.6, fig.4, pl.8, fig.1.

Type-species: *Episema hispana* Rambur, 1848, *Annls Soc. ent. Fr.* (2) **6**: 66 (a junior homonym replaced by *E. ramburi* Zerny, 1927; see below), by monotypy. Spain, Algeria.

Cladocera Rambur is not preoccupied by ‡Cladocera Blainville, 1834, Manuel Actinologie: 670, an incorrect subsequent spelling of Cladocora Ehrenberg & Hemprich, [1834] 1832, Abh. preuss. Akad. Wiss. 1832: 309.

Cladocera Rambur is, however, a junior homonym of Cladocera Hope, 1840, Coleopt. Manual 3: 169, – Insecta, Coleoptera. There is no objective replacement name, but E. hispana is congeneric with Bombyx glaucina Esper, 1789, the typespecies of Episema Ochsenheimer, 1816. The latter is therefore available for use as a subjective replacement name.

E. hispana is a junior primary homonym of Episema trimacula hispana Boisduval, 1834, Icones Hist. Lépid. 2: pl.72, figs 4, 5 – Lepid., Noctuidae. The objective replacement name is Episema ramburi Zerny, 1927, Eos, Madr. 3: 374.

Cladocera in the Noctuidae, Cuculliinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

EPISTAUROPUS Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 625.

Type-species: Stauropus apiculatus Rothschild, 1917, Novit. zool. 24: 245, by original designation. [Malaysia]: Penang.

The date of publication of the part of Seitz containing page 625 is cited on that page as 11.iii.1930 in both the German and English editions.

EPISTEME Hübner, [1820] 1816, Verz. bekannter Schmett.: 180. AGARISTIDAE

Type-species: *Phalaena lectrix* Linnaeus, 1764, *Mus. Ludovicae Ulricae Reginae*: 389, by monotypy. CHINA.

Further information in Nye, 1975, Generic Names Moths World 1: 182.

Episteme has been used as the original generic name for species now placed in the Arctiidae.

‡EPYSTEME; Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 31.

An incorrect subsequent spelling of *Episteme* Hübner [1820]. Hübner, on page 27, established the nominal species '254. *Episteme isse*'. On page 31 he established '265. *Epysteme turbida*' also referring to the genus *Episteme*. On page 43, in the index to genera, species numbers 254 and 265 are both cited as *Episteme*. ‡*Epysteme* must therefore be a misspelling, though Hemming, 1937, *Hübner* 2: 189 treated it as an available name

‡Epysteme is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

EPITALARA Dyar, 1905, Proc. U.S. natn. Mus. 29: 175.

Type-species: *Epitalara reversa* Schaus, 1905, *ibidem* **29**: 205, by original designation (in anticipation of Schaus's description of *reversa*). French Guiana: St Jean.

Epitalara was available but was without an originally included nominal species between its date of publication on 3rd October, 1905, and that of its type-species on 11th October, 1905, by Schaus.

‡EPITAXIS; misspelling. See Epitoxis Wallengren, 1863.

CTEN

EPITHISANOTIA Kiriakoff, 1976, Tierreich 99: 32.

Type-species: Eudryas sanctaejohannis Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 9: 144, by original designation (but cited as E. sanctijohannis Stephens, a nomen nudum; see below). Holotype ô, Type-locality: stated to be [Great Britain]: [England], Horsleydown, but all specimens other than the holotype are from tropical America, (BMNH).

E. sanctaejohannis was proposed as ‡Stae. Johannis which under the Code, Article 26(d), must be written in full and both

words united.

E. sanctaejohannis was first known as ‡E. Sti. Johannis Stephens, 1850, List Specimens lepid. Insects Colln Br. Mus. 5: 312, a nomen nudum based on the same specimen as E. sanctaejohannis Walker but descibed only as 'an exquisitely beautiful nondescript species of Eudryas'.

Epithisanotia in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

EPITOXIS Wallengren, 1863, Wien. ent. Monatschr. 7: 138.

Type-species: *Thyretes amazoula* Boisduval, 1847, *in* Delegorgue, *Voyage Afr. australe* 2: 597, by monotypy. [SOUTH AFRICA]: 'Pays des Amazoulous'.

Epitoxis was described again as new by Wallengren, 1865, K. svenska Vetensk Akad. Handl. 5 (4): 11.

‡EPITAXIS; Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1866.

An incorrect subsequent spelling of *Epitoxis* Wallengren, 1863.

‡EPATOSIS; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling of *Epitoxis* Wallengren, 1863.

EPITROTONOTUS Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 121.

Type-species: Ctenogyna vilis Holland, 1893, Ent. News 4: 343, by original designation. [GABON]: Kangwé [on right bank of Ogooué river just south of equator on 10°50'E].

EPIZARANGA Matsumura, 1920 May 15, Zool. Mag. Tokyo 32: 143.

Type-species: Drymonia permagna Butler, 1881, Trans. ent. Soc. Lond. 1881: 20, by monotypy. JAPAN: Tokyo.

DENSITAS Marumo, 1920 November, J. Coll. Agric. imp. Univ. Tokyo 6: 281 (key), 348.

Type-species: *Drymonia permagna* Butler, 1881, by original designation.

A junior objective synonym of *Epizaranga Matsumura*, 1920 May 15.

EPODONTA Matsumura, 1922, Zool. Mag. Tokyo 34 (402): 522.

Type-species: *Notodonta lineata* Oberthür, 1880, *Etudes d'Entomologie* 5: 61, pl.2, fig.7, by original designation. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

‡EPYSTEME; misspelling. See Episteme Hübner, [1820].

AGARISTIDAE

ERAGISA Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1656.

Type-species: *Eragisa lanifera* Walker, 1858, *ibidem* 15: 1657, by monotypy. [Brazil]: Amazon region.

Eragisa was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901:

ERAMOS Kiriakoff, 1958, Mém. Inst. scient. Madagascar (E) 9: 220.

Type-species: *Eramos viridissima* Kiriakoff, 1958, *ibidem* (E) 9: 222, text-fig.4, pl.4, fig.4, by original designation. MADAGASCAR: Tananarive.

ERASTA Gistl, 1848. See Penthophera Germar, 1812.

ERBESSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 319.

Type-species: *Dioptis sobria* Walker, 1854, *ibidem* 2: 319, by monotypy. [Brazil]: Para.

MYONIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 308.

Type-species: Josia evippe Walker, 1854, ibidem 2: 309, by monotypy. Brazil: Tapajos.

A junior homonym of *Myonia* Dana, 1847, *Ann. J. Sci.* (2) 4: 158, – Fossil Mollusca. There is no objective replacement name, but *J. evippe* is currently treated as congeneric with *Dioptis sobria* Walker, 1854, the type-species of *Erbessa* Walker, 1854; the latter is therefore available for use as a subjective replacement name.

ERCHIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 321.

Type-species: *Dioptis semistria* Walker, 1854, *ibidem* 2: 321, by PRESENT DESIGNATION. BRAZIL.

Erchia was established as a subgenus of Dioptis Hübner, 1818.

ERCONHOLDA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 220.

Type-species: Fentonia mangholda Schaus, 1928, Proc. U.S. natn. Mus. 73 (19): 79, by original designation. Philippines: [Mindanao], Surigao [labelled type 8], Luzon, [Mt Makiling, labelled type 9].

Clarification of the type-locality data was received from Dr

D.C. Ferguson, U.S. National Museum, 10.ix.1977.

ERESSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 149.

Type-species: *Glaucopis confinis* Walker, 1854, *ibidem* 1: 149, by monotypy. INDIA: North.

Eressa was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 115.

ERESSADES Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 531.

Type-species: *Eressades flavipunctata* Bethune-Baker, 1911, *ibidem* (8) 7: 531, by original designation. ANGOLA: N., N'Dalla Tando.

E. flavipunctata is a junior subjective synonym of Thyretes monteiroi Butler, 1876, J. Linn. Soc. (Zool.) 12: 359.

Eressades was established in the Syntomidae, now Ctenuchidae; its type-species was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 21.

ERIKA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 231.

Type-species: *Erika analalava* Griveaud, [1976] 1975, *ibidem* 80: 232, by original designation. MADAGASCAR: N., 37km S. de Vohemar, foret d'Analalava.

ERILYCES Warren, 1895, Novit. zool. 2: 85.

Type-species: Lyces flavissima Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 367, by original designation. VENEZUELA.

Erilyces was established in the Geometridae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 420.

‡ERIOCRYPTA Herrich-Schäffer, 1855. See Aganopis Herrich-Schäffer, [1856]. AGAN

ERIOMASTYX Rothschild & Jordan, 1905, Novit. zool. 12: 469.

Type-species: Eriomastyx latus Rothschild & Jordan, 1905, ibidem 12: 469, by original designation. New Guinea: [Papua New Guinea] British New Guinea.

ERIOPANA Kiriakoff, 1974. See Pristoceraea Karsch, 1895. AGARISTIDAE

ERIOSTEPTA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 2, 30.

Type-species: Eriostepta roseireta Hampson, 1901, ibidem 3: 31, fig.23, by original designation. [BRAZIL]: [Amazonas] Olivença[s].

ERIPHIA Herrich-Schäffer, 1855. See Antichloris Hübner, 1818.

ERIPHIOIDES Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 159.

Type-species: Eriphia ustulata Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.17, by original designation. [Colombia]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

ERITHALES Poey, 1832, Centurie Lépid. Cuba (2): 12.

Type-species: Erithales guacolda Poey, 1832, ibidem (2): text, pl.[14], by monotypy. CUBA.

ERNASSA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1643.

Type-species: Phalaena justina Stoll, 1782, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 186, 250, pl.383, fig.E, by monotypy. SURINAM.

P. justina is a junior subjective synonym of Phalaena sanguinolenta Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 102, 176, pl.252, fig.A.

ERNOLATIA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 131. BOMBYCIDAE

Type-species: Ernolatia signata Walker, 1862, ibidem 6: 131, by monotypy.

Ernolatia was established in the Liparidae, now Lymantriidae; it was placed in the Bombycidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 716.

ERRUCA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 165.

Type-species: Glaucopis devrolii Walker, 1854, ibidem 1: 165, by monotypy. [South America]: [tropical], 'Rio Grande'.

Erruca was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 225.

ERYTHROCLOSTERA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 255.

Type-species: Ichthyura castanea Rothschild, 1915, Lepid. Br. Orn. Un. & Wollaston Expdns Snow Mts sth. Dutch New Guinea: 80, pl.2, fig.60, by original designation. New Guinea: [Irian Jaya], Snow Mts.

ESTHEMA Hübner, [1820] 1816, Verz. bekannter Schmett.: 178.

Type-species: Phalaena bicolora Sulzer, 1776, Geschichte Insecten: 161, pl.22, fig.6, by subsequent designation by Kirby,

1892, Synonymic Cat. Lepid. Heterocera 1: 191. [INDIA] Indien.

See also Hypocrita Hübner [1807].

‡ECTHEMA; Burmeister, 1878, Descr. phys. Répub. Argent.

An incorrect subsequent spelling of Esthema Hübner, [1820].

ESTIGMENE Hübner, [1820] 1816, Verz. bekannter Schmett.:

Type-species: Phalaena acrea Drury, 1773, Illust. nat. Hist. exot. Insects 2: index to 1; 1770, ibidem 1: 7, pl.3, fig.2 (but cited by Hübner as ‡'acria Abb.', an incorrect authorship and incorrect subsequent spelling), by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 226 (but cited as acraea, an unjustified emendation by Kirby, 1892). [U.S.A.]: New York; Maryland; and Virginia.

LEUCARCTIA Packard, 1864, Proc. ent. Soc. Philad. 3: 124. Type-species: Phalaena acrea Drury, 1773, by subsequent designation by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 338 (but cited as acraea, an unjustified emendation by Kirby,

A junior objective synonym of Estigmene Hübner, [1820].

ETOBEMA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 388.

Type-species: Etobema circumdata Walker, 1865, ibidem 32: 388, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 497. New Guinea.

‡ETOBENA; Dallas, 1865, in Günther, Zool. Rec. 2: 611. An incorrect subsequent spelling of *Etobema* Walker, 1865.

‡ETOBENA; misspelling. See Etobema Walker, 1865.

ETOBESA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 471.

Type-species: Etobesa ligneata Walker, 1865, ibidem 32: 471, by monotypy. [Brazil]: [R. Amazon], [Tefé] Ega.

EUAGRA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 323.

Type-species: Dioptis azurea Walker, 1854, ibidem 2: 325, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 179. Brazil.

Euagra was established as a subgenus of Dioptis Hübner, 1818, - Lepid., Dioptidae.

CALLAGRA Butler, 1876, Cistula ent. 2: 111.

Type-species: Dioptis azurea Walker, 1854, by original designation.

A junior objective synonym of Euagra Walker, 1854.

EUANTHIA Kiriakoff, 1962, Tijdschr. Ent. 105: 259. NOTO Type-species: Euanthia venosa Kiriakoff, 1962, ibidem 105: 259, fig.35, by original designation. [TANZANIA] Tanganyika: Shinyanga.

EUARCTIA Staudinger, 1881, Stettin. ent. Ztg 48: 79. ARCT Type-species: Euarctia proserpina Staudinger, 1887, ibidem 48: 80, by monotypy. [U.S.S.R.]: Samarkand Province.

EUBAPHE Hübner, 1823, Zuträge Samml. exot. Schmett. 2:

Type-species: Eubaphe lobula Hübner, 1823, ibidem 2: 20,

pl.[52], figs 299, 300, by monotypy. Brazil.

Eubaphe was established in the Arctiidae (by inference) but is currently placed in the Geometridae. Several species of Arctiidae previously placed in Eubaphe are currently placed in Holomelina Herrich-Schäffer, 1855.

EUBREYERIA Fletcher nom. n. for Breyeria Janse, 1920.

Type-species: *Breyeria dasychiroides* Janse, 1920, *Ann. Transv. Mus.* 7: 214, pl.5, fig.16, pl.8, figs 20-25, pl.14, fig.10, by original designation (for *Breyeria* Janse, 1920). SOUTH AFRICA: [Transvaal], Nelspruit.

BREYERIA Janse, 1920, Ann. Transv. Mus. 7: 213.

Type-species: *Breyeria dasychiroides* Janse, 1920, by original designation.

A junior homonym of *Breyeria* Borre, 1875, *Annls Soc. ent.* Belg. (C.r.) 18: lx, – Palaeodictyoptera. The objective replacement name is *Eubreyeria* Fletcher nom. n.

‡EUCEREA; misspelling. See Eucereon Hübner, [1819].

‡EUCEREEON; misspelling. See Eucereon Hübner, [1819].

EUCEREON Hübner, [1819] 1816, Verz. bekannter Schmett.: 123.

Type-species: Sphinx archias Stoll, 1790, in Cramer, Uitlandsche Kapellen (Papillons exot.) (Aanhangsel): 66, pl.14, figs 6, 7A, 8B, 9C, 10D, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 200. SURINAM.

S. archias is dated from the wrapper of the part and not from the title-page of the volume, 1791.

‡EUCEREEON; Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 265.

An incorrect subsequent spelling of *Eucereon* Hübner, [1819].

‡EUCEREA; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1638.

An incorrect subsequent spelling of Eucereon Hübner, [1819].

EUCEREUM Zerny, 1912, in Wagner, Lepid. Cat. 7: 137. An unjustified emendation of Eucereon Hübner, [1819].

EUCEREUM Zerny, 1912. See Eucereon Hübner, [1819].

EUCERIODES Travassos, 1961, Atas Soc. Biol. Rio de J. 5:

Type-species: *Eucereum wernickei* Draudt, 1917, *în* Seitz, *Gross-Schmett. Erde* 6: 213, pl.30, fig.h, by original designation. BRAZIL: Sta Catarina.

EUCERURA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 260 (key), 294.

Type-species: *Drymonia pica* Butler, 1882, *ibidem* **1882**: 22, by original designation. CHILE.

EUCHAETES Harris, 1841, Rep. Insects Mass. injurious to Vegn: 257.

Type-species: *Phalaena egle* Drury, 1773, *Illust. nat. Hist. exot. Insects* 2: index, 36, pl.20, fig.3, by monotypy. [U.S.A.]: New York.

EUCHAETIAS Lyman, 1902, Can. Ent. 34: 27.

An unnecessary replacement name for *Euchaetes* Harris, 1841, which is not a junior homonym of ‡*Euchaetes* Dejean, [1835], *Cat. Coleoptera* (Edn 2): 340, an unavailable name under the *Code* Article 12, as it was published without description and contained only one species and a variety of it, both also undescribed. The generic name *Euchaetes* was not used again before *Euchaetes* Harris, 1841.

EUCHAETIAS Lyman, 1902. See Euchaetes Harris, 1841.

EUCHARIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 181. ARCT

Type-species: *Phalaena hebe* Linnaeus, 1767, *Syst. Nat.* (Edn 12) 1 (2): 820, by subsequent designation by Kirby, 1880, *Zool. Rec.* 1878: 197. GERMANY.

P. hebe is a junior subjective synonym of Phalaena festiva Hufnagel, 1766, Berl. Mag. 2: 416, 437, — Lepid., Arctiidae.

AMMOBIOTA Wallengren, 1885, Skand. Heterocer-Fjärilar 2: 304.

Type-species: *Phalaena hebe* Linnaeus, 1767, by monotypy. A junior objective synonym of *Eucharia* Hübner, [1820].

EUCHELIA Boisduval, 1828. See Tyria Hübner, [1819].

EUCHILA Billberg, 1820, Enumeratio Insect. Mus. G.J. Billberg: 84.

Type-species: *Phalaena palpina* Clerck, 1759, *Icon. Insect. rariorum*: pl.9, fig.8 (but included by Billberg as *palpina* Linn., an incorrect authorship), by monotypy. Type-locality: not stated, [Europe].

See also Pterostoma Germar, 1812.

ORTHORINIA Boisduval, 1828, Eur. Lepid. Index meth.: 56. Type-species: Phalaena palpina Clerck, 1759 (but included by Boisduval as 'palpina L.', an incorrect authorship), by monotypy.

A junior objective synonym of Euchila Billberg, 1820.

PTILODONTIS Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 28.

Type-species: *Phalaena palpina* Clerck, 1759, by monotypy. A junior objective synonym of *Euchila* Billberg, 1820.

Stephens incorrectly attributed *Ptilodontis* to Hübner and *palpina* to Linnaeus in 1828, *ibidem* 2: 28, and again in 1829 [June], *Nom. Br. Insects*: 39. In 1829 [July], *Syst. Cat. Br. Insects* (2): 43, Stephens proposed *Ptilodontis* as his own new genus.

ODONPTERA Latreille, 1829, in Cuvier, Règne Animal (Edn 2) 5: 409.

Type-species: *Phalaena palpina* Clerck, 1759, by monotypy. A junior objective synonym of *Euchila* Billberg, 1820.

ORTHORHINA Meigen, 1830, Syst. Beschreibung eur. Schmett. 2: 184.

An unjustified emendation of Orthorinia Boisduval, 1828.

‡PHILODONTIS; Blanchard, 1840, in Castelnau, Hist. nat. Insectes 3: 495.

An incorrect subsequent spelling of *Ptilodontis* Stephens, 1828.

ORTHORHINIA Agassiz, 1847, Nomencl. zool. (Index univl.): 265.

An unjustified emendation of Orthorinia Boisduval, 1828.

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Orthorhinia* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

‡PULODONTIS; Butler, 1885, Cistula ent. 3: 127.

An incorrect subsequent spelling of *Ptilodontis* Stephens, 1828.

‡ORTHORINA; Gaede, 1934, in Strand, Lepid. Cat. 59: 168. An incorrect subsequent spelling of Orthorhina Meigen, 1830.

EUCHLAENIDIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 5, 186.

Type-species: *Phaeochlaena transcisa* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 2: 468, by original designation. Type-locality: not known, [Brazil].

EUCHLOROSTOLA Watson nom. n. for Chlorostola Hampson, 1898.

Type-species: *Eupyra corydon* Druce, 1884, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 1: 44, pl.6, fig.13, by original designation (for *Chlorostola* Hampson, 1898). GUATEMALA: Senahu.

CHLOROSTOLA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xviii, 26, 427.

Type-species: *Eupyra corydon* Druce, 1884, by original designation.

A junior homonym of *Chlorostola* Wiese, 1881, *Naturg. Insecten Dtl.* (1) 6: 99, – Coleoptera. The objective replacement name is *Euchlorostola* Watson nom. n.

EUCHONTHA Walker, 1865 [April 29], List Specimens lepid. Insects Colln Br. Mus. 32: 383.

Type-species: *Euchontha sublactigera* Walker, 1865 [April 29], *ibidem* 32: 383, by monotypy. [Colombia]: Bogota.

E. sublactigera is a junior subjective synonym of Devara frigida Walker, [1865 February 11] 1864, ibidem 31: 185.

See also Gnatholophia Felder, 1874, and Macroneurodes Warren, 1900.

‡EUCONTHA; Stretch, 1875, Cistula ent. 2: 18.

An incorrect subsequent spelling of Euchontha Walker, 1865.

EUCHROMIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 121.

Type-species: Sphinx sperchius Cramer, 1777, Uitlandsche Kapellen (Papillons exot.) 2: 79, 150, pl.146, fig.C, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 117. SIERRA LEONE.

S. sperchius is dated from the wrapper of the part and not from the title-page of the volume, 1779. S. sperchius is a junior subjective synonym of Zygaena guineensis Fabricius, 1775, Syst. Ent.: 551.

HIRA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 220.

Type-species: Sphinx sperchius Cramer, 1777, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 293.

A junior objective synonym of *Euchromia* Hübner, [1819]. *Hira* was established as a subgenus of *Euchromia* Hübner, [1819], but this was before the type-species of *Euchromia* had been designated.

EUCLEMENSIA Dognin, 1914 February. See Euclemensoides Strand, 1922.

EUCLEMENSIA Hampson, 1914 [November]. See Euclemensoides Strand, 1922.

EUCLEMENSOIDES Strand, 1922, in Wagner, Lepid. Cat. 26: 739.

Type-species: Euclemensia umbrata Dognin, 1914, Hétérocères nouv. Am. Sud 7: 14, by monotypy (of Euclemensia Dognin, 1914). COLOMBIA: Ouindin, Rio Toche.

Euclemensoides was established as the objective replacement name for Euclemensia Dognin, 1914.

EUCLEMENSIA Dognin, 1914 February, Hétérocères nouv. Am. Sud 7: 14.

Type-species: Euclemensia umbrata Dognin, 1914, by monotypy.

A junior homonym of *Euclemensia* Grote, 1878, *Can. Ent.* 10: 69, – Lepid., Heliodinidae. The objective replacement name is *Euclemensoides* Strand, 1922.

The name *Euclemensia* probably originated from Hampson but was used and made nomenclaturally available by Dognin prior to its proposal as a new generic name by Hampson, 1914 [November].

EUCLEMENSIA Hampson, 1914 [November], Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxv, 702.

Type-species: Euclemensia umbrata Dognin, 1914, by original designation.

A junior homonym of *Euclemensia* Grote, 1878, *Can. Ent.* **10**: 69, – Lepid., Heliodinidae. The objective replacement name is *Euclemensoides* Strand, 1922.

Euclemensia Hampson, 1914 [November], is also a junior homonym and junior objective synonym of Euclemensia Dognin, 1914 February.

EUCLERA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 102, 103 (but attributed to Boisduval, an incorrect authorship).

Type-species: *Sphinx meones* Stoll, 1780, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 73, pl.325, fig.E, by monotypy. SURINAM.

S. meones is dated from the wrapper of the part and not from the title-page of the volume, 1782.

Euclera was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

ANDROCHARTA Felder, C. & Felder, R., 1862, Wien. ent. Monatschr. 6: 231.

Type-species: Sphinx meones Stoll, 1780, by monotypy.

A junior objective synonym of *Euclera* Herrich-Schäffer, 1855.

EUCOCYTIA Rothschild & Jordan, 1905, Novit. zool. 12: 474.

Type-species: *Eucocytia meeki* Rothschild & Jordan, 1905, *ibidem* 12: 474, by original designation. New GUINEA.

EUCONOSIA Watson nom. n. LITH Type-species: Lithosia aspersa Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 104, by original designation (for Conosia Hampson, 1900). BORNEO.

CONOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xx, 83, 542.

Type-species: Lithosia aspersa Walker, 1862, by original designation.

A junior homonym of *Conosia* Van der Wulp, 1880, *Tijdschr*. *Ent.* 23: 159, – Insecta, Diptera. The objective replacement name is *Euconosia* Watson nom. n.

‡EUCONTHA; misspelling. See Euchontha Walker, 1865.

‡EUCREAGA; misspelling. See Eucreagra Strand, 1920.

EUCREAGRA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.33. LITH

Type-species: Eucreagra arculifera Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.33, by monotypy. [SOUTH AFRICA]: Knysna, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

‡EUCREAGA; Strand, 1920, in Wagner, Lepid. Cat. 24: 532. An incorrect subsequent spelling of Eucreagra Felder, 1874.

EUCTENUCHA Grote, 1873, Bull. Buffalo Soc. nat. Sci. 1: 33.

Type-species: Apistosia multifaria Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 479, by original designation (but attributed to Grote & Robinson, an incorrect authorship). [U.S.A.]: California.

EUCYANE Hübner, [1820] 1816, Verz. bekannter Schmett.: 178.

Type-species: *Phalaena celadon* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 55, 148, pl.132, fig.E, by subsequent designation by Bryk, 1931, *in* Wagner, *Lepid. Cat.* 45: 18. SURINAM.

P. celadon is dated from the wrapper of the part and not from the title-page of the volume, 1779.

EUCYCLOPERA Hampson, 1895, Trans. ent. Soc. Lond. 1895: 294.

Type-species: Eucyclopera plagidisca Hampson, 1895, ibidem 1895: 294, by original designation. BHUTAN.

EUCYRTA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.4. ARCT

Type-species: *Eucyrta albicollis* Felder, *ibidem* **2** (Abt.2): pl.102, fig.4, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* **1**:171. [BRAZIL]: [R. Amazon] fl. Amazonas, cited by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 4.

EUDASYCHIRA Möschler, 1887, Abh. senckenb. naturforsch. Ges. 15: 75.

Type-species: *Eudasychira quinquepunctata* Möschler, 1887, *ibidem* **15**: 75, fig.11, by monotypy. [GHANA] Goldküste: Accra.

EUDESMIA Hübner, [1823], Samml. exot. Schmett. 2: pl.[187].

Type-species: *Eudesmia ruficollis* Donovan, 1798, *Epitome nat. Hist. Insects China*: pl.[41], by monotypy. Type-locality: cited as China by Donovan but his specimens probably originated from tropical SOUTH AMERICA.

EUDIOPTIS Prout, 1918. See Paradioptis Hering, 1925.

EUDMOE Hübner, [1819] 1816, Verz. bekannter Schmett.: NOTO

Type-species: *Phalaena arne* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 23, pl.15, fig.E, by monotypy. SURINAM.

P. arne is dated from the wrapper of part 2, not from the title-page of volume 1 which is dated 1779.

EUDOGNINA Fletcher nom. n. for Libyssa Dognin, 1911.

Type-species: Libyssa viridirosea Dognin, 1911, Hétérocères nouv. Am. Sud 2: 42, by original designation (for Libyssa Dognin, 1911). FRENCH GUIANA.

LIBYSSA Dognin, 1911, Hétérocères nouv. Am. Sud 2: 41.

Type-species: Libyssa viridirosea Dognin, 1911, by original designation.

A junior homonym of Libyssa Dallas, 1851, List Specimens hemipt. Insects Colln Br. Mus. 1: 20, — Insecta, Hemiptera. The objective replacement name is Eudognina Fletcher nom. n.

EUDOLICHE Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 660.

Type-species: *Eudoliche vittata* Möschler, 1878, *ibidem* **27** (Abh.): 660, pl.9, fig.28, by monotypy. Surinam: interior of, and Paramaribo.

EUERYTHRA Harvey, 1876, Can. Ent. 8: 5. ARCT Type-species: Euerythra phasma Harvey, 1876, ibidem 8: 5, by original designation. U.S.A.: Texas.

‡ENERYTHRA; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 235.

An incorrect subsequent spelling of Euerythra Harvey, 1876.

EUFENTONIA Matsumura, 1922, Zool. Mag. Tokyo 34: 521.

Type-species: Fentonia nihonica Wileman, 1911, Trans. ent. Soc. Lond. 1911: 286, pl.30, fig.5, by original designation. Japan: Yamato prov., Yoshino.

EUFORBESIA Kiriakoff, 1950, Bull. Annls Soc. r. ent. Belg. 86: 77.

Type-species: *Tanaostyla unimacula* Warren, 1897, *Novit. zool.* 4: 423, by original designation. BOLIVIA: Reyes.

EUGOA Walker, [1858] 1857, List Specimens lepid. Insects Colln Br. Mus. 12: 768.

Type-species: Eugoa aequalis Walker, [1858] 1857, ibidem 12: 768, by monotypy. BORNEO.

EUGONOSIA Schaus, 1899, Jl N.Y. ent. Soc. 7: 214. LITH Type-species: Eugonosia angulifer Schaus, 1899, ibidem 7: 214, by monotypy. [BRAZIL]: Parana, Castro.

The name Eugonosia probably originated from Hampson, but was used and made nomenclaturally available by Schaus prior to its proposal as a new generic name by Hampson, 1900.

EUGONOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 87, 295.

Type-species: Eugonosia angulifer Schaus, 1899, by original designation.

A junior homonym of *Eugonosia* Schaus, 1899. The objective replacement name is *Eugonosia* Schaus, 1899.

EUGONOSIA Hampson, 1900. See Eugonosia Schaus, 1899.

EUGRAPHOSIA Hampson, 1918, Novit. zool. 25: 103. LITH Type-species: Eugraphosia rubrizonea Hampson, 1918, ibidem 25: 104, by original designation. PERU: Contamino, R. Ucayale.

EUHALISIDOTA Grote, 1865, Proc. ent. Soc. Philad. 5: 243.

Type-species: *Euhalisidota luxa* Grote, 1865, *ibidem* 5: 244, pl.4, fig.5, by monotypy. CUBA.

E. luxa is a junior subjective synonym of Halysidota atomosa Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 746 (but proposed as ‡Halesidota, an incorrect subsequent spelling).

EUHAMPSONIA Dyar, 1897, Trans. Am. ent. Soc. 24: 16.

Type-species: *Trabala niveiceps* Walker, 1865, *List Specimens lepid. Insects Colln Br. Mus.* 32: 554, by original designation. [INDIA]: North Hindostan.

EUHARPYIA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 263 (key), 324.

Type-species: *Euharpyia comita* Schaus, 1901, *ibidem* 1901: 324, pl.12, fig.4, by original designation. PERU: Chanchamayo.

‡EUPHARPYIA; Neave, 1939, Nomencl. zool. 2: 343.

An incorrect subsequent spelling of Euharpyia Schaus, 1901. Neave attributed this misspelling to Dognin, 1911, Hétérocères nouv. Am. Sud 2: 39, but at this reference Euharpyia was correctly spelt.

EUHYPARPAX Beutenmüller, 1893, Bull. Am. Mus. nat. Hist. 5: 19.

Type-species: *Euhyparpax rosea* Beutenmüller, 1893, *ibidem* 5: 19, by monotypy. [U.S.A.]: Colorado, Custer Co., West Cliff.

Euhyparpax was established in the Bombycidae; it was placed in the Notodontidae by Neumoegen & Dyar, 1894, Trans. Am. ent. Soc. 21: 204.

EUIMATA Billberg, 1820, Enumeratio Insect. Mus. G.J. Billberg: 91.

Type-species: *Phalaena alciphron* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 58, [147], pl.133, fig.E, by mon-

otypy. [India]: [S.], Tranquebar.

P. alciphron is dated from the wrapper of the part and not from the title-page of the volume, 1779. P. alciphron is a junior subjective synonym of Noctua caricae Fabricius, 1775, Syst. Ent.: 596.

See also Psephea Billberg, 1820.

EUJANSEA Fletcher nom. n. for Pectinophora Janse, 1920.

Type-species: *Pectinophora noctuiformis* Janse, 1920, *Ann. Transv. Mus.* 7: 165, pl.1, fig.7, pl.2, figs 11-17, pl.13, fig.7, by original designation (for *Pectinophora* Janse, 1920). [ZIMBABWE-]RHODESIA: Umtali.

PECTINOPHORA Janse, 1920, Ann. Transv. Mus. 7: 164.

Type-species: *Pectinophora noctuiformis* Janse, 1920, by original designation.

A junior homonym of *Pectinophora* Busck, 1917, *J. agric.* Res. 9: 346, – Lepid., Gelechiidae. The objective replacement name is *Eujansea* Fletcher nom. n.

EULAVINIA Fletcher nom. n. for Lavinia Kiriakoff, 1962.

Type-species: *Phalera lavinia* Fawcett, 1916, *Proc. zool. Soc. Lond.* **1916**: 725, pl.1, fig.20, by original designation (for *Lavinia* Kiriakoff, 1962). [Kenya]: Kedai.

LAVINIA Kiriakoff, 1962, Revue Zool. Bot. afr. 66: 16.

Type-species: *Phalera lavinia* Fawcett, 1916, by original designation.

A junior homonym of Lavinia Girard, 1854, Proc. Acad. nat. Sci. Philad. 7: 137, – Pisces. The objective replacement name is Eulavinia Fletcher nom. n.

EULEECHIA Dyar, 1900, Can. Ent. 32: 347.

Type species: Chalenia histi Obothiir, 1883, Bull. See and

Type-species: *Chelonia bieti* Oberthür, 1883, *Bull. Soc. ent. Fr.* (6) 3: xliii, by original designation (for *Callarctia* Leech, 1899). China: [Szechwan], Ta-tsien-lou.

Euleechia was established as the objective replacement name for Callarctia Leech, 1899.

CALLARCTIA Leech, 1899, Trans. ent. Soc. Lond. 1899: 168. Type-species: Chelonia bieti Oberthür, 1883, by original designation.

A junior homonym of *Callarctia* Packard, 1864, – Lepid., Arctiidae. The objective replacement name is *Euleechia* Dyar, 1900.

NEOCHELONIA Draeseke, 1926, Dt. ent. Z. Iris 40: 47.

Type-species: Chelonia bieti Oberthür, 1883, by subsequent designation by Draudt, 1932, Gross-Schmett. Erde 2 (Suppl.): 89.

A junior objective synonym of Euleechia Dyar, 1900.

EULEPIA Curtis, 1825. See Spiris Hübner, [1819]. ARCT

EULOPHOPTERYX Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 684.

Type-species: *Eulophopteryx splendens* Möschler, 1878, *ibidem* **27** (Abh.): 684, pl.10, fig.42, by monotypy. Surinam.

E. splendens is a junior subjective synonym of *Phalaena* meona Stoll, 1781, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 132, pl.358, fig.B. *P. meona* is dated from the wrapper of part 30, not from the title-page of the volume, dated 1782.

EUMASCHANE Schaus, 1905, Proc. U.S. natn. Mus. 29: 241.

Type-species: *Eumaschane laura* Schaus, 1905, *ibidem* **29**: 241, by monotypy. French Guiana: St Laurent.

EUMELIA Neumoegen, 1893, Can. Ent. 25: 25.

Type-species: Melia danbyi Neumoegen, 1892, ibidem 24: 225, by monotypy (of Melia Neumoegen, 1892). [CANADA]:

British Columbia, Victoria.

Eumelia was established as the objective replacement name for Melia Neumoegen, 1892. Eumelia Neumoegen is not preoccupied by ‡Eumelia Duncan [& Westwood], 1841, in Jardine, Naturalist's Libr. (Edn 1) 33 (Ent.7): pl.29, fig.4, legend, nor by ‡Eumelia Hopffer, 1874, Stettin. ent. Ztg 35: 47; the former is an incorrect (of a multiple) original spelling, the latter an incorrect subsequent spelling of Eumelea Duncan [& Westwood], 1841, ibidem 33 (Ent.7): 215, — Lepid., Geometridae.

MELIA Neumoegen, 1892, Can. Ent. 24: 225.

Type-species: *Melia danbyi* Neumoegen, 1892, by monotypy. A junior homonym of *Melia* Bosc, 1813, *in* Risso, *Nouv. Bull. Sci. Soc. philom. Paris* 3: 233, — Crustacea. The objective replacement name is *Eumelia* Neumoegen, 1893.

EUMENOGASTER Herrich-Schäffer, 1855, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.73, fig.423.

Type-species: *Eumenogaster eumenes* Herrich-Schäffer, 1855, *ibidem* 1(1): wrapper, pl.73, fig.423, by monotypy.

Eumenogaster is dated from the wrapper of the part and not the title-page of the volume, 1850-1858.

‡SPHECODES Herrich-Schäffer, [1858], Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 74 (but attributed to Boisduval, an incorrect authorship).

An unavailable name under Article 11(d) of the *Code*, first published in the synonymy of *Eumenogaster* Herrich-Schäffer, 1855, and not subsequently treated as an available name.

EUMETOPONA Fitch, 1856. See Datana Walker, 1855.

EUNADUNA Dognin, 1901, Naturaliste 15: 249. NOTO Type-species: Eunaduna cerurata Dognin, 1901, ibidem 15: 249, by original designation. [ARGENTINA]: Tucuman.

E. cerurata is a junior subjective synonym of Thosea annulipes Berg, 1878, An. Soc. cient. argent. 5: 186.

See also Tecmessa Burmeister, 1878.

EUNOMIA Hübner, 1818, Zuträge Samml. exot. Schmett. 1:

Type-species: Zygaena colombina Fabricius, 1793, Ent. Syst. 3 (1): 403 (but cited as ‡columbina, an incorrect subsequent spelling), by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 146. [WEST INDIES]: Americae merid. insulis.

Incorrect type-species designation: Zygaena andromacha Fabricius, 1775, a nominal species not originally included in Eunomia, and not linked with one of the originally included names when cited by Kirby, in Rye, 1878, Zool. Rec. 13 (Insecta): 165.

Not preoccupied by ‡Eunomia Rafinesque, 1815, Analyse Nat.: 135, - Vermes. An undescribed genus having no originally included species.

‡GLAUCOPIS Hübner, 1808, Erste Zuträge Samml. exot. Schmett.: 3-6.

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1966, *Bull. zool. Nom.* 23 (Opinion 789): 214. Placed on the Official Index of Rejected and Invalid Generic Names in

Zoology: Name No.1844.

The five originally included nominal species were later figured and are all currently placed in Ctenuchidae genera. The first of these species ‡G. psophis (figs 9, 10) was later published by Hübner (1818, Zuträge Samml. exot. Schmett. 1: 8, pl.[2], figs 9, 10) as 'Eunomia Columbina', an incorrect subsequent spelling of Zygaena colombina Fabricius, 1793.

MARISSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 174.

Type-species: Zygaena colombina Fabricius, 1793, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 200.

A junior objective synonym of Eunomia Hübner, 1818.

Marissa was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 200.

EUNOTELA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 263 (key), 325.

Type-species: *Eunotela pallida* Schaus, 1901, *ibidem* 1901: 325, pl.12, fig.5, by original designation. [BRAZIL]: Parana, Castro.

EUNYSTALEA Grote, 1895, Abh. naturw. Ver. Bremen 14: 48.

Type-species: *Nystalea indiana* Grote, 1884, *Papilio* 4: 7, by original designation. [U.S.A.]: Florida, Indian River.

‡EUNYSTATEA; Neave, 1939, Nomencl. zool. 2: 339.

An incorrect subsequent spelling of Eunystalea Grote, 1895.

‡EUNYSTATEA; misspelling. See Eunystalea Grote, 1895.

‡EUPHARPYIA; misspelling. See Euharpyia Schaus, 1901.

EUPHASTIA Dognin, 1911, Hétérocères nouv. Am. Sud 2: 35.

Type-species: *Euphastia ophidera* Dognin, 1911, *ibidem* 2: 35, by original designation. French Guiana: St Laurent du Maroni.

E. ophidera is a junior subjective synonym of Rifargia nubila Druce, 1901, Ann. Mag. nat. Hist. (7) 7: 437.

‡EUPHROSTIS; misspelling. See Euproctis Hübner, [1819].

EUPLAGIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 180.

Type-species: *Phalaena hera* Linnaeus, 1767, *Syst. Nat.* (Edn 12) 1 (2): 834, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 261, but cited as *Phalaena quadripunctaria* Poda, 1761, a nominal species not originally included in *Euplagia*. Kirby, however, placed *quadripunctaria* as the senior synonym of *hera*, a nominal species originally included in *Euplagia*. Under the *Code*, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. ITALY, GERMANY.

P. hera is a junior subjective synonym of Phalaena quadripunctaria Poda, 1761, Insecta Mus. Graecensis: 89.

EUPLESIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, figs 9, 10. ARCT

Type-species: Euplesia vittigera Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.9, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 166, but cited as Glaucopis sphingidea Perty, 1833, a nominal species not originally included in Euplesia. Kirby, however, placed sphingidea, as the senior synonym of vittigera, a nominal species originally

included in *Euplesia*. Under the *Code* Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. [BRAZIL]: [R. Amazon] fl. Amazonas.

E. vittigera is a junior subjective synonym of Glaucopis sphingidea Perty, in Martius 1833, Delectus Anim. articulatorum Brasil: 158, fig.

EUPLEXIDIA Hampson, 1896, Fauna Br. India (Moths) 4: 461.

Type-species: *Euplexidia noctuiformis* Hampson, 1896, *ibidem* 4: 461, fig.246, by original designation. INDIA: [Assam], Khasis.

Hampson attributed both *Euplexidia* and *noctuiformis* to 'Swinh. MS' and placed them in the Notodontidae. They were subsequently placed in the Noctuidae by Hampson, 1908, *Cat. Lepid. Phalaenae Br. Mus.* 7: 110.

EUPLEXIDIA Swinhoe, 1897, Ann. Mag. nat. Hist. (6) 19: 165.

Type-species: Euplexidia noctuiformis Hampson, 1896, by original designation.

A junior homonym of *Euplexidia* Hampson, 1896. The generic name was used and unintentionally made nomenclaturally available by Hampson prior to its proposal and generic description by Swinhoe for the same concept. The objective replacement name is *Euplexidia* Hampson, 1896.

EUPLEXIDIA Swinhoe, 1897. See Euplexidia Hampson, 1896. NOCTUIDAE

EUPLOCIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 172.

Type-species: *Phalaena membliaria* Cramer, 1780, *Uitlandsche Kapellen (Papillons exot.)* 3: 139, 175, pl.269, figs C, D (but cited as *membliare*, an unjustified emendation by Hübner, [1819]), by monotypy. [EAST INDIES] Indes Orientales.

P. membliaria is dated from the wrapper of the part and not

from the title-page of the volume, 1782.

‡EUPREPIA; misspelling. See Arctia Schrank, 1802. ARCT

EUPREPIA Agassiz, 1846. See Arctia Schrank, 1802. ARCT

EUPROCTIDION Holland, 1893, Psyche, Camb. 6: 414.

Type-species: *Euproctidion gabunica* Holland, 1893, *ibidem* 6: 414, 567, pl.10, fig.19, by monotypy. [GABON]: [Ogooué] Ogové R. valley (see page 373).

EUPROCTILLA Aurivillius, 1904, Ark. Zool. 2 (4): 54.

Type-species: *Euproctilla insignis* Aurivillius, 1904, *ibidem* 2 (4): 54, fig.43, by monotypy. [Cameroun] Kamerun.

Euproctilla when established contained a second nominal species 'Euproctilla? disjuncta n. sp.' but this was doubtfully included and, under the Code Article 68(c), is not eligible for fixation as type-species.

EUPROCTILLINA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 141.

Type-species: Artaxa mesomelaena Holland, 1893, Psyche, Camb. 6: 417, 567, pl.10, fig.20, by original designation. [GABON]: [Ogooué] Ogové R. valley (see page 373).

Euproctillina is here adopted as the correct original spelling, under the Code, Article 32(b), from a multiple original spelling that included ‡Euprotillina.

The date of publication of the part of Seitz containing page 141 is cited on page 137 as 1.xi.1926 in both the German and the English editions.

‡EUPROTILLINA; Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 141, 576.

An incorrect (of a multiple) original spelling of *Euproctillina* Hering, 1926.

Although proposed and indexed in the German edition as ‡ Euprotillina gen. nov.' the spelling Euproctillina was used in the heading of the page and the new genus was compared with the preceding genus Euproctilla Aurivillius. In the English edition published on the same day the genus was proposed as Euproctillina but indexed as ‡Euprotillina.

EUPROCTILLOPSIS Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 142.

Type-species: *Euproctillopsis affinis* Hering, 1926, *ibidem* 14: 143, pl.28, row f, by original designation. [CAMEROUN] Kamerun: Lolodorf.

The date of publication of the part of Seitz containing page 142 is cited on page 137 as 1.xi.1926 in both the German and the English editions.

EUPROCTIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 159.

Type-species: *Phalaena chrysorrhoea* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 502, by subsequent designation by Moore, [1883] 1882-3, *Lepid. Ceylon* 2: 89. Type-locality: not stated, [Europe].

‡LEUCOMA Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4)(Opinion 97): 19. Also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140.

Only included species: *Bombyx auriflua* [Denis & Schiffermüller], 1775, (a junior subjective synonym of *Phalaena similis* Fuessly, 1775), which is congeneric with *Phalaena chrysorrhoea* Linnaeus, 1758, the type-species of *Euproctis* Hübner, [1819].

PORTHESIA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 65.

Type-species: *Phalaena chrysorrhoea* Linnaeus, 1758, by subsequent designation by Westwood, 1840, *Introd. mod. Classif. Insects* 2 (Synopsis Genera Br. Insects): 92.

Invalid type-species designation: Phalaena chrysorrhoea Linnaeus, 1758, cited by Boisduval, 1836, Hist. nat. Insectes (Spec. gén. Lépid.) 1: 143. Boisduval in his lengthy 154 page Introduction, reviewed earlier classifications and designated up to three different type-species for each generic name. In his 'Exposé de notre méthode' from page 155-690, no type-species designations were made for the genera he himself used. Under the Code, Article 69(a)(iii), the type-designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species.' Boisduval's type designations although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not in the past been accepted by Hemming, or other authors.

Porthesia was again proposed by Stephens, 1829 [June], Nom. Br. Insects: 40; and 1829 [July], Syst. Cat. Br. Insects (2): 53.

‡EUPHROSTIS; Lucas, 1890, Proc. Linn. Soc. N.S.W. (2) 4: 1090.

An incorrect subsequent spelling of *Euproctis* Hübner, [1819].

EUPROCTOIDES Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 544.

Type-species: *Euproctoides miniata* Bethune-Baker, 1911, *ibidem* (8) 7: 544, by original designation. [ANGOLA]: [Vila Salazar] N'Dalla Tando.

EUPROCTOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxiii, 615.

Type-species: *Euproctosia cretata* Hampson, 1914, *ibidem* 1: 615, fig.185, by original designation. [GHANA] Gold Coast: Bibianaha.

‡EUPROTILLINA; misspelling. See Euproctillina Hering, 1926.

EUPSEUDOSOMA Grote, 1865, Proc. ent. Soc. Philad. 5: 240.

Type-species: Charidea nivea Herrich-Schäffer, [1855 June] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.51, fig.279, by monotypy. BRAZIL.

C. nivea is a junior subjective synonym of Phalaena involuta Sepp, [1855], Surinamsche Vlinders 1: 253.

EUPSYCHOMA Grote, 1865, Proc. ent. Soc. Philad. 4: 317.

Type-species: *Eupsychoma geometrica* Grote, 1865, *ibidem* 4: 318, by monotypy. U.S.A.: Colorado.

E. geometrica is a junior subjective synonym of Phalaena plantaginis Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 501.

See also Parasemia Hübner, [1820].

EUPYDNA Fletcher nom. n. for Pydna Walker, 1856. NOTO Type-species: Pydna testacea Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1754, by monotypy (of Pydna Walker, 1856). [INDIA]: Hindostan.

PYDNA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1753.

Type-species: Pydna testacea Walker, 1856, by monotypy.

A junior homonym of *Pydna* Herrich-Schäffer, 1855, *Syst. Bearbeitung Schmett. Eur.* 6: 127, 128, – Lepid., Geometridae. The objective replacement name is *Eupydna* Fletcher nom. n.

According to Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, and to Hemming, 1937, Hübner 1: 584, 589, Heft 68 (containing pages 81-152 of volume 6 of Herrich-Schäffer's work) was published in 1855.

EUPYRA Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.13, figs 54-57.

Type-species: *Eupyra imperialis* Herrich-Schäffer, [1853] 1850-1858, *ibidem* 1 (1): wrapper, pl.13, fig.54 (but attributed to Boisduval, an incorrect authorship), by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 124. Venezuela.

EURATA Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.13, fig.52.

Type-species: *Eurata picta* Herrich-Schäffer, [1853] 1850-1858, *ibidem* 1 (1): wrapper, pl.13, fig.52, by monotypy (but attributed to Boisduval, an incorrect authorship). VENEZUELA.

The name Eurata probably originated from the same Boisduval manuscript name from which Eurota Walker, 1854, was derived.

See also Eurota Walker, 1854.

EURODES Turner, 1899, Trans. Proc. R. Soc. S. Aust. 23:

Type-species: Eurodes micrommata Turner, 1899, ibidem 23: 12, by monotypy. AUSTRALIA: Northern Territory, Port Darwin.

EUROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 88, 330.

Type-species: Setinochroa trimacula Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 14, 85, pl.157, figs 27, 28, by original designation. [SRI LANKA] Ceylon: Nawala-pittia.

EUROTA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 180.

Type-species: Glaucopis pictula Walker, 1854, ibidem 1: 181,

by monotypy. VENEZUELA.

G. pictula is a junior subjective synonym of Eurata picta Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.13, fig.52.

Eurota was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 285.

See also Eurata Herrich-Schäffer, [1853].

EUROZONOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxvii, 788.

Type-species: Eurozonosia fulvinigra Hampson, 1914, ibidem 1: 789, fig. 258, by original designation. [Kenya] Br. E. Afr.: N. Kavirondo, Mumias.

EURYLOMIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.26.

Type-species: Eurylomia curvinervis Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.26, by monotypy. MEXICO: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 5.

E. curvinervis is a junior subjective synonym of Lithosia cordula Boisduval, 1870, Considérations Lépid. Guatemala: 95.

EURYNORA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxvii, 788.

Type-species: *Bitecta flaveola* Rothschild, 1912, *Novit. zool.* 19: 223, by original designation. New Guinea: [Papua New Guinea], Mambare R., Biagi.

EURYPEPLA Turner, 1922. See Eurypeplella Hedicke, 1923.

EURYPEPLELLA Hedicke, 1923, Dt. ent. Z. 1923: 226. LITH Type-species: Eurypepla pteridaula Turner, 1922, Proc. R. Soc. Vict. 35: 31, by monotypy (of Eurypepla Turner, 1922). AUSTRALIA: MacIntyre Range, Queensland National Park.

Eurypeplella was established as the objective replacement name for Eurypepla Turner, 1922.

name for Eurypepia Turner, 1922.

EURYPEPLA Turner, 1922, Proc. R. Soc. Vict. 35: 30.

Type-species: Eurypepla pteridaula Turner, 1922, by monotypy.

A junior homonym of *Eurypepla* Boheman, 1854, *Monogr. Cassididarum* 2: 236, – Insecta, Coleoptera. The objective replacement name is *Eurypeplella* Hedicke, 1923.

EURYPTIDIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 91, 230.

Type-species: Euryptidia univitta Hampson, 1900, ibidem 2: 230, fig. 152, by original designation. BRAZIL: Rio [de] Janeiro.

EURYSTAURA Janse, 1920, Ann. Transv. Mus. 7: 216. NOTO Type-species: Eurystaura brunnea Janse, 1920, ibidem 7: 216, pl.9, figs 5-9, pl.14, fig.11, by original designation. SOUTH AFRICA: [Natal], Durban.

EURYSTAURELLA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 42.

Type species: *Eurystaurella subaequa* Kiriakoff, 1962, *ibidem* (8vo Sci. zool.) **114**: 42, fig.54, by original designation. [ZAIRE]: Sankuru, Katako-Kombe.

EURYSTAURIDIA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 24.

Type-species: Eurystaura olivacea Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 429, pl.71 row c, by original designation. Type-localities: cited as 'Kamerun, Ruwenzori,

Lindi'. The 'Type 6' in BMNH is from [ZAIRE] west Semliki valley, 20 miles S.W. of Boga.

In the text of the English edition of Seitz, the reference to the figure of *E. olivacea* is incorrectly cited as pl.71 row n.

EURYSTAUROPSIS Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 430.

Type-species: *Eurystauropsis albidilinea* Gaede, 1928, *ibidem* 14: 430, pl.71 row d, by monotypy. NIGERIA: southern.

The date of publication of the part of Seitz containing page 430 is cited on page 425 as 30.iii.1928 in both the German and the English editions.

EUSCHAUSIA Dyar, 1898, Can. Ent. 30: 81.

Type-species: Halysidota argentata Packard, 1864, Proc. ent. Soc. Philad. 3: 129 (but cited as ‡Halesidota, an incorrect subsequent spelling), by subsequent designation (for Schausia Dyar, 1897) by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 145 (but cited as ‡Halisidota, an incorrect subsequent spelling). U.S.A.: Cal[ifornia], Gulf of Georgia.

Euschausia was established as the objective replacement

name for Schausia Dyar, 1897.

SCHAUSIA Dyar, 1897, Can. Ent. 29: 212, 217.

Type-species: *Halysidota argentata* Packard, 1864, by subsequent designation by Hampson, 1901, *Cat. Lepid. Phalaenae Br. Mus.* 3: 145.

A junior homonym of *Schausia* Karsch, 1895, *Ent. Nachr.* 21: 346, – Lepid., Agaristidae. The objective replacement name is *Euschausia* Dyar, 1898.

EUSCOTUROPSIS Bryk, 1930, in Strand, Lepid. Cat. 42: 17.

Type-species: Scoturopsis elongata Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 508, by original designation. PERU: Vilcanota.

EUSHACHIA Matsumura, 1925, Zool. Mag. Tokyo 37: 404.

Type-species: Eushachia auripennis Matsumura, 1925, ibidem 37: 404, pl.7, fig.2, by original designation, but cited as ‡auripenuis, an incorrect (of a multiple) original spelling. TAIWAN.

E. auripennis is a subspecies of Niganda aurata Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 64.

See also Midasia Kiriakoff, 1962.

EUSTEMA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 260 (key), 285.

Type-species: Stilpnotia dara Druce, 1894, Ann. Mag. nat. Hist. (6) 13: 355, by original designation. MEXICO: Orizaba.

EUSTEMIDES Dyar, 1908, Proc. ent. Soc. Wash. 9: 46.

Type-species: Eustema carama Druce, 1904, Ann. Mag. nat. Hist. (7) 13: 248, by original designation. Peru: south-eastern, Santo Domingo.

EUTAENIA Wallengren, 1875. See Paralacydes Aurivillius, 1899.

EUTANE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 531.

Type-species: *Eutane terminalis* Walker, 1854, *ibidem* 2: 531, by monotypy. [AUSTRALIA] New Holland: specimens from unstated localities and from New South Wales.

EUTELESIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 91, 398.

Type-species: Nola vulgaris Druce, 1885, Biologia cent.-am.

(Zool.) Lepid.-Heterocera 1: 141, pl.13, fig.22, by original designation. Costa Rica: Irazu, Rio Sucio.

EUTHEMONIA Stephens, 1828. See Diacrisia Hübner, [1819]. ARCT

EUTHORACIA Hulstaert, 1923, Revue zool. afr. 11: 411.

Type-species: Caryatis syntomina Butler, 1878, Ann. Mag. nat. Hist. (5) 2: 456, by original designation. NIGERIA: Old Calabar.

EUTHYONE Watson nom. n. for Thyone Walker, 1854. LITH Type-species: Thyone simplex Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 553, by monotypy (of Thyone Walker, 1854). Honduras.

THYONE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 552.

Type-species: Thyone simplex Walker, 1854, by monotypy. A junior homonym of Thyone Philippi, 1840, Arch. Naturgesch. 6 (1): 190, - Crustacea. The objective replacement name is Euthyone Watson nom. n.

EUTIMIA Wallengren, 1858, Ofvers. K. VetenskAkad. Förh. Stockh. 15: 211.

Type-species: Eutimia marpissa Wallengren, 1858, ibidem 15: 211, by original designation. [South Africa]: Caffraria.

E. marpissa was again described as a new species by Wallengren, 1860, Wien, ent. Monatschr. 4: 164.

See also Parauchmis Holland, 1897.

‡EUTOMIA; misspelling. See Eutomis Hübner, [1819]. CTEN

EUTOMIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 122.

Type-species: Sphinx minceus Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 109, pl.347, fig.A, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 100. [South Africa]: [Cape of Good Hope] Kaap de Goede Hoop.

S. minceus is dated from the wrapper of the part and not from the title-page of the volume, 1782.

Eutomis was attributed incorrectly, to himself by Boisduval, 1870, Considérations Lepid. Guatemala: 80.

‡EUTOMIA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling of *Eutomis* Hübner, [1819].

EUTORNOPERA Hampson, 1895, Trans. ent. Soc. Lond. 1895:

Type-species: Eutornopera argentifascia Hampson, 1895, ibidem 1895: 280, fig., by original designation. BHUTAN.

EUTRICHOLOBA Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 334.

Type-species: Eutricholoba signata Kiriakoff, 1965, ibidem 101: 335, fig.3, by original designation. CAMEROUN: Lomié.

EUTROTONOTUS Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 440.

Type-species: Eutrotonotus albidilinea Gaede, 1928, ibidem 14: 440, pl.72 row g, by monotypy. MADAGASCAR.

The date of publication of the part of Seitz containing page 440 is cited on page 433 as 30.v.1928 in the German edition, and as 26.xi.1928 in the English edition.

EUVERNA Neumoegen & Dyar, 1893, Ent. News 4: 141.

Type-species: Seirarctia clio Packard, 1864, Proc. ent. Soc. Philad. 3: 120, by original designation. U.S.A.: California.

EUXOGA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 692. NOTO

Type-species: Euxoga argenteopunctata Möschler, 1878, ibidem 27 (Abh.): 692, pl.10, fig.50, by monotypy. SURINAM.

EUZEUGAPTERYX Dyar, 1905, Proc. U.S. natn. Mus. 29: 175.

Type-species: Euzeugapteryx speciosa Schaus, 1905, ibidem 29: 205, by original designation (in anticipation of Schaus's description of speciosa). FRENCH GUIANA: Maroni River, St Jean.

Euzeugapteryx was nomenclaturally available but was without an available included species between its date of publication 1905 October 3rd and that of its type-species 1905 October 11th.

EUZORA Turner, 1915, Proc. R. Soc. Qd 27: 22. LYMA Type-species: Porthesia collucens Lucas, 1890, Proc. Linn. Soc. N.S.W. (2) 4: 1090, by original designation. AUSTRALIA: Queensland, Brisbane.

P. collucens was established in the subgenus ‡Euphrostis, an incorrect subsequent spelling of Euproctis Hübner, [1819].

EVAGRA Boisduval, 1870. See Agyrta Hübner, [1820].

EVIUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 639.

Type-species: Phalaena hippia Stoll, 1790, in Cramer, Uitlandsche Kapellen (Papillons exot.) (Aanhangsel): 57, 183, pl.12, fig.8, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 197. SURINAM.

P. hippia is dated from the wrapper of the part and not from

the title-page of the volume, 1791.

EXAERETA Hübner, [1820]. See Dicranura R.L., 1817. NOTO

EXANTONGILA Fletcher nom. n. for Antongila Kiriakoff,

Type-species: Desmeocraera antongilensis Viette, 1955, Bull. mens. Soc. linn. Lyon 24: 221, figs 1-3, by original designation (for Antongila Kiriakoff, 1958). MADAGASCAR: north-eastern, Maroantsetra.

Prof. Dr S.G. Kiriakoff was informed of the homonymy and has agreed to the proposal of this replacement name.

ANTONGILA Kiriakoff, 1958, Mém. Inst. scient. Madagascar (E) 9: 224.

Type-species: Desmeocraera antongilensis Viette, 1955, by

original designation.

A junior homonym of Antongila Roewer, 1931, Z. wiss. Zool. 138: 170, - Arachnida. The objective replacement name is Exantongila Fletcher nom. n.

See Dicranura R.L., 1817. #EXCERATA: misspelling.

EXEMASIA Dyar, 1920, Insecutor Inscit. menstr. 8: 31.

Type-species: Exemasia ochropasa Dyar, 1920, ibidem 8: 31, by original designation. MEXICO: Mexico City.

‡EXERETA; misspelling. See Dicranura R.L., 1817. NOTO

EXILISIA Toulgoët, 1958, Mém. Inst. scient. Madagascar (E) **9**: 181.

Type-species: Philenora punctata Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 507, 508, pl.33, fig.22, by original designation. MADAGASCAR.

EXITELICA Turner, 1921, Proc. Linn. Soc. N.S.W. 46: 160. ARCT

Type-species: Leptosoma aegrotum Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 145, pl.5, fig.15, by monotypy. AUSTRALIA: New South Wales.

EXOTROCHA Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 691, 693.

Type-species: Phalaena liboria Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 106, pl.345, fig.D, by

monotypy. SIERRA LEONE.

P. liboria is dated from the wrapper of the part and not from the title-page of the volume, 1782. P. liboria is a junior subjective synonym of Phalaena fulvia Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 509.

See also Clerckia Aurivillius, 1882.

EYPREPIA Ochsenheimer, 1810. See Arctia Schrank, 1802. ARCT

EYRALPENUS Butler, 1875, Cistula ent. 2: 35. Type-species: Spilosoma testacea Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 670, by original designation. South Africa: [Durban] Port Natal, and [Zululand] Zoolu Country.

FABRESEMA Holloway, 1979, Series Ent. 15: 367. Type-species: Fabresema gerardi Holloway, 1979, ibidem 15: 369, text-figs 77, 79, pl.70, fig.4, by original designation. NEW CALEDONIA: Les Dalmates.

FANALA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 227.

Type-species: Dasychira abbreviata Kenrick, 1914, Trans. ent. Soc. Lond. 1913: 596, pl.32, fig.36, by original designation. MADAGASCAR: Central.

FANAMBANA Kiriakoff & Viette, 1969, Lambillionea 68: 3.

Type-species: Fanambana pyralidina Kiriakoff & Viette, 1969, ibidem 68: 6, figs 1, 2, by original designation. MADA-GASCAR: northern, 37km south of Vohémar.

FARIGIA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 260 (key), 283.

Type-species: Heterocampa sagana Druce, 1894, Ann. Mag. nat. Hist. (6) 13: 357, by original designation. MEXICO: Teapa in Tabasco.

FASSLIA Dognin, 1911, Hétérocères nouv. Am. Sud 2: 15.

Type-species: Fasslia hampsoni Dognin, 1911, ibidem 2: 15, by monotypy. Colombia: near Cali, Monte Socorro.

FELDERIA Kirby, 1892. See Acreagris Felder, 1874. **ZYGAENIDAE**

FELIA Walker, 1864, J. Proc. Linn. Soc. (Zool.) 7: 197.

Type-species: Felia intermixta Walker, 1864, ibidem (Zool.) 7: 197, by monotypy. Borneo: Sarawak.

FENTONIA Butler, 1881, Trans. ent. Soc. Lond. 1881: 20.

Type-species: Fentonia laevis Butler, 1881, ibidem 1881: 20, by original designation. JAPAN: Yokohama.

F. laevis is a junior subjective synonym of Harpyia ocypete Bremer, 1861, Bull. Acad. imp. Sci. St. Pétersb. 3: 481.

See also Urocampa Staudinger, 1892, and Neoshachia Matsumura, 1925.

FENTONINA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 421. NOTO

Type-species: Fentonina punctum Gaede, 1928, ibidem 14: 421, pl.70 row d, by monotypy. [TANZANIA]: Lindi.

The date of publication of the part of Seitz containing page 421 is cited as 29.iii.1928 on page 417 of both the German and the English editions.

FLETCHERINIA Griveaud, 1964, Faune Madagascar 17: 126.

Type-species: Fletcherinia decaryi Griveaud, 1964, ibidem 17: 127, pl.1, fig.87, by original designation. MADAGASCAR: Central, région de l'Ankaizina.

FOCILLISTIS Hampson, 1926, Descr. new Genera Species Lepid. Phalaenae Subfamily Noctuinae Br. Mus.: 599. NOCTUIDAE

Type-species: Zethes salsoma Swinhoe, 1902, Ann. Mag. nat. Hist. (7) 10: 499, by original designation.

Further information in Nye, 1975, Generic Names Moths World 1: 209.

‡FOCILLSTIS; Nye, 1975, Generic Names Moths World 1:

An incorrect subsequent spelling of Focillistis Hampson, 1926.

‡FOCILLSTIS; misspelling. See Focillistis Hampson, 1926.

FODINOIDEA Saalmüller, 1884, Lepid. Madagascar (1): 154.

Type-species: Fodinoidea staudingeri Saalmüller, 1884, ibidem (1): 154, pl.5, fig.63, by monotypy. MADAGASCAR: S., 'Betsileo-land'.

Fodinoidea was established in the Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 82; it was placed in the Hypsidae by Toulgoët, 1957, Annls Soc. ent. Fr. 125: 73; it is retained in the Arctiidae, and placed in the Arctiinae.

†FORMAFENTONIA: misspelling. See Formofentonia Matsumura, 1925. NOTO

FORMOFENTONIA Matsumura, 1925, Zool. Mag. Tokyo 37: 396.

Type-species: Formofentonia rotundata Matsumura, 1925, ibidem 37: 396, pl.6, fig.10, by monotypy. TAIWAN.

From a multiple original spelling in which both Formofentonia and ‡Formafentonia were used, we, as first revisers, adopt Formofentonia, the spelling now in current use.

F. rotundata is a subspecies of Stauropus orbifer Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 152.

‡FORMAFENTONIA; Matsumura, 1925, Zool. Mag. Tokyo 37: 396.

An incorrect (of a multiple) original spelling of Formofentonia Matsumura, 1925.

FORMOTENSHA Matsumura, 1925, Zool. Mag. Tokyo 37: 397.

Type-species: Formotensha marginalis Matsumura, 1925, ibidem 37: 397, by original designation. TAIWAN.

FREGELLA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 272.

Type-species: Euchromia semiluna Walker, 1854, ibidem 1: 272, by monotypy. Type-locality: not stated, [America]: [tropical].

Fregella was established as a subgenus of Euchromia Hübner, [1819].

FURCULA Lamarck, 1816, Hist. nat. Anim. sans Vertèbres 3:

Type-species: Phalaena furcula Clerck, 1759, Icon. Insect.

rariorum 1: pl.9, fig.9 (but included by Lamarck as Furcula salicis, an unnecessary objective replacement name; see below), by absolute tautonymy. TYPE-LOCALITY: not stated, [EUROPE].

Furcula salicis Lamarck, 1816, ibidem 3: 581, was established unnecessarily as an objective replacement name for Phalaena furcula Clerck, but attributed incorrectly to Fabricius.

HARPYIAS Hübner, [1819] 1816, Verz. bekannter Schmett.: 148.

Type-species: *Phalaena furcula* Clerck, 1759, by subsequent designation by Lempke, 1951, *Ent. Ber., Amst.* 13: 295.

A junior objective synonym of Furcula Lamarck, 1816.

FUSADONTA Matsúmura, 1920, Zool. Mag. Tokyo 32: 146.

Type-species: *Notodonta basilinea* Wileman, 1911, *Trans. ent. Soc. Lond.* **1911**: 292, pl.30, fig.23, by monotypy. Japan: [Honshu], Yamato prov., Yoshino.

Fusadonta was again proposed by Matsumura, 1921, Thou-

sand Insects Japan (Additam.) 4: 795.

‡FUSADONTA; misspelling. See Takadonta Matsumura, 1920.

FUSAPTERYX Matsumura, 1920, Zool. Mag. Tokyo 32: 148.

Type-species: Lophopteryx ladislai Oberthür, 1879, Diagnoses Espèces nouv. Lépid. ile Askold: 13, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

Fusapteryx was again proposed by Matsumura, 1921, Thou-

sand Insects Japan (Additam.) 4: 814.

GADIANA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 33: 854.

Type-species: Gadiana rufescens Walker, 1865, ibidem 33:

854, by monotypy. [COLOMBIA]: Bogota.

Gadiana was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 333.

GAEDEA Romieux, 1943. See Gaedeina Romieux, 1945.

GAEDEINA Romieux, 1945, Mitt. schweiz. ent. Ges. 19: 274.

Type-species: *Gaedea alternata* Romieux, 1943, *ibidem* 19: 104, pl.9, fig.7, by original designation (for *Gaedea* Romieux, 1943). [ZAIRE] Congo Belge: Haut-Katanga, Shinkolobwe.

Gaedeina was established as the objective replacement name for Gaedea Romieux, 1943.

GAEDEA Romieux, 1943, Mitt. schweiz. ent. Ges. 19: 103.

Type-species: Gaedea alternata Romieux, 1943, by original designation.

A junior homonym of *Gaedea* Hering, 1924, *Dt. ent. Z.* **1924**: 273, – Lepid., Zygaenidae. The objective replacement name is *Gaedeina* Romieux, 1945.

GALANTHELLA Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 183.

Type-species: Norraca arctipennis Holland, 1893, Psyche, Camb. 6: 488, by original designation. [EQUATORIAL GUINEA]: [Rio Muni, Rio Benito] Benita. The type-locality was stated to be Benita, 'a mission station upon the coast 110 miles north of the Equator'.

GALETHALEA Butler, 1876, J. Linn. Soc. (Zool.) 12: 424.

CTEN

Type-species: Halysidota pica Walker, 1855, List Specimens

lepid. Insects Colln Br. Mus. 3: 743, by original designation (but cited as ‡Halesidota, an incorrect subsequent spelling by Walker, 1855). [Brazil]: Rio [de] Janeiro.

GALLABA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 457.

Type-species: Gallaba duplicata Walker, 1865, ibidem 32: 458, by monotypy. [Australia]: [Queensland], Moreton Bay.

GALLIENICA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 227.

Type-species: *Parorgyia maligna* Butler, 1882, *Cistula ent*. 3: 17, by original designation (but cited by Griveaud in ‡*Panorgyia*, an incorrect subsequent spelling). MADAGASCAR: Ankafana.

GALONA Karsch, 1895, Ent. Nachr. Berlin 21: 363. NOTO Type-species: Galona serena Karsch, 1895, ibidem 21: 363, pl.3, fig.1, by subsequent designation by Janse, 1920, Ann. Transv. Mus. 7: 205. ANGOLA.

GALOS Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 172.

Type-species: *Galos franki* Kiriakoff, 1959, *ibidem* **95**: 172, fig.18, by original designation. [ZAIRE] Congo Belge: Sankuru, Lusambo.

GALTARA Walker, [1863] 1862, List Specimens lepid. Insects Colln Br. Mus. 26: 1688.

Type-species: *Galtara purata* Walker, [1863] 1862, *ibidem* **26**: 1688, by monotypy. South Africa.

Galtara was established in the Geometridae; it was placed in the Arctiidae by Hampson, 1900, Ann. S. Afr. Mus. 2: 62.

BITHRA Fawcett, 1918, Proc. zool. Soc. Lond. 1917: 235.

Type-species: Bithra aganice Fawcett, 1918, ibidem 1917: 235, pl.1, fig.11, by monotypy. [KENYA] Br. E. Afr.: [Teita Hills], Kedai.

A junior homonym of Bithra Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 621, — Lepid., Notodontidae. There is no objective replacement name but the typespecies of Bithra Fawcett is currently considered to be congeneric with the type-species of Galtara Walker. The latter is therefore available for use as a subjective replacement name.

GAMPOLA Moore, 1878, Proc. zool. Soc. Lond. 1878: 26.

Type-species: *Gampola fasciata* Moore, 1878, *ibidem* **1878**: 27, pl.2, fig.18, by monotypy. [Sri Lanka] Ceylon.

‡GAMPONA; Lucas, 1901, Arch. Naturgesch. 64 (2): 624.
An incorrect subsequent spelling of Gampola Moore, 1878.

‡GAMPONA; misspelling. See Gampola Moore, 1878.

GANDHARA Moore, 1878, Proc. zool. Soc. Lond. 1878: 15.

Type-species: Lithosia serva Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 506, by monotypy. NEPAL.

GANGAMELA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 140.

Type-species: Gangamela saturata Walker, [1865] 1864, ibidem 31: 140, by monotypy. BRAZIL.

UXELA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1982.

Proposed, unnecessarily, as an objective replacement name for *Gangamela* Walker, [1865] which is not preoccupied by ‡*Gangemela* Walker, 1865, *ibidem* 32: 443, an incorrect subsequent spelling of *Gaugemela* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 138, – Lepid., Notodontidae.

‡UZELA; Neave, 1940, Nomencl. zool. 4: 624.
An incorrect subsequent spelling of Uxela Walker, 1866.

‡GANGAMELA; misspelling. See Gaugamela Walker, 1862.

GANGARIDES Moore, 1866, Proc. zool. Soc. Lond. 1865: 821.
NOTO

Type-species: Gangarides dharma Moore, 1866, ibidem 1865: 821, pl.43, fig.7, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 810. INDIA: N.E. Bengal.

Gangarides was established in the Lasiocampidae; it was placed in the Eupterotidae by Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 42; it was transferred to the Notodontidae by Grunberg, 1912, in Seitz, Gross-Schmett. Erde 2: 311.

GANGARIDOPSIS Grünberg, 1912, in Seitz, Gross-Schmett. Erde 2: 294.

Type-species: Gangarides citrina Wileman, 1911, Trans. ent. Soc. Lond. 1911: 283, pl.31, fig.3, by monotypy. JAPAN: Nikko.

The date of publication of the part of Seitz containing page 294 is cited on page 289 as 20.iii.1912 in both the German and English editions.

GARDINIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 286.

Type-species: Chrysocale magnifica Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 96, by original designation. [COLOMBIA]: Bogota.

GARDINICA Strand, 1932, Folia zool. hydrobiol. 4: 146.

Proposed, unnecessarily, as an objective replacement name for *Gardinia* Kirby, 1892, which is not preoccupied by ‡*Gardinia* Desmarest, in Chenu, 1877, *Encycl. Hist. nat. Crust. Moll. Zoophytes*: 164, an incorrect subsequent spelling of *Gadinia* Gray, 1824, *Phil. Mag.* 63: 275, – Mollusca.

GARDINICA Strand, 1932. See Gardinia Kirby, 1892. PERI

GARGETTA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 455.

Type-species: Gargetta costigera Walker, 1865, ibidem 32: 455, by monotypy. [INDIA]: Darjeeling.

GARGETTIANA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 617.

Type-species: Gargetta melanosticta Joicey & Talbot, 1917, Ann. Mag. nat. Hist. (8) 20: 61, pl.2, fig.5, by original designation. New Guinea: [Irian Jaya], Wandammen Mts.

The date of publication of the part of Seitz containing 617 is cited on that page as 10.iii.1930 in both the German and the English editions.

GARGETTINA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 28.

Type-species: *Psalisodes longivitta* Kiriakoff, 1962, *Annls Mus. r. Afr. cent.* (8vo Sci. zool.) **114**: 45, fig.58, by original designation. [ZAIRE]: Sankuru, Katako-Kombe.

GARGETTOSCRANCIA Strand, 1912, Arch. Naturgesch. 78 (A) 6: 162.

Type-species: Scrancia albolineata Strand, 1912, ibidem 78 (A) 6: 162, by monotypy. [EQUATORIAL GUINEA]: [Rio Muni], Nkolentangan.

Gargettoscrancia Was established as a subgenus of Scrancia Holland, 1893.

GARTHA Schaus, 1892, Am. Lepid. Illust. new & rare Species:

Type-species: Gartha dalsa Schaus, 1892, ibidem: 4, pl.1, fig.15, by monotypy. Brazil: Sta Catarina, Ecuador.

G. dalsa is a junior subjective synonym of Lagaria acuminata

Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1608.

GARUDINIA Moore, [1882] 1882-3, *Lepid. Ceylon* **2**: 59.

Type-species: *Tospitis latana* Walker, 1863, *List Specimens lepid. Insects Colln Br. Mus.* **28**: 427, by monotypy. [SRI LANKA] Ceylon.

GARUDINISTIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 82, 280.

Type-species: *Tospitis eburneana* Walker, 1863, *List Specimens lepid. Insects Colln Br. Mus.* 28: 428, by original designation. Borneo: Sarawak.

GARUDINODES Bethune-Baker, 1908, Novit. zool. 15: 195.

Type-species: *Garudinodes bicolorana* Bethune-Baker, 1908, *ibidem* 15: 195, by original designation. New Guinea: [Papua New Guinea], Ekeikei.

GASTROCHRYSIA Kiriakoff, 1954, Revue Zool. Bot. afr. 50: 186.

Type-species: *Gastrochrysia lineolata* Kiriakoff, 1954, *ibidem* 50: 187, pl.2, fig.23, by original designation. [ZAIRE]: Katanga, [Lubumbashi] Elizabethville.

GAUDEATOR Dyar, 1914, Proc. U.S. natn. Mus. 47: 166.

Type-species: Gaudeator paidicus Dyar, 1914, ibidem 47: 166, by original designation. PANAMA: Cabima.

GAUGAMELA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 138.

Type-species: Gaugamela atrifrons Walker, 1862, ibidem (Zool.) 6: 138, by monotypy. Borneo: Sarawak.

G. atrifrons is a junior subjective synonym of Ichthyura dorsalis Walker, 1862, ibidem 6: 134.

‡GANGAMELA; Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 443.

An incorrect subsequent spelling of *Gaugamela* Walker, 1862.

GAZALINA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 398.

Type-species: *Gazalina venosata* Walker, 1865, *ibidem* 32: 398, by monotypy. [INDIA]: Hindostan.

G. venosata is a junior subjective synonym of Dasychira apsara Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 341.

Gazalina was established in the Liparidae, now Lymantridae; it was transferred to the Notodontidae by Grünberg, 1912, in Seitz, Gross-Schmett. Erde 2: 317; and transferred to the Thaumetopoeidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

GELASTOCERA Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 476.

Type-species: Gelastocera exusta Butler, 1877, ibidem (4) 20: 476, by original designation. JAPAN: Hakodaté.

Gelastocera was established in the Notodontidae; it was placed in the Noctuidae by Hampson, 1894, Fauna Br. India (Moths) 2: 427.

GEODENA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1691. GEOMETRIDAE

Type-species: Geodena quadrigutta Walker, 1856, ibidem 7: 1691, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 429. Type-locality: not known. Specimens associated with the holotype & in BMNH are from SIERRA LEONE, GHANA and NIGERIA.

Geodena was established in the Lithosiidae, now Arctiidae Lithosiinae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, *ibidem* 1: 429; it was transferred to the Aganaidae, now Arctiidae Aganainae, by Swinhoe, 1903, *Trans. ent. Soc. Lond.* 1903: 497; and was placed in the Geometridae by Bethune-Baker, 1911, *Ann. Mag. nat. Hist.* (8) 7: 560.

GERARCTIA Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 448. NOCTUIDAE

Type-species: *Gerarctia poliotis* Hampson, 1905, *ibidem* (7) **15**: 449, by original designation. CANARY ISLANDS: Tenerife.

Gerarctia was established in the Arctiidae; it was transferred to the Noctuidae by Draudt, 1935, in Seitz, Gross-Schmett. Erde 3 (Suppl.): 208.

GERBA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 270.

Type-species: Gerba quadrifasciata Walker, [1865] 1864, ibidem 31: 270, by monotypy. Mexico: [Oaxaca] Oajaca.

GERIDIXIS Dyar, 1914, Proc. U.S. natn. Mus. 47: 170.

Type-species: Geridixis minx Dyar, 1914, ibidem 47: 171, by original designation. PANAMA: Trinidad River.

GERIOJENNSA Watson nom. n. for Jorgensenia Schaus, 1924.

Type-species: Jorgensenia cunegunda Schaus, 1924, Proc. U.S. natn. Mus. 65 (7): 26, by original designation (for Jorgensenia Schaus, 1924). ARGENTINA: Esquina Grande.

JORGENSENIA Schaus, 1924, Proc. U.S. natn. Mus. 65 (7): 26.

Type-species: Jorgensenia cunegunda Schaus, 1924, by original designation.

A junior homonym of *Jorgensenia* Kieffer, 1913, *in* Wytsman, *Genera Insect.* **152**: 118, 148, – Insecta, Diptera. The objective replacement name is *Geriojennsa* Watson nom. n.

GETTA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 139.

Type-species: Getta niveifascia Walker, [1865] 1864, ibidem 31: 140, by monotypy. [Brazil]: Amazon region, Nanta.

G. niveifascia is a junior subjective synonym of Josia clite Walker, 1854, ibidem 2: 312.

GHORIA Moore, 1878, Proc. zool. Soc. Lond. 1878: 12. LITH Type-species: Ghoria albocinerea Moore, 1878, ibidem 1878: 13, pl.1, fig.10, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 316. INDIA: Darjeeling.

‡CHORIA; Schaus, 1894, Proc. zool. Soc. Lond. 1894: 240. An incorrect subsequent spelling of Ghoria Moore, 1878.

GINALDIA Schaus, 1905, Proc. U.S. natn. Mus. 29: 282.

Type-species: Ginaldia davidsoni Schaus, 1905, ibidem 29: 282, by monotypy. French Guiana: Maroni River, St Laurent.

GINSHACHIA Matsumura, 1929, Insecta matsum. 4: 44, 45.

Type-species: *Ginshachia elongata* Matsumura, 1929, *ibidem* 4: 44, pl.1, fig.14, by original designation. TAIWAN: Horisha.

GISARA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 261 (key), 310.

Type-species: Symmerista procne Schaus, 1892, Proc. zool. Soc. Lond. 1892: 336, by original designation. [BRAZIL]: Rio de Janeiro, Corcovado.

GISSARUS Nye nom. n. for Ocneropsis Kozhanchikov, 1950.

LYMA

Type-species: *Ocneropsis relictus* Kozhanchikov, 1950, Fauna SSSR (N.S.) **42** (Insecta) Lepid. 12: 377, fig.183, by original designation (for *Ocneropsis* Kozhanchikov, 1950). U.S.S.R.: [Tadzhikistan], Gissar Mts, R. Varzob valley, Kondara caves.

OCNEROPSIS Kozhanchikov, 1950, Fauna SSSR (N.S.) 42 (Insecta) Lepid. 12: 376.

Type-species: Ocneropsis relictus Kozhanchikov, 1950, by

original designation.

A junior homonym of *Ocneropsis* Uvarov, 1942, *Trans. Am. ent. Soc.* 67: 348, – Orthoptera. The objective replacement name is *Gissarus* Nye nom. n.

‡GLAUCOPIS Hübner, [1806]. See Syntomis Ochsenheimer, 1808.

GLAUCOPIS Fabricius, 1807. See Marmax Rafinesque, 1815. ZYGAENIDAE

‡GLAUCOPIS Hübner, 1808. See Eunomia Hübner, 1818. CTEN

GLAUCOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 91, 229.

Type-species: Glaucosia agyllia Hampson, 1900, ibidem 2: 229, fig.151, by original designation. BRAZIL: [Espirito Santo] Espiritu Santo.

GLAUCOSTOLA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 3, 87.

Type-species: Leucopsumis guttipalpis Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1649, by original designation. BRAZIL.

GLISSA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 186.

Type-species: Glissa bifacies Walker, [1865] 1864, ibidem 31: 186, by monotypy. [Brazil]: [Tefé] Ega.

G. bifacies is a junior subjective synonym of Josia prolifera Walker, 1854, ibidem 2: 302.

Glissa was established in the Chrysaugidae, now Pyralidae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 405.

See also Adelphoneura Felder, 1874.

GLUPHISIA Boisduval, 1828, Eur. Lepid. Index meth.: 56.

Type-species: *Bombyx crenata* Esper, 1785, *Die Schmett.* 3 (23): 245, pl.47, figs 3, 4, by monotypy. GERMANY: Frankfurtam-Main.

GLYPHIDIA Agassiz, 1847, Nomencl. zool. (Index univl.): 163, 164.

An unjustified emendation of Gluphisia Boisduval, 1828.

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Glyphidia* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

GLYPHISIA Spuler, 1903, in Hofmann, Schmett. Eur. 1: 94. An unjustified emendation of Gluphisia Boisduval, 1828, and a junior homonym of Glyphisia Stephens, 1829, Nom. Br. Insects: 48, – Lepid., Tortricidae.

GLUPHISIODES Kiriakoff, 1954, Revue Zool. Bot. afr. 49: 294.

Type-species: Gluphisiodes bivittata Kiriakoff, 1954, ibidem 49: 294, pl.1, fig.1, pl.2, fig.12, by original designation. [ZAIRE] Congo Belge: Katanga, [Lubumbashi] Elisabethville.

GLYPHIDIA Agassiz, 1847. See Gluphisia Boisduval, 1828.

GLYPHISIA Spuler, 1903. See Gluphisia Boisduval, 1828. NOTO

GNAMPTONYCHIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 86, 251.

Type-species: Atolmis flavicollis Druce, 1885, Biologia cent.am. (Zool.) Lepid.-Heterocera 1: 132, pl.13, fig.6, by original designation. MEXICO: Durango, Milpas.

GNATHOLOPHIA Felder, 1874, in Felder & Rogenhofer. Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.8.

Type-species: Gnatholophia longinervis Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.8, by monotypy. French Guiana, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 5.

G. longinervis is a form of Devara frigida Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 185.

See also Euchontha Walker, 1865, and Macroneurodes Warren, 1900.

‡GRATHOLOPHIA; Prout, 1918, Novit. zool. 25: 414.

An incorrect subsequent spelling of Gnatholophia Felder, 1874.

GNOPHAELA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 330.

Type-species: Gnophaela aequinoctialis Walker, 1854, ibidem 2: 331, by monotypy. South America.

Gnophaela Walker probably originated from Boisduval who attributed it to himself in Boisduval, 1870, Considérations Lepid. Guatemala: 87.

‡GONOPHAELA; Druce, 1884, Biologia cent.-am. (Zool.) Lepid.-Heterocera 3: pl.12, fig.7.

An incorrect subsequent spelling of Gnophaela Walker, 1854.

GNOPHAELA Herrich-Schäffer, [1855]. See Paraethria Hampson, 1898.

GNOPHRIA Stephens, 1829. See Atolmis Hübner, [1819].

GNOPHRIOIDES Heylaerts, 1891, Annls Soc. ent. Belg. 35

Type-species: Gnophrioides flaviplaga Heylaerts, 1891, ibidem 35 (Bull.): ccccxii, by original designation. JAVA: Préanger.

GOACAMPA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 263

Type-species: Goacampa variabilis Schaus, 1901, ibidem **1901**: 331, pl.12, fig.6, by original designation. Oaxaca.

GOAXIS Schaus, 1901, Trans. ent. Soc. Lond. 1901: 262 (key),

Type-species: Goaxis singularis Schaus, 1901, ibidem 1901: 316, pl.11, fig.13, by original designation. BRAZIL: Petropolis.

GODASA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 271.

Type-species: Godasa maculatrix Walker, [1865] 1864, ibidem 31: 272, by monotypy. Holotype &, Sierra Leone, (BMNH).

G. maculatrix is a junior subjective synonym of Noctua sidae Fabricius, 1794, Ent. Syst. 3 (2): 23.

Godasa was established in the Callimorphidae, now Arctiidae; it was placed in the Agaristidae by Jordan, 1896, in Rothschild & Jordan, Novit. zool. 3: 47; it was placed in the Arctiidae, Hypsinae by Gaede, 1926, in Seitz, Gross-Schmett. Erde 14: 92; and is here again transferred to the Agaristidae on the basis of tympanic and venational characters.

Godasa in the Agaristidae was not included in Nye, 1975, Generic Names Moths World 1.

GOGANA Walker, 1866. See Choerotricha Felder, 1874. LYMA

‡GOGANE; misspelling. See Choerotricha Felder, 1874.

GONEPTERONIA Kenrick, 1917, Trans. ent. Soc. Lond. 1917: NOCTUIDAE

Type-species: Gonepteronia bipunctata Kenrick, 1917, ibidem 1917: 95, pl.3, fig., by monotypy. Type(s) MADAGASCAR,

Gonepteronia in the Noctuidae, Ophiderinae is cited here as it was not included in Nye, 1975, Generic Names Moths World

GONERDA Moore, 1879, Proc. zool. Soc. Lond. 1879; 395.

Type-species: Gonerda perornata Moore, 1879, ibidem 1879: 395, pl.32, fig.2, by monotypy. [INDIA/PAKISTAN]: [Kashmir] Cashmere.

GONIOSEMA Turner, 1899, Trans. Proc. R. Soc. S. Aust. 23:

Type-species: Chiriphe anguliscripta Lucas, 1890, Proc. Linn. Soc. N.S.W. (2) 4: 1079, by monotypy. AUSTRALIA: Queensland, Brisbane.

GONOCLOSTERA Butler, 1877, Ann. Mag. nat. Hist. (4) 20:

Type-species: Gonoclostera latipennis Butler, 1877, ibidem (4) 20: 476, by original designation. JAPAN: Yokohama.

G. latipennis is a subspecies of Pygaera timoniorum Bremer, 1861, Bull. Acad. imp. Sci. St Pétersb. 3: 482.

GONODONTA Hübner, 1818, Zuträge Samml. exot. Schmett.

Type-species: Gonodonta uncina Hübner, 1818, ibidem 1: 11, pl.[6], figs 35, 36, by subsequent designation by Grote, 1902, Allg. Z. Ent. 7: 472. BRAZIL.

Gonodonta is included here as it is linked with a usage of Ptilodon before it became an available name in the Notodontidae.

‡PTILODON Hübner, 1808, Erste Zuträge Samml. exot. Schmett.: 4.

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1966, Bull. zool. Nom. 23 (Opinion 789): 216.

Placed on the Official Index of Rejected and Invalid Generic Names in Zoology with Name No.1881.

The only included name ‡Ptilodon uncium Hübner, 1808, was subsequently made nomenclaturally available as Gonodonta uncina Hübner, 1818.

‡GONOPHAELA; misspelling. See Gnophaela Walker, 1854.

GONOTREPHES Hampson, 1909, Ann. Mag. nat. Hist. (8)

Type-species: Thyrarctia friga Druce, 1906, Ann. Mag. nat. Hist. (7) 17: 406, by original designation. Peru: Carabaya, Quinton.

GOPHA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 81. NOTO

Type-species: Gopha mixtipennis Walker, 1862, ibidem (3) 1: 81, by monotypy. Brazil.

GORGONIDIA Dyar, 1898, Jl N.Y. ent. Soc. 6: 36. ARCT Type-species: Zatrephes garleppi Druce, 1898, Ann. Mag. nat. Hist. (7) 1: 148, by subsequent designation by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 39. BOLIVIA.

GOZARTA Walker, 1869. See Bardaxima Walker, 1858.

GRAFIRIA Draudt, 1932, in Seitz, Gross-Schmett. Erde 6: 1006.

Type-species: *Rifargia steinbachi* Rothschild, 1917, *Novit. zool.* **24**: 255, pl.6, fig.4, by original designation. BOLIVIA: eastern, Buenavista.

The date of publication of the part of Seitz containing page 1006 is cited on page 1001 as 25.xii.1932 in both the German and the English editions.

GRAMMARCTIA Aurivillius, 1899, Ent. Tidskr. 20: 236, 243.

Type-species: Setina bilinea Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 237, by original designation. [SOUTH AFRICA]: Caffraria.

GRAMMIA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 261.

Type-species: *Bombyx quenseli* Paykull, 1793, *Skr. Naturh.-Selsk. Kiøbenhavn* 2 (2): 99, fig., by monotypy. Sweden.

ORODEMNIAS Wallengren, 1885, Skand. Heterocer-Fjärilar 2 (3): 315.

Type-species: *Bombyx quenseli* Paykull, 1793, by monotypy. A junior objective synonym of *Grammia* Rambur, 1866.

‡ORDEMNIAS; Aue, 1927, Lepid. Rdsch. 1: 127.

An incorrect subsequent spelling of *Orodemnias* Wallengren, 1885.

GRAMMOA Aurivillius, 1904, Ark. Zool. 2 (4): 51. LYMA Type-species: Grammoa striata Aurivillius, 1904, ibidem 2 (4): 52, fig.39, by monotypy. [CAMEROUN] Kamerun.

GRANGULA Kiriakoff, 1967, Tijdschr. Ent. 110: 57. NOTO Type-species: Stauropus apicalis Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 59, by original designation. INDIA: Darjeeling.

GRANGULINA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 377.

Type-species: *Grangulina sumatrana* Kiriakoff, 1974, *ibidem* 17: 378, text-fig.3, pl.1, fig.3, by original designation. SUMATRA: northern, Deli, Dolok Merangir.

GRAPHEA Schaus, 1894, Proc. zool. Soc. Lond. 1894: 232.

Type-species: *Graphea marmorea* Schaus, 1894, *ibidem* 1894: 232, by monotypy. VENEZUELA: Aroa.

GRAPHELYSIA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 408.

Type-species: *Elysius strigillata* Rothschild, 1910, *Novit. zool.* 17: 187, pl.14, fig.13, by original designation. Peru: south-east, Carabaya, La Union.

GRAPHIDURA Strand, 1911, Jb. nassau. Ver. Naturk. 64: 121.

Type-species: *Notodonta argenteomaculata* Aurivillius, 1892, *Ent. Tidskr.* **13**: 196, fig.3, by original designation. CAMEROUN/ GABON.

The type-locality of the type-species was not stated, but it was described in a paper entitled 'Schmetterlinge aus Gabun und Camerun'.

GRAPHIDURINA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 254.

Type-species: *Graphidurina auriplaga* Kiriakoff, 1964, *ibidem* 14: 255, pl.1, fig.7, by original designation. China: Kwangtung, Linping.

GRAPHOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: viii, 82, 97.

LITH
The appears Chaphagia bilingata Hampson, 1900, ibiday.

Type-species: *Graphosia bilineata* Hampson, 1900, *ibidem* 2: 97, fig.32, by original designation. New Guinea: [Irian Jaya], [Kapaor] Kapaur.

GRAPTASURA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvii, 90, 421.

Type-species: Cyme polygrapha Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.7, by original designation. [SULAWESI] Celebes: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

GRAPTOPHALERA Kiriakoff, 1967, Biologia gabon. 3: 351.

Type-species: *Graptophalera bicolor* Kiriakoff, 1967, *ibidem* 3: 351, fig.38, by original designation. GABON: Bélinga.

‡GRATHOLOPHIA; misspelling. See Gnatholophia Felder, 1874.

GRIPHOCERURA Kiriakoff, 1958, Mém. Inst. scient. Madagascar (E) 9: 224.

Type-species: Stauropus malgassica Kenrick, 1917, Trans. ent. Soc. Lond. 1917: 85, pl.1, by original designation. MADAGASCAR.

GROTEA Moore, 1866. See Sebastia Kirby, 1892. ARCT

GRUCIA Dyar, 1916, Proc. U.S. natn. Mus. 51: 5.

Type-species: Grucia monacheicauda Dyar, 1916, ibidem 51: 6, by original designation. MEXICO: Hidalgo, Guerrero Mill.

‡CRUCIA; Neave, 1939, Nomencl. zool. 1: 880. An incorrect subsequent spelling of Grucia Dyar, 1916.

GUNDLACHIA Herrich-Schäffer, 1866. See Burtia Grote, 1866. CTEN

GURNA Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 123.

Type-species: *Dysauxes indica* Moore, 1879, *Proc. zool. Soc. Lond.* **1879**: 390, by original designation. [INDIA]: Bombay.

GYARA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 915. PERI

Type-species: Gyara fenestrata Walker, 1855, ibidem 4: 916, by monotypy. [Brazil]: Para and Rio [de] Janeiro.

GYLLA Bryk, 1953, Ark. Zool. (2) 5: 218. LITH Type-species: Gylla crotalistria Bryk, 1953, ibidem (2) 5: 218, by original designation. Peru: Amazonas, Rio Autaz.

GYMNASURA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvii, 86, 425.

Type-species: Calligenia saginaea Turner, 1899, Trans. Proc. R. Soc. S. Aust. 1899: 11, by original designation. AUSTRALIA: Queensland, Cairns.

GYMNELIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 152.

Type-species: Glaucopis laennus Walker, 1854, ibidem 1: 154, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 136. Brazil: Rio [de] Janeiro.

Gymnelia Walker probably originated from Boisduval who attributed it to himself in Boisduval, 1870, Considérations Lépid. Guatemala: 80.

NOTO

Gymnelia was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 189.

BOMBILIODES Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xii, 24, 179.

Type-species: Glaucopis laennus Walker, 1854, by original designation.

A junior objective synonym of Gymnelia Walker, 1854.

‡BOMBYLIODES; Neave, 1939, Nomencl. zool. 1: 449.

An incorrect subsequent spelling of *Bombiliodes* Hampson, 1898.

GYMNOCHROMA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 92, 537.

Type-species: *Nudaria fulvipicta* Hampson, 1896, *Fauna Br. India* (Moths) **4**: 501, by original designation. [INDIA]: [Assam], Khasia Hills.

GYMNOPODA Felder, 1874. See Apeplopoda Watson nom.
CTEN

GYNAEPHORA Hübner, [1819] 1816, Verz. bekannter Schmett.: 161.

Type-species: *Phalaena selenitica* Esper, 1789, *Die Schmett.* 3 (Suppl.) 3 Abschn.: 18, pl.82, fig.5, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 473. [France]: [Moselle], Lemberg.

CLADOPHORA Geyer, 1832, in Hübner, Zuträge Samml. exot. Schmett. 4: 33.

Type-species: *Phalaena selenitica* Esper, 1789, by subsequent designation by Collenette, 1935, *Stylops* 4: 243.

Cladophora is a junior objective synonym of Gynaephora Hübner, [1819].

GYNOPHORA Agassiz, 1847, Nomencl. zool. (Index univl.): 169.

An unjustified emendation of Gynaephora Hübner, [1819]. The 'Index universalis' was published as Fascicle 12 of the Nomenclator zoologicus; Gynophora is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

GYNOPHORA Agassiz, 1847. See Gynaephora Hübner, [1819].

HABROCHROMA Turner, 1940, Proc. R. Soc. Qd 51: 100.

Type-species: Asura ectophaea Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 451, pl.31, fig.9, by monotypy. AUSTRALIA: [Northern Territory], Port Darwin.

HABROPHYLLA Turner, 1921, *Proc. Linn. Soc. N.S.W.* **45**: 476 (key), 482.

Type-species: *Euproctis euryzona* Lower, 1902, *Trans. Proc. R. Soc. S. Aust.* **26**: 213, by monotypy. Australia: South Australia, Henley Beach.

HADESINA Warren, 1900, Novit. zool. 7: 126. DIOP Type-species: Hadesina limbaria Warren, 1900, ibidem 7: 127, by original designation. ECUADOR: Chimbo.

Hadesina was established in the Geometridae, Cyllopodinae, now Sterrhinae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 418.

HAEMANOTA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 3, 87.

Type-species: *Haemanota rubriceps* Hampson, 1901, *ibidem* 3: 87, fig.62, by original designation. BRAZIL: [Espirito Santo] Espiritu Santo.

HAEMAPHLEBIA Hampson, 1911. See Haemaphlebiella Collins, 1962.

HAEMAPHLEBIELLA Collins, 1962, Ann. Mag. nat. Hist. (13) 5: 125.

Type-species: *Baritius venata* Rothschild, 1909, *Ann. Mag. nat. Hist.* (8) 4: 210, by original designation (for *Haemaphlebia* Hampson, 1911). Surinam: Maroewym Valley, Aroewarwa Creek, [Brazil]: Upper Amazon, Fonte Boa.

Haemaphlebiella was established as the objective replacement name for Haemanhlebia Hampson, 1911

ment name for Haemaphlebia Hampson, 1911.

HAEMAPHLEBIA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 407.

Type-species: *Baritius venata* Rothschild, 1909, by original designation.

A junior homonym of *Haemaphlebia* Hampson, 1910, *Cat. Lepid. Phalaenae Br. Mus.* 10: 3, 203, – Lepid., Noctuidae. The objective replacement name is *Haemaphlebiella* Collins, 1962.

HAEMATERION Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.49, fig.262.

Type-species: *Haematerion braco* Herrich-Schäffer, [1855] 1850-1858, *ibidem* 1 (1): wrapper, pl.49, fig.262 (but attributed to Boisduval, an incorrect authorship), by monotypy. Brazil.

Haematerion Herrich-Schäffer probably originated from Boisduval who attributed it to himself in Boisduval, 1870, Considérations Lépid. Guatemala: 80.

HAEMATOMIS Schaus, 1899, Jl N.Y. ent. Soc. 7: 215. LITH Type-species: Haematomis uniformis Schaus, 1899, ibidem 7: 215, by monotypy. MEXICO: Oaxaca.

This generic name probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson, 1900.

HAEMATOMIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 86, 514.

Type-species: Lithosia mexicana Druce, 1885, Biologia cent.am. (Zool.) Lepid.-Heterocera 1: 131, pl.13, figs 2, 3, by original designation. Mexico: Cordova; Durango State, Guatemala: San Geronimo, Pantaleon.

A junior homonym of *Haematomis* Schaus, 1899. There is no objective replacement name but the type-species of *Haematomis* Hampson, 1900, is currently considered to be congeneric with the type-species of *Haematomis* Schaus, 1899. The latter is therefore available for use as a subjective replacement name

‡HAEMATOMONIS; Dyar, [1903] 1902, Bull. U.S. natn. Mus. 52: 80.

An incorrect subsequent spelling of *Haematomis* Hampson, 1900.

HAEMATOMIS Hampson, 1900. See *Haematomis* Schaus, 1899.

‡HAEMATOMONIS; misspelling. See Haematomis Schaus, 1899.

HAGAPTERYX Matsumura, 1920, Zool. Mag. Tokyo 32: 149.

Type-species: *Lophopteryx admirabilis* Staudinger, 1887, *in* Romanoff, *Mém. Lépid.* 3: 224, pl.12, fig.9, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

Hagapteryx was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 812, 813.

‡HALESIDOTA; misspelling. See Halysidota Hübner, [1819].

‡HALIACTINIDIA; misspelling. See Heliactinidia Hampson, 1901.

‡HALISIDOTA; misspelling. See Halysidota Hübner, [1819].

‡HALIURA; misspelling. See Heliura Butler, 1876. CTEN

HALONE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 540.

Type-species: *Halone sobria* Walker, 1854, *ibidem* 2: 540, by monotypy. [South Africa]: [Durban] Port Natal.

HALSEYELLA Nye nom. n. for Crinola Leech, 1890. LYMA Type-species: Crinola flavicollis Leech, 1890, Entomologist 23: 111, by monotypy. China: Chang Yang.

This name is proposed in appreciation of the help given by Miss Sheila H. Halsey in the compilation of this catalogue.

CRINOLA Leech, 1890, Entomologist 23: 111.

Type-species: Crinola flavicollis Leech, 1890, by monotypy. A junior homonym of Crinola Rafinesque, 1815, Analyse Nature: 151, – Vermes. The objective replacement name is Halseyella Nye nom. n.

HALURGIA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 95, 97.

Type-species: *Halurgia aurorina* Herrich-Schäffer, 1855, *ibidem* 6: 97, by monotypy. Type-locality: not stated; identity of species uncertain.

Halurgia was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, Corresp Bl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

HALYSIDOTA Hübner, [1819] 1816, Verz. bekannter Schmett.: 170.

Type-species: *Phalaena tessellaris* J.E Smith, 1797, *in* Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 149, pl.75, but included by Hübner as 'Halisidota Tesselaris Abb.', an incorrect (of a multiple) original spelling of the generic name, and an incorrect subsequent spelling of the specific name, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 209, but cited as ‡*Halisidota*, an incorrect (of a multiple) original spelling. [U.S.A.]: Georgia.

Halysidota was the spelling adopted by Berg, 1882, An. Soc. cient. argent. 13: 214, as first reviser, from a multiple original spelling which included both Halysidota and ‡Halisidota. Hübner opened his list of species with the heading '... Halysidotae', the Latin plural of Halysidota; he then included three species, the first being cited as 'Halisidota Tesselaris Abb.'.

‡HALISIDOTA; Hübner, [1819] 1816, Verz. bekannter Schmett.: 170.

An incorrect original spelling of *Halysidota* Hübner, [1819]. ‡*Halisidota* was the spelling rejected by Berg, 1882, *An. Soc. cient. argent.* 13: 214 from a multiple original spelling.

ALYSIDOTA Agassiz, 1847, Nomencl. zool. (Index univl.): 15, 172.

An unjustified emendation of *Halysidota* Hübner, [1819]. The 'Index universalis' was published as Fascicle 12 of the

Nomenclator zoologicus; Alysidota is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

‡HALESIDOTA; Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 732.

An incorrect subsequent spelling of *Halysidota* Hübner, [1819].

HAMIDONTA Dognin, 1911, Hétérocères nouv. Am. Sud 2: 36.

Type-species: *Hamidonta unca* Dognin, 1911, *ibidem* 2: 36, by original designation. FRENCH GUIANA: St Jean du Maroni.

HAMMATOPHORA Westwood, 1843. See *Phalera* Hübner, [1819].

HAMPSONITA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 211.

Type-species: Cerura esmeralda Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 458, by original designation. [SOUTH AFRICA]: Pretoria.

HANOISIELLA de Joannis, 1928, Annls Soc. ent. Fr. 97: 274.

Type-species: *Hanoisiella exilicosta* de Joannis, 1928, *ibidem* 97: 274, pl.4, fig.3, by original designation. [VIETNAM]: Hanoi.

HAPIGIA Guenée, 1852, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 6: 375.

Type-species: *Hapigia nodicornis* Guenée, 1852, *ibidem* 6: 376, pl.12, fig.7, by monotypy. French Guiana: Cayenne.

From a multiple original spelling in which *Hapigia* was used twice (on pages 375 and 376), and ‡*Hapygia* was used three times (in the legend to pl.12; in the 'Explication des Planches' part 1, page 6; and in the index in volume 7, page 408, also published in 1852), we have adopted *Hapigia*, the spelling in current use, as the correct original spelling.

Hapigia was established in the Noctuidae; it was placed in the Notodontidae by Druce, 1887, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 243.

‡HAPYGIA; Guenée, 1852, Hist. nat. Insectes (Spec. gén. Lépid.) 6: legend to pl.12; Explication des Planches (1):6; and 1852, ibidem 7: 408.

An incorrect (of a multiple) original spelling of *Hapigia* Guenée, 1852.

HAPIGIODES Dyar, 1910, Proc. U.S. natn. Mus. 38: 259.

Type-species: *Hapigia xolotl* Schaus, 1892, *Proc. zool. Soc. Lond.* **1892**: 339, by original designation. Mexico: Paso de San Juan.

HAPLOA Hübner, [1820] 1816, Verz. bekannter Schmett.: 182.

Type-species: *Phalaena clymene* Brown, 1776, *New Illust. Zool.*: 96, pl.[38], by subsequent designation by Hampson, 1901, *Cat. Lepid. Phalaenae Br. Mus.* 3: 492. Type-locality: cited as Ceylon, but this was probably an error as *clymene* Brown is distributed only in U.S.A.

Invalid type-species: Bombyx colona Hübner, [1803], a nominal species not originally included in Haploa, and not linked with one of the originally included nominal species when cited by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 252. In his synonymy of colona, Kirby included Noctua clymene Esper, 1794, a secondary homonym of P. clymene Brown. Kirby did cite P. clymene Brown as a separate species so his designation of colona Hübner was probably an inadvertent error.

HAPLONERITA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 411.

Type-species: Idalus simplex Rothschild, 1909, Novit. zool.

16: 23, pl.4, fig.10, by original designation. Surinam: Maroewym Valley, Aroewara Creek.

HAPLOPSEUSTIS Meyrick, 1902 April, Trans. ent. Soc. Lond. 35: 34.

Type-species: *Haplopseustis erythrias* Meyrick, 1902 April, *ibidem* 35: 34, by monotypy. Syntypes 28, Australia: North Australia, Port Darwin, Queensland, Brisbane, (BMNH).

Haplopseustis was established in the Caradrinidae, now Noctuidae; this placing was confirmed by Turner, 1920, Proc. Linn. Soc. N.S.W. 45: 475; it was placed in the Lymantriidae by Strand, 1925, in Seitz, Gross-Schmett. Erde 10: 362, but, as stated on page 36, this part of the work was ready in 1918; publication was delayed due to World War II, so Strand had not seen Turner's 1920 work. Haplopseustis was placed in the Noctuidae, Acontiinae by Nye, 1975, Generic Names Moths World 1: 230.

See also Acnissa Turner, 1902 December.

‡HOPLOPSEUSTIS; Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 77.

An incorrect subsequent spelling of *Haplopseustis* Meyrick, 1902.

‡Hoplopseustis is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡HOPLOPSEUSTRIAS; Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 77.

An incorrect subsequent spelling of *Haplopseustis* Meyrick, 1902.

‡Hoplopseustrias is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

HAPLOZANA Aurivillius, 1901, Ent. Tidskr. 22: 121. NOTO Type-species: Haplozana nigrolineata Aurivillius, 1901, ibidem 22: 122, fig.23, by monotypy. [ZAIRE] Congo: between Kassongo and Stanley Falls.

See also Hypophiala Janse, 1920.

‡HAPODONTA; misspelling. See Hupodonta Butler, 1877.

‡HAPYGIA; misspelling. See Hapigia Guenée, 1852. NOTO

HARAPA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 47.

LYMA

Type-species: *Harapa testacea* Moore, 1879, *ibidem* (1): 47, pl.2, fig.15, by monotypy. INDIA: [W. Bengal, Darjeeling] Darjiling.

HARDINGIA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 260 (key), 293.

Type-species: *Hardingia roberti* Schaus, 1901, *ibidem* **1901**: 293, pl.11, fig.10, by original designation. COLOMBIA.

HARMA Walker, [1858]. See Hippia Möschler, 1878. NOTO

HARPANDRYA Bryk, 1913, Societas ent. 28: 100. NOTO Type-species: Harpandrya aeola Bryk, 1913, ibidem 28: 100, text-fig., by monotypy. CAMEROUN: Lolodorf.

‡HARPYA; misspelling. See Harpyia Ochsenheimer, 1810.

HARPYIA Ochsenheimer, 1810, Schmett. Eur. 3: 19. NOTO Type-species: Bombyx milhauseri Fabricius, 1775, Syst. Ent.: 577, by subsequent designation by Blanchard, [1847], in Cuvier, Règne Animal (Disciples Edn) (Insectes) 2: title-page and pl.153 bis. [GERMANY]: Dresden.

On the title-page of each volume of the Disciples Edition, it was stated in French that the edition was 'accompanied by plates representing the types of all the genera, the distinctive

characters of the different groups and the modification of structure on which this classification is based'.

The dates of publication of the parts of the Disciples Edition were discussed and listed by Cowan, 1976, J. Soc. Bibliphy nat. Hist. 8 (1): 32-64.

Invalid designation of type-species: Phalaena vinula Linnaeus, 1758, was designated by Boisduval, 1836, Hist. nat. Insectes (Spec. gén. Lépid.) 1: 134. In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different typespecies for each generic name. In his 'Exposé de notre méthode', pages 155-690, no type-species designation was made for any of the genera that he himself used. Under the Code, Article 69(a)(iii), the type-species designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other authors.

Acceptance of Boisduval's designation in this case would place *Harpyia* Ochsenheimer, 1810, as a junior objective synonym of *Cerura* Schrank, 1802.

Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 238, incorrectly attributed to himself the authorship of Harpyia Ochsenheimer, 1810, but misspelt the name as ‡Harpya.

HOPLITIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 147.

Type-species: Bombyx terrifica [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 63, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 595, but cited as milhauseri Fabricius which Kirby placed as the senior synonym of 'terrifica Borkhausen', an incorrect authorship; see below. [AUSTRIA]: Vienna district.

A junior homonym of *Hoplitis* Klug, 1807, *Magazin Insektenk*. (*Illiger*) 6: 225, – Insecta, Hymenoptera. There is no objective replacement name, but *B. terrifica* is a junior subjective synonym of *Bombyx milhauseri* Fabricius, 1775, the typespecies of *Harpyia* Ochsenheimer, 1810; the latter is thus available for use as a subjective replacement name.

When Kirby designated as type-species *Bombyx milhauseri* Fabricius, 1775, a nominal species not originally included in *Hoplitis*, he also (on the same page) placed *terrifica*, a nominal species originally included in *Hoplitis*, as a junior subjective synonym of *milhauseri*. Under the *Code*, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species.

‡HARPYA; Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 238.

An incorrect subsequent spelling of, and an incorrect authorship for *Harpyia* Ochsenheimer, 1810.

‡HOPLITES; Agassiz, 1846, Nomencl. zool. (Nom. syst. Generum Anim.) Lepid.: 36; also 1847, ibidem (Index univl.): 185.

An incorrect subsequent spelling of *Hoplitis* Hübner, [1819]. The Lepidoptera part of *Nomencl. zool*. (Nom. syst. Generum Anim.) was included in Fascicles 9 and 10 issued in one wrapper dated 1846. The Index universalis formed Fascicle 12 of the work; the wrapper is dated 1847, the title-page is dated 1846.

‡Hoplites Agassiz, 1846, was placed by the International Commission on Zoological Nomenclature on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.289 as '(emend. of Hoplitis Hübner, [1819]) (suppressed under the Plenary Powers for the purposes both of the Law of Priority and of the Law of Homonymy)'. There is no justification for regarding this usage by Agassiz as an emendation,

but the result is the same as either way this usage is not nomenclaturally available.

HYBOCAMPA Lederer, 1853, Verh. zool.-bot. Ver. Wien 2 (Abh.): 78.

Type-species: Bombyx milhauseri Fabricius, 1775, by monotypy.

A junior objective synonym of Harpyia Ochsenheimer, 1810.

‡ARPYIA; Balducci, 1905, Boll. Soc. ent. ital. 36: 117.

An incorrect subsequent spelling of *Harpyia* Ochsenheimer, 1810.

‡HOPLISIA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 428.

An incorrect subsequent spelling of Hoplitis Hübner, [1819].

HARPYIAS Hübner, [1819]. See Furcula Lamarck, 1816.

HASSLERIA Jörgensen, 1934, Dt. ent. Z. Iris 48: 64. LITH Type-species: Hassleria majas Jörgensen, 1934, ibidem 48: 65, by original designation. PARAGUAY: Cordillera de Villarrica, Mburero.

HATIMA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 450.

Type-species: *Hatima semirufescens* Walker, 1865, *ibidem* 32: 450, by monotypy. U.S.A.

HEBENA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1722.

Type-species: *Hebena venosa* Walker, 1856, *ibidem* 7: 1723, by monotypy. [South Africa]: [Durban] Port Natal.

H. venosa is a junior subjective synonym of Automolis lateritia Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.51, fig.274.

Hebena was established in the Arctiidae; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 34.

See also Hexaneura Wallengren, 1860.

HECTOBROCHA Meyrick, 1886, *Proc. Linn. Soc. N.S.W.* (2) 1: 692, 706.

Type-species: *Hectobrocha pentacyma* Meyrick, 1886, *ibidem* (2) 1: 707, by monotypy. AUSTRALIA: Queensland.

HECTOGAMA Meyrick, 1889, Trans: ent. Soc. Lond. 1889: 465.

Type-species: *Hectogama dissozona* Meyrick, 1889, *ibidem* 1889: 465, by monotypy. New Guinea.

IIELGA Kiriakoff, 1962, Tijdschr. Ent. 105: 257. NOTO Type-species: Helga infans Kiriakoff, 1962, ibidem 105: 258, fig.30, by original designation. [ZIMBABWE-]RHODESIA: Nyamandhlovu.

IIELIACTINIDIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 5, 205.

Type-species: *Bepara chiguinda* Druce, 1885, *Proc. zool. Soc. Lond.* 1885: 524, pl.32, fig.10, by original designation. ECUADOR: Chiguinda.

‡HALIACTINIDIA; Grünberg, 1910, Arch. Naturgesch. 73 (2): 245.

An incorrect subsequent spelling of *Heliactinidia* Hampson, 1901.

HELIOCAES Turner, 1940. See Cheliosea Watson nom. n. ARCT

HELIORABDIA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 402.

Type-species: *Chrysorabdia taiwana* Wileman, 1910, *Ento-mologist* 43: 178, by original designation. [TAIWAN] Formosa: Arizan.

Hampson, 1911, established *Heliorabdia* for the male syntype of *C. taiwana* Wileman, here designated as the LECTOTYPE of *C. taiwana* Wileman, 1910, (the female syntype of *C. taiwana* Wileman was a different species named *Chrysorabdia vilemani* Hampson, 1911, *ibidem* (8) 8: 401).

HELIOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 87, 275.

Type-species: *Pallene jucunda* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 2: 543, by original designation. [AUSTRALIA]: New South Wales, Sydney.

HELIOZONA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 5, 218.

Type-species: Satara lianga Semper, 1899, Reisen Archipel Philipp. Semper (2) 6: 490, pl.55, fig.8, by original designation. Philippines: E. Mindanao, Lianga.

HELIURA Butler, 1876, J. Linn. Soc. (Zool.) 12: 417. CTEN Type-species: Heliura solicauda Butler, 1876, ibidem 12:

418, by original designation. HONDURAS.

H. solicauda Butler is a junior subjective synonym of Eucereon rhodophila Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1638, but cited as ‡Eucerea, an incorrect subsequent spelling.

‡HELUIRA; Schaus, 1889, Entomologica am. 5: 90.
An incorrect subsequent spelling of Heliura Butler, 1876.

‡HALIURA; Grünberg, 1913, Arch. Naturgesch. 78 (B) 7: 113. An incorrect subsequent spelling of Heliura Butler, 1876.

‡HELUIRA; misspelling. See Heliura Butler, 1876. CTEN

HEMEROCAMPA Dyar, 1897, Can. Ent. 29: 13 (key), 15.

Type-species: *Phalaena leucostigma* Smith, 1797, *in* Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 157, pl.79, by original designation. [U.S.A.]: Georgia and Virginia.

HEMEROPHANES Collenette, 1953, Ann. Mag. nat. Hist. (12) 6: 571.

Type-species: *Dasycampa ianthina* Janse, 1915, *Ann. Transv. Mus.* 5: 26, pl.3, figs 3, 4, by original designation (of *Dasycampa* Janse, 1915). South Africa: Barberton and Woodbush Village.

D. ianthina is a junior subjective synonym of Heteronygmia flammeola Distant, 1899, Ann. Mag. nat. Hist. (7) 4: 360.

Hemerophanes was proposed as the objective replacement name for Dasycampa Janse, 1915, but Collenette cited as typespecies, H. flammeola Distant, the senior subjective synonym of the type-species.

DASYCAMPA Janse, 1915, Ann. Transv. Mus. 5: 25.

Type-species: Dasycampa ianthina Janse, 1915, by original designation.

A junior homonym of *Dasycampa* Guenée, 1837, *Annls Soc.* ent. Fr. 6: 224, – Lepid., Noctuidae. The objective replacement name is *Hemerophanes* Collenette, 1953.

HEMICERAS Guenée, 1852, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 6: 379.

Type-species: *Hemiceras cadmia* Guenée, 1852, *ibidem* 6: 383, pl.13, fig.2, by subsequent designation by Grote, 1874, *Bull. Buffalo Soc. nat. Sci.* 2: 29. NORTH AMERICA.

Hemiceras was established in the Noctuidae; it was placed in the Notodontidae by Druce, 1887, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 249.

See also Comidava Walker, [1863].

HEMIFENTONIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 152.

Type-species: Parafentonia inconspicua Kiriakoff, 1964, Bonn. zool. Beitr. 14: 278, abb.4, fig.35, pl.3, fig.37, by original designation (for Parafentonia Kiriakoff, 1964). CHINA: Chekiang, east Tien-mu-Shan.

Hemifentonia was established as the objective replacement

name for Parafentonia Kiriakoff, 1964.

PARAFENTONIA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 277. Type-species: Parafentonia inconspicua Kiriakoff, 1964, by original designation.

A junior homonym of Parafentonia Roepke, 1944, Natuurh. Maandbl. 33: 19, - Lepid., Notodontidae. The objective replacement name is Hemifentonia Kiriakoff, 1967.

‡HEMIHYALAE; misspelling. See Hemihyalea Hampson, 1901.

HEMIHYALEA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: ix, 6, 129.

Type-species: Phaegoptera cornea Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.14, figs 61, 62, by original designation. VENEZUELA.

‡HEMIHYALAE; Strand, 1919, in Wagner, Lepid. Cat. 22:

An incorrect subsequent spelling of Hemihyalea Hampson, 1901.

HEMIPECTEROS Schaus, 1920, Insecutor Inscit. menstr. 8: 152.

Type-species: Hemipecteros arthemis Schaus, 1920, ibidem 8: 153, by original designation. [GUYANA] British Guiana.

HEMIPSILIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 91, 532.

Type-species: Nudaria coavestis Hampson, 1894, Fauna Br. India (Moths) 2: 124, by original designation. [INDIA]: Sikkim.

HEMIRHIPIDIA Kiriakoff, 1955, Mitt. münch. ent. Ges. 44-45: 256.

Type-species: Rhipidarctia danieli Kiriakoff, 1955, ibidem 44-45: 256, text-fig.1, pl.4, fig.10, by original designation. [CAMEROUN] Kamerun: Debundscha.

Hemirhipidia was established as a subgenus of Rhipidarctia

Kiriakoff, 1953.

HEMITUERTA Kiriakoff, 1976. **98**: 106. Tierreich AGARISTIDAE

Type-species: Paratuerta nana Hampson, 1916, in Poulton, Proc. zool. Soc. Lond. 1916: 112, pl.1, fig.17, by original designation. Syntypes 18, 19, [Somali Republic] Somaliland: Mandera, (BMNH).

Hemituerta in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

HEMONIA Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 425.

Type-species: Hemonia orbiferana Walker, 1863, ibidem 28:

426, by monotypy. [SRI LANKA] Ceylon.

Hemonia was established in the Tortricidae; it was placed in the Lithosiidae, now Arctiidae Lithosiinae, by Moore, [1887] 1884-7, Lepid. Ceylon 3: 537.

HENOSIS Wallengren, 1865. See Zastonia Herrich-Schäffer, [1858].

HEORTA Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1664.

Type-species: Heorta roseoalba Walker, 1858, ibidem 15: 1665, by monotypy. [Brazil]: [R. Amazon], [Tefé] Ega.

Heorta was established in the Noctuidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 579.

HEPATOPHALERA Kiriakoff, 1960, Revue Zool. Bot. afr. 61:

Type-species: Hepatophalera decipiens Kiriakoff, 1960, ibidem 61: 57, fig.9, by original designation. [ZAIRE]: Sankuru, Katako-Kombe.

HEPTAPTOSIS Talbot, 1929, Bull. Hill Mus. Witley 3: 59.

Type-species: Heptaptosis joicevi Talbot, 1929, ibidem 3: 61, text-fig.1, pl.2, fig.7, by original designation. [WEST AFRICA]: [Gulf of Guinea], Principe Island.

HERACULA Moore, [1866] 1865, Proc. zool. Soc. Lond. 1865:

Type-species: Heracula discivitta Moore, [1866] 1865, ibidem 1865: 804, pl.43, fig.2, by monotypy. [INDIA]: [Himachal Pradesh], Darjeeling.

HERAIA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 57.

Type-species: Stauropus thalassina Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 470, by original designation. [ZIM-BABWE- RHODESIA: Salisbury.

HERBERTINA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 263 (key), 329.

Type-species: Lophopteryx eumeta Druce, 1900, Ann. Mag. nat. Hist. (7) 5: 514, by original designation. COLOMBIA: Sierra del Libano.

HERBULOTEDRA Thiaucourt, 1978, Lambillionea 78: 49.

Type-species: Urgedra palmeri Druce, 1911, Ann. Mag. nat. Hist. (8) 8: 146, by original designation. Colombia: western, San Antonio.

In the original combination of the type-species the generic name was published as ‡Urgerda, an incorrect subsequent spelling.

HEREA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 197.

Type-species: Glaucopis metaxanthus Walker, 1854, ibidem 1: 198, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 367. [Brazil]: Para.

Herea was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 367.

‡HERMINA; misspelling. See Hyrmina Walker, 1854.

HERONINA Kiriakoff, 1955, Mitt. münch. ent. Ges. 44-45:

Type-species: Anace herona Druce, 1887, Proc. zool. Soc. Lond. 1887: 669, by original designation. [CAMEROUN]: [Mongo ma Lobah] Mongo-ma Lubah [4°12'N 9°11'E].

Heronina was established as a subgenus of Balacra Walker, 1856.

HESTIARCHA Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 691, 736.

Type-species: Hestiarcha pyrrhopa Meyrick, 1886, ibidem (2) 1: 736, by monotypy. Australia: South Australia, Port Lincoln.

HESUDRA Moore, 1878, Proc. zool. Soc. Lond. 1878: 12. LITH Type-species: *Hesudra divisa* Moore, 1878, *ibidem* **1878**: 12, pl.1, fig.4, by monotypy. INDIA: Darjeeling.

HESYCHOPA Turner, 1940, Proc. R. Soc. Qd 51: 70. LITH Type-species: Lithosia chionora Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 702, by original designation. AUSTRALIA: New South Wales, Sydney.

HETERALLACTIS Meyrick, 1886, *Proc. Linn. Soc. N.S.W.* (2) 1: 691, 703.

Type-species: *Heterallactis euchrysa* Meyrick, 1886, *ibidem* (2) 1: 703, by monotypy. Australia: Queensland, Brisbane.

HETEROCAMPA Doubleday, 1841, Entomologist 1: 55.

Type-species: *Heterocampa astarte* Doubleday, 1841, *ibidem* 1: 57, by monotypy. U.S.A.: East Florida, St John's Bluff. See also *Balia* Doubleday, 1869, and *Trichotis* Felder, 1874.

HETERODONTA Duponchel, [1845]. See Spatalia Hübner, [1819].

HETERODONTIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.25.

Type-species: *Heterodontia tricolor* Felder, 1874, *ibidem* 2 (Abt.2): pl.105, fig.25, by monotypy. [Brazil]: Amazon R., cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 5.

‡HETEROMELESE Zikan, 1968. See Melese Walker, 1854.

‡HETEROMORPHA Hübner, [1806]. See Diloba Boisduval, [1837].

HETEROMORPHA Hübner, 1822. See Diloba Boisduval, [1837].

HETERONYGMIA Holland, 1893, Psyche, Camb. 6: 414.

Type-species: *Heteronygmia rhodapicata* Holland, 1893, *ibidem* 6: 416, 567, pl.10, fig.15, by subsequent designation by Bryk, 1934, *in* Strand, *Lepid. Cat.* 62: 308. [GABON]: [Ogooué] Ogové R. valley (see page 373).

HETEROTROPA Turner, 1940, Proc. R. Soc. Qd 51: 115.

Type-species: *Heterotropa fastosa* Turner, 1940, *ibidem* 51: 115, by monotypy. Australia: Darwin; Cape York; and Kuranda.

HEXAFRENUM Matsumura, 1925, Zool. Mag. Tokyo 37: 400.

Type-species: Hexafrenum maculifer Matsumura, 1925, ibidem 37: 400, pl.6, fig.4, by original designation. TAIWAN.

In the species-heading, the type-species name was published as ‡'Hexafrenum macurifer', each name an incorrect (of a multiple) original spelling.

HEXANEURA Wallengren, 1860, Wien. ent. Monatschr. 4: 164.

Type-species: *Hexaneura cinnamomea* Wallengren, 1860, *ibidem* 4: 164, by original designation, under the *Code*, Article 68(a)(i). [SOUTH AFRICA]: Caffraria.

H. cinnamomea is a junior subjective synonym of Automolis lateritia Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.51, fig.274.

Hexaneura was established among Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 433; it was placed in

the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 143; and was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 34.

See also Hebena Walker, 1856.

HIERA Druce, 1885, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 142.

Type-species: *Hiera gyge* Druce, 1885, *ibidem* 1: 142, pl.14, fig.10, by monotypy. Panama: Volcan de Chiriqui.

HIEROPHANTA Meyrick, 1894. See Leucodonta Staudinger, 1892.

HIGENA Matsumura, 1925, Zool. Mag. Tokyo 37: 394.

Type-species: *Higena plumigera* Matsumura, 1925, *ibidem* 37: 394, pl.7, fig.5, by original designation. TAIWAN.

HILEMA Agassiz, 1847. See Eilema Hübner, [1819]. LITH

HIMALA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 57.

Type-species: Redoa argentea Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 827, by monotypy. [INDIA]: Assam.

HIMERARCTIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 36.

Type-species: *Automolis docis* Hübner, [1831] 1825, *Zuträge Samml. exot. Schmett.* 3: 32, pl.[93], figs 537, 538, by original designation. [French Guiana]: Cayenne.

HIMEROPTERYX Staudinger, 1887, in Romanoff, Mém. Lépid. 3: 228.

Type-species: *Himeropteryx miraculosa* Staudinger, 1887, *ibidem* 3: 228, pl.12, fig.10, by monotypy. [U.S.S.R.]: Raddefka.

‡HIPOCRITA Hübner, [1806]. See Tyria Hübner, [1819].

‡HIPOCRITA Hübner, 1808. See Apistosia Hübner, 1818.

Name not used again in Lepidoptera except as a misspelling of *Hypocrita* Hübner, [1807].

‡HIPOCRITA; misspelling. See Hypocrita Hübner, [1807].

‡HIPOGYMNA Hübner, [1806]. See Penthophera Germar, 1812.

HIPPIA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 693.

Type-species: *Phalaena mumetes* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 128, pl.82, fig.A, by monotypy. SURINAM.

P. mumetes is dated from the wrapper of part 7, not from the title-page of volume 1 which is dated 1779.

HARMA Walker, [1858] 1857, List Specimens lepid. Insects Colln Br. Mus. 13: 1078 (key), 1105.

Type-species: *Harma vittipalpis* Walker, [1858] 1857, *ibidem* 13: 1106, by monotypy. [Dominican Republic]: St Domingo.

A junior homonym of *Harma* Doubleday, [1848] 1850-1852, in Doubleday & Westwood, *Genera diurn. Lepid.* (2): pl.40, – Lepid., Nymphalidae. There is no objective replacement name, but *H. vittipalpis* is congeneric with *Phalaena mumetes* Cramer, 1775, the type-species of *Hippia* Möschler, 1878; the latter is thus available for use as a subjective replacement name.

Harma Walker, [1858], was established in the Noctuidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 573.

HIPPOLA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 222. CTEN

Type-species: Euchromia ferox Walker, 1854, ibidem 1: 223, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 305, but cited as ‡ 'ipomeae', an incorrect subsequent spelling of Glaucopis ipomoeae Harris, 1839, a nominal species not originally included in Hippola Walker. Hampson, on page 307, placed ipomoeae as the senior synonym of ferox, a nominal species originally included in Hippola. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. [U.S.A.] United States.

E. ferox is a junior subjective synonym of Glaucopis ipomoeae

Harris 1839, Am. J. Sci. 36: 316.

Hippola was established as a subgenus of Euchromia Hübner, [1819].

See also Syntomeida Harris, 1839.

HIPPURARCTIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 32.

Type-species: *Hippurarctia vicina* Kiriakoff, 1953, *ibidem* **26**: 33, by original designation. [ZAIRE]: Kivu.

HIRA Walker, 1854. See Euchromia Hübner, [1819]. CTEN

HIRADONTA Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 31, 36.

Type-species: *Hiradonta takaonis* Matsumura, 1924, *ibidem* 9: 31, by original designation. [JAPAN]: Honshu, Mt Takao near Tokyo; Chichibu by S. Hirayama.

HIRSUTOPALPIS Bethune-Baker, 1904, Novit. zool. 11: 375.

Type-species: *Hirsutopalpis fasciata* Bethune-Baker, 1904, *ibidem* 11: 375, by original designation. New Guinea: [Papua New Guinea], Ekeikei; Dinawa.

Hirsutopalpis was established in the Notodontidae; it was placed in the Noctuidae, Ophiderinae by Nye, 1975, Generic

Names Moths World 1: 242.

‡HISTIAEA; misspelling. See Histioea Walker, 1854.

‡HISTIOCA; misspelling. See Histioea Walker, 1854.

HISTIOEA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 216.

Type-species: Euchromia proserpina Hübner, 1823, Zuträge Samml. exot. Schmett. 2: 10, pl.[39], figs 221, 222, by subsequent designation by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 310 (but cited for ‡Histiaea, an incorrect subsequent spelling). BRAZIL.

Histioea was established as a subgenus of Euchromia Hübner, [1819].

‡HISTIOCA; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1624.

An incorrect subsequent spelling of Histioea Walker, 1854.

‡HISTIAEA; Druce, 1896, Ann. Mag. nat. Hist. (6) 18: 29.

An incorrect subsequent spelling of *Histioea* Walker, 1854; also used by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: xv, 25, 310.

‡HISTIXEA; Rothschild, 1911, Novit. zool. 18: 39.

An incorrect subsequent spelling of Histioea Walker, 1854.

‡HISTIXEA; misspelling. See Histioea Walker, 1854.

HOBAPROMEA Watson nom. n. for Baeomorpha Turner, 1940.

Type-species: *Baeomorpha cleta* Turner, 1940, *Proc. R. Soc. Qd* 51: 115, by monotypy (of *Baeomorpha* Turner, 1940). Australia: N. Queensland, Cape York.

BAEOMORPHA Turner, 1940, Proc. R. Soc. Qd 51: 115.

Type-species: Baeomorpha cleta Turner, 1940, by monotypy. A junior homonym of Baeomorpha Brues, 1937, in Carpenter, Univ. Toronto Stud. geol. Ser. 40: 41, — Insecta, Hymenoptera. The objective replacement name is Hobapromea Watson nom. n.

HOBARTINA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 207.

Type-species: Scythrophanes amblyiodes Turner, 1931, Proc. Linn. Soc. N.S.W. 56: 330, by original designation. AUSTRALIA: Tasmania, Hobart.

HOLARCTIA Smith, 1938, Revision of the Genus Apantesis Walker. Univ. Illinois: 6.

Type-species: Arctia turbans Christoph, 1892, Horae Soc. ent. Ross. 26: 460, by original designation. [U.S.S.R.]: E. Siberia, S.W. of Irkutsk, Tunga-Alpen.

HOLOCHREA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 87, 268.

Type-species: *Holochrea xanthogramma* Hampson, 1900, *ibidem* 2: 269, fig.187, by original designation. [INDONESIA]: [Sulawesi], [Sangihe I.] Sangir.

HOLOCRASPEDON Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 14, 83.

Type-species: *Holocraspedon nigropunctum* Hampson, 1893, *ibidem* 9: 14, 84, pl.158, fig.4, by monotypy. [Sri Lanka] Ceylon: Nawala-pittia.

HOLOCRASPEDUM Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 91, 336.

An unjustified emendation of *Holocraspedon* Hampson, 1893.

A junior homonym of *Holocraspedum* Cragin, 1893, *Rep. geol. Surv. Texas* 4: 190, – Mollusca. The objective replacement name is *Holocraspedon* Hampson, 1893.

HOLOCRASPEDUM Hampson, 1900. See Holocraspedon Hampson, 1893.

HOLOMELINA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 95, 97.

Type-species: *Eubaphe aurantiaca* Hübner, [1831] 1825, *Zuträge Samml. exot. Schmett.* 3: 9, pl.[71], figs 411, 412, by subsequent designation by Hampson, 1901, *Cat. Lepid. Phalaenae Br. Mus.* 3: 188. [U.S.A.]: Pennsylvania.

Holomelina was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

HOLOPHAEA Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 408.

Type-species: *Holophaea caerulea* Druce, 1898 May, *ibidem* (7) 1: 408, by monotypy. ECUADOR.

The name *Holophaea* probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson, 1898 [December].

HOLOPHAEA Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xiv, 26, 265.

Type-species: *Ichoria lycone* Druce, 1884, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 1: 47, pl.6, fig.9, by original designation. Costa Rica: Irazu; Rio Sucio, Panama: Chiriqui Volcano.

A junior homonym of *Holophaea* Druce, 1898. There is no objective replacement name but the type-species of *Holophaea* Hampson, 1898 [December], is currently considered to be congeneric with the type-species of *Holophaea* Druce, 1898 May, the latter is available for use as a subjective replacement name.

‡HOLOPHOEA; Zerny, 1912, in Wagner, Lepid. Cat. 7: 155.
An incorrect subsequent spelling of Holophaea Hampson, 1898.

HOLOPHAEA Hampson, 1898 [December]. See Holophaea Druce, 1898 May.

‡HOLOPHOEA; misspelling. See Holophaea Druce, 1898.

‡HOMALOMERIA; misspelling. See Homoeomeria Wallengren, 1865.

HOMAROA Collenette, 1955, Naturaliste malgache 7: 176.

Type-species: *Leucoma brontonepha* Collenette, 1931, *Trans. ent. Soc. Lond.* **79**: 342, pl.13, fig.14, by original designation. MADAGASCAR: Antsianaka and Lake Alàotra.

HOMOCENTRIDIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 144.

Type-species: *Fentonia concentrica* Oberthür, 1911, *Etudes Lépid. comp.* **5** (1): 336, pl.83, fig.777, by original designation. [CHINA]: [Szechwan], Ta-tsien-Lou.

HOMOCHIRA Hampson, 1905, Ann. S. Afr. Mus. 3: 409.

Type-species: *Dasychira rendalli* Distant, 1897, *Ann. Mag. nat. Hist.* (6) **20**: 203, by original designation. [SOUTH AFRICA]: Transvaal, Barberton.

HOMOCHROA Spuler, 1906. See Omochroa Rambur, 1866.

HOMOEOCERA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.26. CTEN

Type-species: *Homoeocera crassa* Felder, 1874, *ibidem* 2 (Abt.2): pl.102, fig.26, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-105: 5.

HOMOEOGNATHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, figs 20, 21.

Type-species: *Homoeognatha aganais* Felder, 1874, *ibidem* **2** (Abt.2): pl.106, fig.21, by subsequent designation by Hampson, 1905, *Ann. S. Afr. Mus.* **3**: 414. [SOUTH AFRICA]: [Natal] Natalia, cited by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 5.

HOMOEOMERIA Wallengren, 1865, K. svenska Vetensk-Akad. Handl. (N.F.) 5 (4): 36.

Type-species: Leucoma flavicapilla Wallengren, 1860, Wien. ent. Monatschr. 4: 163, by monotypy. [South Africa]: Caffraria.

‡HOMALOMERIA; Strand, 1910, in Seitz, Gross-Schmett. Erde 2: 109.

An incorrect subsequent spelling of *Homoeomeria* Wallengren, 1865.

HOMONEURONIA Dyar, 1905, Proc. U.S. natn. Mus. 29: 174.

Type-species: *Homoneuronia modesta* Schaus, 1905, *ibidem* 29: 188, by original designation (in anticipation of Schaus's description of *modesta*). FRENCH GUIANA: Maroni River, St Jean.

Homoneuronia was available but was without an available included species between its date of publication 1905 October 3rd and that of its type-species 1905 October 11th by Schaus.

HOMOPSYCHE Butler, 1882, Ann. Mag. nat. Hist. (5) 10: 226.

Type-species: *Homopsyche nudarioides* Butler, 1882, *ibidem* (5) **10**: 226, by monotypy. [New Guinea]: [Bismarck Archipelago], New Britain.

H. nudarioides is a junior subjective synonym of Chamaita trichopteroides Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6:

See also Chamaita Walker, 1862.

HONDELLA Moore, [1883] 1882-3, Lepid. Ceylon 2: 144.

Type-species: *Ptilomacra juvenis* Walker, 1855, *List Specimens lepid. Insects Colln Br. Mus.* 5: 1099, by monotypy (but cited by Moore as ‡*Philomacra*, an incorrect subsequent spelling). [Sri Lanka] Ceylon.

HONVEDA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 154 (key), 169.

Type-species: *Pydna fasciata* Moore, 1879, in Hewitson & Moore, *Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson* (1): 66, by monotypy. INDIA: Darjeeling.

Honveda was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 71.

HOPLARCTIA Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: xii, 54. CTEN

Type-species: Ammalo nantana Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 32: 282, by original designation. PERU: [Nauta] Nanta.

‡HOPLISIA; misspelling. See Harpyia Ochsenheimer, 1810.

‡HOPLITES; misspelling. See Harpyia Ochsenheimer, 1810.

HOPLITINA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10:

Type-species: Stauropus notodontina Rothschild, 1917, Novit. zool. 24: 246, pl.5, fig.1, by original designation. [INDIA]: Assam, Khasia Hills.

The date of publication of the part of Seitz containing page 639 is cited on page 633 as 21.ii.1930 in the German edition; and as 17.v.1930 in the English edition.

HOPLITIS Hübner, [1819]. See Harpyia Ochsenheimer, 1810.

#HOPLOPSEUSTIS; misspelling. See Haplopseustis Meyrick, 1902. NOCTUIDAE

#HOPLOPSEUSTRIAS; misspelling. See Haplopseustis
Meyrick, 1902.
NOCTUIDAE

HOPPIANA Hering, 1925, in Hering & Hopp, Dt. ent. Z. Iris 39: 197.

Type-species: *Hoppiana aspasta* Hering, 1925, *ibidem* 39: 198, by original designation. Colombia: Choco, Rio Micay.

HORAIA Nakamura, 1973. See Pulia Kiriakoff, 1974.

HORAMA Hübner, [1819] 1816, Verz. bekannter Schmett.: 125.

Type-species: *Sphinx pretus* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 121, pl.175, figs E, F, by monotypy. WEST INDIES: [Virgin Is], St Thomas.

S. pretus is dated from the wrapper of the part and not from

the title-page of the volume, 1779.

MASTIGOCERA Harris, 1839, Am. J. Sci. 36: 315.

Type-species: *Mastigocera vespina* Harris, 1839, *ibidem* **36**: 315, by monotypy. West Indies: [Virgin Is], St Thomas.

M. vespina is a junior subjective synonym of Sphinx pretus Cramer, 1777, Uitlandsche Kapellen (Papillons exot.) 2: 121,

pl.175, figs E, F.

A junior homonym of *Mastigocera* Dejean, [1835], *Cat. Coléoptères* (Edn 2): 345, – Insecta, Coleoptera. There is no objective replacement name but the type-species of *Mastigocera* Harris, 1839, is a junior subjective synonym of the type-species of *Horama* Hübner, [1819]. The latter is therefore available for use as a subjective replacement name.

Neave, 1939, Nomencl. zool. 2: 160, cited Drucea Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 130 as a replacement name, but Kirby was not employing Drucea as such. Kirby proposed Drucea as a new genus for the concept Mastigocera Harris, 1839, sensu Boisduval, 1870, Considerations

Lépid. Guatemala: 81.

‡HORAMIA; Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 251.

An incorrect subsequent spelling of *Horama* Hübner, [1819].

HORAMELLA Draudt, 1915, in Seitz, Gross-Schmett. Erde 6: 145.

Type-species: *Horamella fassli* Draudt, 1915, *ibidem* 6: 145, pl.21, row i, by original designation. Colombia: eastern.

The date of publication of the part of Seitz containing page 145 is cited on that page as 30.iv.1915 in both the German and English editions.

‡HORAMIA; misspelling. See Horama Hübner, [1819].

HORISHACHIA Matsumura, 1929, Insecta matsum. 4: 39, 40.

Type-species: *Horishachia infusca* Matsumura, 1929, *ibidem* 4: 39, pl.1, fig.19, by original designation. TAIWAN: Horisha. *H. infusca* is a junior subjective synonym of *Phalera obscura* Wileman, 1910, *Entomologist* 43: 138.

HUNYADA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 151 (key), 161.

Type-species: *Pydna hunyada* Swinhoe, 1903, *Ann. Mag. nat. Hist.* (7) **11**: 504, by original designation. JAVA: southern.

Hunyada was established as a subgenus of Rosiora Kiriakoff, 1962; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 105.

HUPODONTA Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 475.

Type-species: *Hupodonta corticalis* Butler, 1877, *ibidem* (4) **20**: 475, by original designation. Japan: Yokohama.

H. corticalis is a subspecies of Anodonta pulcherrima Moore, 1866, Proc. zool. Soc. Lond. 1865: 814, pl.43, fig.4.

‡HAPODONTA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 429.

An incorrect subsequent spelling of Hupodonta Butler, 1877.

HYALAETHEA Butler, 1887, Ann. Mag. nat. Hist. (5) 19: 216.

Type-species: *Hyalaethea woodfordii* Butler, 1887, *ibidem* (5) **19**: 216, by monotypy. Solomon Islands: Alu.

HYALARCTIA Schaus, 1901 March, Ann. Mag. nat. Hist. (7) 7: 268.

Type-species: *Hyalarctia sericea* Schaus, 1901, *ibidem* (7) 7: 268, by monotypy. [Brazil]: Parana, Castro.

The name *Hyalarctia* probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson four months later.

HYALARCTIA Hampson, 1901 [July], Cat. Lepid. Phalaenae Br. Mus. 3: x, 5, 143.

Type-species: *Hyalarctia sericea* Schaus, 1901, by original designation.

A junior homonym of *Hyalarctia* Schaus, 1901. The objective replacement name is *Hyalarctia* Schaus, 1901.

HYALARCTIA Hampson, 1901 [July]. See Hyalarctia Schaus, 1901 March.

HYALEUCEREA Butler, 1875, Ann. Mag. nat. Hist. (4) 15: 399.

Type-species: Glaucopis erythrotelus Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 147, by original designation. [BRAZIL]: Para.

‡HYALEUCEREON; Schaus, 1901, Jl N.Y. ent. Soc. 9: 44.
An incorrect subsequent spelling of Hyaleucerea Butler, 1875.

‡HYALEUCEREON; misspelling. See Hyaleucerea Butler, 1875.

‡HYALOCA; misspelling. See Hyalocoa Hampson, 1901.

HYALOCOA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 5, 202.

Type-species: *Lithosia diaphana* Eversmann, 1848, *Bull. Soc. imp. Nat. Moscou* **21** (3): 212, by original designation. [U.S.S.R.]: Siberia, Irkutsk region.

‡HYALOCA; Bang-Haas, 1927, Horae macrolepid. Reg. palaearct. 1: 77.

An incorrect subsequent spelling of *Hyalocoa* Hampson, 1901.

HYALOMIS Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 429.

Type-species: Desmidocnemis thyria Druce, 1898, Ann. Mag. nat. Hist. (7) 1: 207, by subsequent designation by Collins, 1962, Ann. Mag. nat. Hist. (13) 5: 125. ECUADOR: Balsapamba.

Hampson, 1905, was not proposing *Hyalomis* as a replacement name for *Desmidocnemis* Möschler, 1872, *Stettin. ent. Ztg* 33: 346, – Lepid., Ctenuchidae, but was establishing a new genus for species placed incorrectly in *Desmidocnemis*, by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 363.

HYALOPERINA Aurivillius, 1904, Ark. Zool. 2 (4): 63.

LYMA

Type-species: *Hyaloperina nudiuscula* Aurivillius, 1904, *ibidem* **2** (4): 63, by monotypy. [Congo] Kongogebiet.

HYALOPIS Herrich-Schäffer, 1855. See Laemocharis Herrich-Schäffer, [1854].

HYALURGA Hübner, [1819] 1816, Verz. bekannter Schmett.:

Type-species: Phalaena fenestra Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 505, by monotypy (but cited as fenestrigera, an unjustified emendation by Hübner, [1819]). [WEST INDIES]

DITAXIS Boisduval, 1870, Considérations Lépid. Guatemala:

Type-species: Phalaena vinosa Drury, 1773, Illust. nat. Hist. exot. Insects 2: index to vol.1; 1770, ibidem 1: 47, pl.23, fig.4 [described and figured but without scientific name until 1773], by subsequent designation by Bryk, in Strand, Lepid. Cat. 45: 41. WEST INDIES: Antigua.

A junior homonym of Ditaxis M'Lachlan, 1867, J. Linn. Soc. (Zool.) 9: 262, – Neuroptera. There is no objective replacement name but the type-species of Ditaxis Boisduval, 1870, is currently considered to be congeneric with the type species of Hyalurga Hübner, [1819]. The latter is therefore available for use as a subjective replacement name.

CLETORNA Boisduval, 1870, Considérations Lépid. Guatemala: 86.

Type-species: Phalaena fenestra Linnaeus, 1758, by original designation (but cited as #fenestrata, an incorrect subsequent spelling).

A junior objective synonym of Hyalurga Hübner, [1819].

HYARIAS Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 184.

Type-species: Spilosoma metarhoda Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1695, by original designation. PHILIPPINES: Manila.

HYBOCAMPA Lederer, 1853. See Harpyia Ochsenheimer,

HYDA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 181.

Type-species: Glaucopis basilutea Walker, 1854, ibidem 1: 182, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 148. [Brazil]: Pernambuco.

Incorrect type-species: Glaucopis singularis Walker, 1854,

cited by Hampson, 1920, Novit. zool. 26: 253.

Hyda was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now the Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 178.

XANTHYDA Hampson, 1920, Novit. zool. 26: 253.

Type-species: Glaucopis basilutea Walker, 1854, by monotypy.

A junior objective synonym of Hyda Walker, 1854.

Xanthyda was established, unnecessarily, for the concept Hyda Walker, 1854, because an incorrect type-species was accepted and cited by Hampson, 1920. Xanthyda was linked by Hampson with its single included species by the entry 'p.178', which referred to Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 178.

HYDRUSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 255.

Type-species: Euchromia bicolor Walker, 1854, ibidem 1: 255, by monotypy. Australia.

Hydrusa was established as a subgenus of Euchromia Hübner, [1819].

HYELA Walker, 1854. See Phoenicoprocta Druce, 1898.

HYLAEORA Doubleday, [1849] 1848, Proc. zool. Soc. Lond. 1848: 117.

Type-species: Hylaeora eucalypti Doubleday, [1849] 1848, ibidem 1848: 118, Annulosa pl.5, by monotypy. Australia.

‡HYLEORA; Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1045.

An incorrect subsequent spelling of Hylaeora Doubleday, [1849].

HYLEMERA Butler, 1878, Ann. Mag. nat. Hist. (5) 2: 293. GEOMETRIDAE

Type-species: Hylemera tenuis Butler, 1878, ibidem (5) 2: 294, by original designation. MADAGASCAR: Fianarantsoa.

Hylemera was established in the Nyctemeridae, now Arctiidae; its type-species was placed in the Geometridae by Swinhoe, 1904, Trans. ent. Soc. Lond. 1904: 537.

‡HYLEORA; misspelling. See Hylaeora Doubleday, [1849].

HYPAGOPTERA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 83, 179.

Type-species: Hypagoptera rufeola Hampson, 1900, ibidem 2: 179, fig. 95, by original designation. [South Africa]: Natal, Eastcourt, Cape Province, Transkei.

‡HYPOGOPTERA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 422.

An incorrect subsequent spelling of Hypagoptera Hampson,

HYPAMBADRA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 385.

Type-species: Hypambadra speculigera Kiriakoff, 1974, ibidem 17: 386, text-fig.9, pl.1, fig.6, by original designation. SUMATRA: northern, 30 km S.W. of Siantar.

‡HYPANTRIA; misspelling. See Hyphantria Harris, 1841.

HYPAREVA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 90, 191.

Type-species: Hypareva pogonoda Hampson, 1900, ibidem 2: 192, fig.112, by original designation. MEXICO: Teapa, GUA-TEMALA, PANAMA, [FRENCH GUIANA]: Cayenne.

HYPARPAX Hübner, [1825] 1806, Samml. exot. Schmett. 2: pl.[168].

Type-species: Phalaena aurora Smith, 1797, in Smith & Abbot, Nat. Hist. rarer lepid. Insects Georgia 2: 173, pl.87, by monotypy. [U.S.A.]: Georgia.

See also Sangata Walker, 1860.

HYPASURA Daniel, 1952, Bonn. zool. Beitr. 3: 89. Type-species: Hypasura hoenei Daniel, 1952, ibidem 3: 89, by original designation (but cited as \$h\overline{o}nei, an incorrect original spelling). CHINA: Hunan, Hoeng Shan.

HYPATIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 147.

Type-species: Hysia delecta Butler, 1876, J. Linn. Soc. (Zool.) 12: 396, by original designation. [BRAZIL]: Para.

H. delecta was established for the concept Euchromia astyoche Geyer, 1832, sensu Walker, 1854; see below under

Hypatia was established as the objective replacement name for Hysia Walker, 1854, which was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae. Hypatia and Hysia were retained in the Zygaenidae by Kirby, 1892, ibidem 1: 147; they were placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 266.

HYSIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 172.

Type-species: Hysia delecta Butler, 1876 (but included by Walker as Glaucopis astyoche (Geyer) a misidentification; see below), by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 147 (for Hypatia Kirby, 1892).

H. delecta was established for the concept Euchromia astyoche Geyer, 1832, in Hübner, Zuträge Samml. exot. Schmett. 4: 41, sensu Walker, 1854, ibidem 1: 173. This is a case of a misidentified type-species which under he Code, Article 70(a), should be referred to the Commission.

Hysia was established as a subgenus of Glaucopis Fabricius,

1807.

A junior homonym of *Hysia* Mulsant, 1850, *Annls Soc. Agric. Lyon* (2) **2**: 47, – Coleoptera. The objective replacement name is *Hypatia* Kirby, 1892.

HYPERAESCHRA Butler, 1880, Ann. Mag. nat. Hist. (5) 6: NOTO

Type-species: *Hyperaeschra pallida* Butler, 1880, *ibidem* (5) 6: 65, by original designation. [INDIA]: Darjeeling.

HYPERAESCHRELLA Strand, 1916, Arch. Naturgesch. 81
(A) 12: 154.

Type-species: *Hyperaeschra kosemponica* Strand, 1916, *ibidem* **81**(A) 12: 153, by original designation. Talwan: Kosempo.

Hyperaeschrella was established as a subgenus of Hyperaeschra Butler, 1880.

HYPERANDRA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 3, 69.

Type-species: Creatonotus appendiculatus Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: wrapper, pl.82, figs 475, 476, by monotypy. [BRAZIL]: Rio [de Janeiro].

C. appendiculatus was established in Creatonotus Agassiz, 1847, an unjustified emendation of Creatonotos Hübner, [1819].

‡HYPERANTHINA; misspelling. See Hyphantria Harris, 1841.

HYPERBOREA Grum-Grshimailo, 1900, Ezheg. zool. Muz.

Type-species: *Hyperborea czekanowskii* Grum-Grshimailo, 1900, *ibidem* 4: 464, by monotypy. [U.S.S.R.]: Siberia, R. Tunguska valley.

‡HYPERBOREAE; Sheljuzhko, 1918, Neue Beitr. syst. Insektenk. 1: 104.

An incorrect subsequent spelling of *Hyperborea* Grum-Grshimailo, 1900.

‡HYPERBOREAE; misspelling. See Hyperborea Grum-Grshimailo, 1900.

‡HYPERCOMPA; misspelling. See Hypercompe Hübner, [1819].

‡HYPERCOMPE Hübner, [1806]. See Arctia Schrank, 1802.

‡HYPERCOMPE Hübner, 1808. See Cycnia Hübner, 1818.

‡HYPERCOMPE Hübner, 1818. See Hypercompe Hübner, [1819].

HYPERCOMPE Hübner, [1819], Samml. exot. Schmett. 1: pl.[191].

Type-species: *Phalaena icasia* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 130, 149, pl.181, fig.E (but included by Hübner as *Hypercompe eridane* (Cramer) a misidentification; see below), by subsequent designation by International Commission on Zoological Nomenclature, 1966, *Bull. zool. Nom.* 23 (Opinion 789): 213. SURINAM.

P. icasia is dated from the wrapper of the part and not from the title-page of the volume, 1779.

The species figured by Hübner, [1819]: pl.[191] was misidentified and was not *Phalaena eridanus* Cramer, 1775, *ibidem* 1: 107, but was *Phalaena icasia* Cramer, 1777, *fide* Seitz, 1919, *Gross-Schmett. Erde* 6: 319. In accordance with the *Code*, Article 70(a), the Commission designated *Phalaena icasia* as type-species.

Hypercompe Hübner, [1819], was placed on the Official List of Generic Names in Zoology: Name No.1738 by the International Commission on Zoological Nomenclature, 1966, ibidem 23 (Opinion 789): 213.

‡HYPERCOMPE Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 7, 33.

Not nomenclaturally available as a genus-group name. Hübner used Hypercompe, on both pages, in a suprageneric sense for one of the names of his tribes (Namen der Stämme), listed on pages [33], [34]. Hübner listed his genus-group names on pages [35], [36].

‡HYPERCOMPE; Billberg, 1820, Enumeratio Insect. Mus. G.J. Billberg: 91.

An incorrect subsequent spelling of *Hypercompe* Hübner, [1819].

‡Hypercompa was also used by Rambur, [1866], Cat. syst. Lépid. Andalousie (2): 230; and by Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 55, 67.

HYPERMAEPHA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 85, 351.

Type-species: Cisthene sanguinea Butler, 1878, Trans. ent. Soc. Lond. 1878: 53, by original designation. [BRAZIL]: [Amazonas], Rio Jutahi.

HYPERPHARA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xix, 31, 468.

Type-species: *Melanochroia junctura* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 211, by original designation. [COLOMBIA]: Bogota.

HYPERTHAEMA Schaus, 1901 March, Ann. Mag. nat. Hist. (7) 7: 265.

Type-species: *Hyperthaema albipuncta* Schaus, 1901, *ibidem* (7) 7: 265, by subsequent designation by Collins, 1962, *Ann. Mag. nat. Hist.* (13) 5: 122. BRAZIL: São Paulo.

The name *Hyperthaema* probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson, 1901 [July].

HYPERTHAEMA Hampson, 1901 [July], Cat. Lepid. Phalaenae Br. Mus. 3: viii, 6, 88.

Type-species: Halysidota sanguineata Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 309 (but cited as ‡Halesidota, an incorrect subsequent spelling), by original designation (but cited as ‡Halesidota). [COLOMBIA]: Bogota. A junior homonym of Hyperthaema Schaus, 1901 March.

A junior homonym of *Hyperthaema* Schaus, 1901 March. There is no objective replacement name but the type-species of *Hyperthaema* Hampson, 1901 [July] is currently considered to be congeneric with the type-species of *Hyperthaema* Schaus, 1901 March. The latter is therefore available for use as a subjective replacement name.

‡HYPOTHAEMA; Jörgensen, 1935, An. Mus. argent. Cienc. nat. 38: 109.

An incorrect subsequent spelling of *Hyperthaema* Schaus, 1901 March.

HYPERTHAEMA Hampson, 1901 [July]. See Hyperthaema Schaus, 1901 March.

HYPERTHAGYLLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 92, 192.

Type-species: Agylla mira Butler, 1878, Trans. ent. Soc.

Lond. 1878: 58, pl.3, fig.2, by original designation. [BRAZIL]: [Amazonas], Rio Javary and Rio Jutahi.

HYPEUGOA Leech, 1899, Trans. ent. Soc. Lond. 1899: 189.

Type-species: *Hypeugoa flavogrisea* Leech, 1899, *ibidem* **1899**: 190, by original designation. CHINA: [Szechwan], Tatsien-lu.

HYPHANTRIA Harris, 1841, Rep. Insects Mass. injurious to Vegn: 255.

Type-species: *Hyphantria textor* Harris, 1841, *ibidem*: 255, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 234. U.S.A.: Massachusetts.

H. textor is a junior subjective synonym of Phalaena cunea Drury, 1773, Illust. nat. Hist. exot. Insects 2: index to vol. 1; 1770, ibidem 1: 36, pl.18, fig.4 [described and figured but without name until 1773].

‡HYPANTRIA; Clemens, 1861, Proc. Acad. nat. Sci. Philad. 1860: 530.

An incorrect subsequent spelling of Hyphantria Harris, 1841.

‡HYPERANTHINA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 421.

An incorrect subsequent spelling of Hyphantria Harris, 1841.

HYPHILARA Kirby, 1897. See *Hyphilare* Hübner, [1821]. NOCTUIDAE

HYPHILARE Hübner, [1821] 1816, Verz. bekannter Schmett.: 239.

Type-species: Noctua albipuncta [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 84, by subsequent designation by Hampson, 1905, Cat. Lepid. Phalaenae Br. Mus. 5: 436.

Further information in Nye, 1975, Generic Names Moths World 1: 252.

HYPHILARA Kirby, 1897, Allen's Naturalist's Libr. (Handbk Order Lepid.) 5: 15.

An unjustified emendation of *Hyphilare* Hübner, [1821]. *Hyphilara* in the Noctuidae, Hadeninae is cited here as it was not included in Nye, 1975, *Generic Names Moths World* 1.

HYPHORAEA Agassiz, 1847. See Hyphoraia Hübner, [1819].

HYPHORAIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 182.

Type-species: *Phalaena aulica* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 505, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 261. Europe.

HYPHORAEA Agassiz, 1847, Nomencl. zool. (Index univl.): 190.

An unjustified emendation of *Hyphoraia* Hübner, [1819].

The 'Index universalis' was published as Fascicle 12 of the *Nomenclator zoologicus*; *Hyphoraea* is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

HYPIDALIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 3, 70.

Type-species: *Idalus enervis* Schaus, 1894, *Proc. zool. Soc. Lond.* 1894: 229, by original designation. [BRAZIL]: Parana, Castro.

HYPIDOTA Schaus, 1904, Trans. Am. ent. Soc. 30: 137.

Type-species: *Hypidota neurias* Schaus, 1904, *ibidem* 30: 137, by monotypy. [Brazil]: Parana, Castro.

HYPOCHARIS Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xv, 26, 271.

Type-species: *Laemacharis clusia* Druce, 1897, *Ann. Mag. nat. Hist.* (6) **20**: 303, by original designation. [BRAZIL]: Amazon region, 'Maranham'.

HYPOCLADIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xvii, 31, 362.

Type-species: *Dycladia militaris* Butler, 1877, *Illust. typical Specimens Lepid. Heterocera Colln Br. Mus.* 1: ix, 33, pl.16, fig.1, by original designation. [BRAZIL]: [Para], Santarem.

HYPOCRISIAS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 7, 216.

Type-species: *Purius punctatus* Druce, 1884, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 1: 100, by original designation. GUATEMALA: Las Mercedes.

HYPOCRITA Hübner, [1807], Samml. exot. Schmett. 1: pl.[187].

Type-species: *Hypocrita dichroa* Hübner, [1807], *ibidem* 1: pl.[187], by monotypy. Type-locality: not stated, [tropical AMERICA].

H. dichroa is a junior subjective synonym of Phalaena bicolora Sulzer, 1776, Geschichte Insecten: 161, pl.22, fig.6.

See also Esthema Hübner, [1820].

‡HIPOCRITA; Hübner, [1810], Samml. exot. Schmett. 1: pl.[189].

An incorrect subsequent spelling of *Hypocrita* Hübner, [1807].

‡Hipocrita was placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1859.

‡HYPOCRITA Hübner, 1808. See Metastatia Butler, 1876. CTEN

‡HYPOGOPTERA; misspelling. See Hypagoptera Hampson, 1900.

‡HYPOGYMA; misspelling. See Lymantria Hübner, [1819].

HYPOGYMNA Billberg, 1820. See Lymantria Hübner, [1819].

‡HYPOGYMNIA; misspelling. See Lymantria Hübner, 1819.

‡HYPOLACHNA; misspelling. See Hypolochma Felder, 1861.

HYPOLOCHMA Felder, 1861, Sher. Akad. Wiss. Wien 43 (Abt.1): 40.

NOCTUIDAE

Type-species: *Hypolochma sericea* Felder, 1861, *ibidem* 43 (Abt.1): 41, by monotypy. Type(s) &, [Indonesia]: [Moluccas], [Ambon I.] Amboina, (BMNH).

Hypolochma was established in the Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 923, but has not been used by later authors; it is here transferred to the Noctuidae, Sarrothripinae, near Barasa Walker, 1862, on the advice of, and with acknowledgement to our colleague Mr A.H. Hayes.

‡HYPOLACHNA; Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 441.

An incorrect subsequent spelling of *Hypolochma* Felder, 1861.

HYPOMOLIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xv, 3, 414.

Type-species: *Idalus evippus* Druce, 1898, *Ann. Mag. nat. Hist.* (7) 1: 212, by original designation. BOLIVIA.

HYPONERITA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xv, 3, 440.

Type-species: *Idalus lavinia* Druce, 1890, *Proc. zool. Soc. Lond.* **1890**: 496, pl.42, fig.6, by original designation. [GUYANA] British Guiana.

HYPOPHIALA Janse, 1920, Ann. Transv. Mus. 7: 156 (key), 192.

Type-species: *Hypophiala melanogramma* Janse, 1920, *ibidem* 7: 193, pl.1, figs 26, 27, pl.4, figs 20-24, pl.14, fig.3, by original designation. [ZIMBABWE-]RHODESIA: Salisbury.

H. melanogramma is a junior subjective synonym of Haplozana nigrolineata Aurivillius, 1901, Ent. Tidskr. 22: 122, taxt. fig. 23

C

See also Haplozana Aurivillius, 1901.

HYPOPREPIA Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 21.

Type-species: *Hypoprepia fucosa* Hübner, [1831] 1825, *ibidem* 3: 21, pl.[81], figs 471, 472, by monotypy. [U.S.A.]: Georgia.

HYPOSHADA Hampson, 1909, Ann. Mag. nat. Hist. (8) 4: 352.

Type-species: Scoliacma pellopis Bethune-Baker, 1908, Novit. zool. 15: 193, by original designation (but cited as ‡pelopis, an incorrect subsequent spelling). New Guinea: [Papua New Guinea], Owgarra and Biagi.

HYPOSICCIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 90, 406.

Type-species: Aemene amnaea Swinhoe, 1894, Ann. Mag. nat. Hist. (6) 14: 437, by original designation. [INDIA]: [Assam], Shillong.

HYPOSTAURELLA Kiriakoff, 1967, Biologia gabon. 3: 335.

Type-species: *Hypostaurella arcuata* Kiriakoff, 1967, *ibidem* 3: 335, fig.23, by original designation. GABON: Makokou.

HYPOSTAUROPUS Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 397.

Type-species: *Quadricalcarifera saitonis* Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19 (1): 12, pl.1, fig.10, by original designation. TAIWAN: Horisha, Baibara.

‡HYPOTHAEMA; misspelling. See Hyperthaema Schaus, 1901.

HYPOTUERTA Kiriakoff, 1976, Tierreich 98: 109.

Type-species: *Tuerta transiens* Hampson, 1901, *Cat. Lepid. Phalaenae Br. Mus.* 3: 624, pl.53, fig.15, by original designation. Holotype 9, [NIGERIA]: Old Calabar, (BMNH).

Hypotuerta in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

HYPSA Hübner, [1819] 1816, Verz. bekannter Schmett.: 172.

Type-species: *Phalaena silvandra* Stoll, 1781, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 155, 252, pl.369, fig.D, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 388. [INDIA]: [S.], Coramandel.

P. silvandra is dated from the wrapper of the part and not

from the title-page of the volume, 1782.

P. silvandra is a junior subjective synonym of Phalaena heliconia Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 511.

ASPA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 452.

Type-species: *Phalaena heliconia* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 511, by monotypy. INDIA.

A junior homonym of Aspa Adams & Adams, [1853] 1858, Genera recent Mollusca 1: 106, — Mollusca. There is no objective replacement name but P. heliconia is the senior subjective synonym of P. silvandra Stoll, the type-species of Hypsa Hübner, [1819]. The latter is therefore available for use as a subjective replacement name.

Aspa was established as a subgenus of Hypsa Hübner, [1819].

HYPSOIDES Butler, 1882, Cistula ent. 3: 1. THAU
Type-species: Hypsoides bipars Butler, 1882, ibidem 3: 2, by

monotypy. MADAGASCAR: Forests of Ancaya.

Hypsoides was established in the Chalcosiidae, now Zygaenidae, Chalcosiinae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 578; and was placed in the Thaumetopoeidae by Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 397.

HYRMINA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 331.

Type-species: *Dioptis areolata* Walker, 1854, *ibidem* 2: 332, by PRESENT DESIGNATION. BRAZIL.

Hyrmina was established as a subgenus of Dioptis Hübner, 1818.

‡HERMINA; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1653.

An incorrect subsequent spelling of Hyrmina Walker, 1854.

HYSIA Walker, 1854. See Hypatia Kirby, 1892. CTEN

HYSIBADA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 497.

Type-species: *Hysibada varipes* Walker, 1865, *ibidem* 32: 498, by monotypy. [South Africa]: Natal.

Hysibada was established in the Limacodidae; it was placed in the Bombycidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 716; and was placed in the Lymantriadae, now Lymantriidae, by Hampson, 1905, Ann. S. Afr. Mus. 3: 411.

See also Woerdenia Snellen, 1872.

IANASSA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1101.

Type-species: *Ianassa lignicolor* Walker, 1855, *ibidem* 5: 1101, by monotypy. Type-locality: not stated, [U.S.A.]: [Atlantic States].

See also Xylinodes Packard, 1864.

PHYA Druce, 1887, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 242.

Proposed unnecessarily as an objective replacement name for *Ianassa* Walker, 1855, which Druce believed to be preoccupied by *Janassa* Münster, 1839, *Beitr. Petrefacten-Kunde* 1: 46, 114, – Pisces.

ICAMBOSIDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 400.

Type-species: *Icambosida nigrifrons* Walker, 1865, *ibidem* 32: 401, by monotypy. [INDIA] Hindostan.

‡ICHARIA; misspelling. See Ichoria Butler, 1876. CTEN

ICHORIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 370. CTEN Type-species: Euchromia quadrigutta Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 245, by original designation. MEXICO.

‡ICHARIA; Rothschild, 1911, Novit. zool. 18: 36.
An incorrect subsequent spelling of Ichoria Butler, 1876.

ICHTHYOSOMA Felder, 1874. See Trichomaplata Wing, [1850].

ICHTHYURA Hübner, [1819] 1816, Verz. bekannter Schmett.: 162.

Type-species: *Phalaena anastomosis* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 506, by subsequent designation by Hampson, [1893] 1892, *Fauna Br. India* (Moths) 1: 172. Type-LOCALITY: not stated, [EUROPE].

ICTA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 922.

Type-species: *Icta fulviceps* Walker, 1855, *ibidem* **4**: 922, by monotypy. [Australia] New Holland.

IDALIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 149.

Type-species: *Phalaena regina* Cramer, 1780, *Uitlandsche Kapellen (Papillons exot.)* 3: 142, 176, pl.272, fig.A, by monotypy. SURINAM.

P. regina is dated from the wrapper of the part and not from the title-page of the volume, 1782.

IDALUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 645.

Type-species: *Phalaena admirabilis* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* **2**: 11, [147], pl.103, fig.G, by monotypy. Surinam.

P. admirabilis is dated from the wrapper of the part and not from the title-page of the volume, 1779.

EMPUSA Hübner, [1819] 1816, Verz. bekannter Schmett.: 170.

Type-species: *Phalaena admirabilis* Cramer, 1777, by monotypy.

A junior homonym of *Empusa* Illiger, 1798, *in* Kugelann, *Verz. Käfer Preussens*: 499, – Insecta, Orthoptera. The objective replacement name is *Idalus* Walker, 1855.

IDIOCTETUS Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.88, fig.504.

Type-species: *Idioctetus leucanioides* Herrich-Schäffer, [1856] 1850-1858, *ibidem* 1 (1): wrapper, pl.88, fig.504, by monotypy. America: [tropical] Am. aequin.

IDOPTERUM Hampson, 1894, Fauna Br. India (Moths) 2: viii, 42, 103.

Type-species: *Idopterum ovale* Hampson, 1894, *ibidem* 2: 103, fig.69, by original designation. Burma: Bernardmyo.

ILEMA Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 341. LYMA

Type-species: *Melia costalis* Walker, 1855, *List Specimens lepid. Insects Colln Br. Mus.* 4: 808, by monotypy (of *Melia* Walker, 1855). JAVA.

Ilema was proposed as the objective replacement name for *Melia* Walker, 1855.

MELIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 808.

Type-species: *Melia costalis* Walker, 1855, by monotypy. A junior homonym of *Melia* Bosc, 1813, *in* Risso, *Nouv. Bull. sci. Soc. philomath. Paris* 3: 233, — Crustacea. The objective replacement name is *Ilema* Moore, [1860].

ILEMA Hampson, 1900. See Eilema Hübner, [1819]. LITH

ILEMODES Hampson, 1900, Ann. S. Afr. Mus. 2: 53. ARCT Type-species: Ilemodes heterogyna Hampson, 1900, ibidem
2: 53, by original designation. South Africa: Natal, and [Cape Province] Cape Colony.

ILIPA Walker, 1854, List Specimens lepid. Insects Colln Br.
Mus. 1: 166. CTEN

Type-species: Glaucopis braconoides Walker, 1854, ibidem

1: 166, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 143. HONDURAS.

Ilipa was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 225.

ILLICE Walker, 1859, List Specimens lepid. Insects Colln Br. Mus. 19: 1019.

Type-species: *Illice batialis* Walker, 1859, *ibidem* **19**: 1019, by monotypy. [Brazil]: Rio [de] Janeiro.

ILLIPULA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 237.

Type-species: Sesia melanochloros Sepp, [1848], Surinaamsche Vlinders (2): 145, pl.69, by monotypy. SURINAM.

S. melanochloros is a junior subjective synonym of Zygaena eriphia Fabricius, [1777], Genera Insect.: 276.

Illipula was established as a subgenus of Euchromia Hübner, [1819].

See also Antichloris Hübner, 1818, and Copaena Herrich-Schäffer, [1855].

IMAIDA Toxopeus, 1948, Treubia 19: 448.
Type-species: Imaida sinuata Toxopeus, 1948, ibidem 19: 449, figs 1, 7b, by original designation. Borneo: [Kalimantan], Samarinda.

IMAUS Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 54. LYMA
Type-species: Lymantia munda Walker, 1855, List Species

Type-species: Lymantria munda Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 875, by original designation. [Bangladesh]: [E. Bengal], [Sylhet] Silhet.

INDUBA Kiriakoff, 1975, Arnoldia, (Rhodesia) 7 (15): 7.

Type-species: *Induba nigrescens* Kiriakoff, 1975, *ibidem* 7 (15): 8, fig.6, by original designation. [ZIMBABWE-]RHODESIA: Matetsi, Delaware ranch.

INNISCA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 153 (key), 176.

Type-species: *Pydna eupatagia* Hampson, [1893] 1892, *Fauna Br. India* (Moths) 1: 141, by original designation. [INDIA]: Sikkim.

Innisca was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 78.

INOPSIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.6.

Type-species: *Inopsis catoxantha* Felder, 1874, *ibidem* 2 (Abt.2): pl.106, fig.6, by monotypy. Mexico, cited by Felder, 1875, *ibidem* 2 (Abt.2): Erklärung to pls 75-107: 5 (as ‡*Jnopsis*, an incorrect subsequent spelling).

Felder, 1874, in his script legends to his plates, did not distinguish between the capital letters I and J; the spelling *Inopsis* has been adopted by authors.

‡JNOPSIS; Felder, 1875, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): Erklärung to pls 75-107: 5. An incorrect subsequent spelling of Inopsis Felder, 1874.

INOUELLA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 193.

Type-species: *Lophopteryx umbrosa* Leech, 1898, *Trans. ent. Soc. Lond.* **1898**: 313, by original designation. China: western, Ni-tou.

INOUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1095.

Type-species: Inous nigripalpis Walker, 1855, ibidem 5: 1096, by monotypy. South Africa.

Inous was established in the Notodontidae; it was placed in the Limacodidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 542.

IONTHAS Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxvii, 777.

Type-species: Ionthas ataracta Hampson, 1914, ibidem 1: 777, fig.254, by original designation. [AUSTRALIA]: Queensland, Warwick.

The name Ionthas and that of its type-species were attributed to Turner by Hampson.

‡JONTHAS; Strand, 1920, in Wagner, Lepid. Cat. 24: 802. An incorrect subsequent spelling of Ionthas Hampson, 1914.

IOSTAURA Kiriakoff, 1970, Ann. Transv. Mus. 26 (9): 187.

Type-species: Iostaura jansei Kiriakoff, 1970, ibidem 26 (9): 188, fig.6, by original designation. South Africa: Natal, St Lucia.

IPHIGENIELLA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 47.

Type-species: Iphigeniella prasina Kiriakoff, 1962, ibidem (8vo Sci. zool.) 114: 48, fig.61, by original designation. [ZAIRE]: east Ituri valley, 20 miles south of Irumu.

IRA Neumoegen, 1890. See Cubira Watson nom. n. **CTEN**

IRANADA Wiltshire, 1977, Scientific Results of the Oman Flora and Fauna Survey 1975 (Lepid.): 174.

Type-species: Phothedes secunda Ershov, 1874, in Fedchenko, Reise in Turkestan 2 (5) 3: 52, pl.3, fig.49, by original designation. Lectotype &, [U.S.S.R.]: Turkestan, Kisil-Kum desert, (ZI, Leningrad) designated by Wiltshire (in press).

Iranada in the Noctuidae, Ophiderinae, is cited here as it

was not included in Generic Names Moths World 1.

IRIDOPLITIS Kiriakoff, 1955, Revue Zool. Bot. afr. 52: 338.

Type-species: Iridoplitis iridescens Kiriakoff, 1955, ibidem 52: 339, fig.15, pl.2, fig.15, by original designation. [ZAMBIA] N.W. Rhodesia: Mwengwa.

IROPOCA Turner, 1904, Trans. ent. Soc. Lond. 1904: 477.

Type-species: Teara rotundata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 851, by monotypy. [Aus-TRALIA]: [Tasmania] Van Dieman's Land.

ISANTHRENE Hübner, [1819] 1816, Verz. bekannter Schmett.: 125.

Type-species: Glaucopis incendiaria Hübner, [1809] 1806, Samml. exot. Schmett. 1: pl.[166], by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 132. Type-Locality: not stated, [South America].

\$SPHECOMORPHA Hübner, [1806], Tentamen determina-

tionis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) Opinion 97: 19; also ibidem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140.

Only included species Sphecomorpha incendiaria Hübner, 1806, ibidem: [1], later established as Glaucopis incendiaria Hübner, [1809].

‡ISARES Herrich-Schäffer, [1858]. See Macrobrochis Herrich-Schäffer, 1855. ARCT

ISCHNARCTIA Bartel, 1903, Dt. ent. Z. Iris 16: 206. ARCT

Type-species: Ischnarctia brunnescens Bartel, 1903, ibidem 16: 207, by monotypy. [Tanzania]: [Tanganyika], Unyika.

ISCHNOCAMPA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.22. ARCT

Type-species: Ischnocampa sordida Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.22, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 6 (as ‡ Jschnocampa, an incorrect subsequent spelling).

Felder, 1874, in his script legends to his plates, did not distinguish between the capital letters I and J; the spelling *Ischnocampa* has been adopted by authors.

‡JSCHNOCAMPA; Felder, 1875, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): Erklärung to pls

An incorrect subsequent spelling of Ischnocampa Felder, 1874.

‡ISCHNOGNATHA Boisduval, 1870. See Ischnognatha Felder, 1874. ARCT

ISCHNOGNATHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.2.

Type-species: Ischnognatha semiopalina Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.2, by monotypy. [French Guiana] Guyana gall., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6 (as ‡ Jschnognatha, an incorrect subsequent spelling).

Felder, 1874, in his script legends to his plates, did not distinguish between the capital letters I and J; the spelling

Ischnognatha has been adopted by authors.

‡ISCHNOGNATHA Boisduval, 1870, Considérations Lépid. Guatemala: 99 (but attributed to Felder).

A nomen nudum, first published in the synonymy of Trichromia Hübner, [1819] (but attributed to Herrich-Schäffer, an incorrect authorship), and unavailable under the Code,

The only originally included species, ‡Trichromia semiopalina Felder, a nomen nudum, was later established as Ischnognatha semiopalina Felder, 1874. Boisduval presumably had access to Felder's plates and legends (to which Boisduval refers) prior to their publication by Felder in 1874.

‡JSCHNOGNATHA; Felder, 1875, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): Erklärung to pls 75-107: 6.

An incorrect subsequent spelling of Ischnognatha Felder, 1874.

ISIA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1698.

Type-species: Isia intricata Walker, 1856, ibidem 7: 1698, by monotypy. Brazil.

ISINE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 545.

Type-species: Isine trigutta Walker, 1854, ibidem 2: 545, by monotypy. Sierra Leone.

Isine was established in the 'Lithosiidae'; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 429; and was transferred back to the 'Lithosiidae' by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497

ISIONDA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1657.

Type-species: Isionda pyraloides Walker, 1856, ibidem 7: 1658, a junior secondary homonym, see below, by monotypy. [Brazil]: Tapayos.

Isionda pyraloides Walker, 1856, is currently included in

Myonia Walker, 1854, where it is a junior secondary homonym of Chrysauge pyraloides Walker, 1854, ibidem 2: 376; the objective replacement name, Myonia assimilis Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 506, is the valid name for the type-species of Isionda Walker, 1856.

Under the *Code*, Article 59(b)(i), a junior secondary homonym rejected before 1961 is permanently rejected and cannot be restored unless the use of the replacement name is contrary

to existing usage.

Isionda was established in the Pericopidae, now Arctiidae; it was placed in the Cyllopodidae, now Geometridae, Sterrhinae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 409; and was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 399.

ISOCTENIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.99, fig.15.

Type-species: *Isoctenia quadristrigata* Felder, 1874, *ibidem* 2 (Abt.2): pl.99, fig.15, by monotypy. [French Guiana] Guyana gall., cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 6.

I. quadristrigata is a junior subjective synonym of Turuenna dirphioides Walker, 1865, the type-species of Turuenna Walker, 1865.

Felder, 1874, in his script legends to his plates, did not distinguish between the capital letters I and J; the spelling *Isoctenia* has been adopted by authors.

Isoctenia when established was not placed in a family; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 834; and placed in the Lymantriidae by Bryk, 1934, in Strand, Lepid. Cat. 62: 323.

See also Turuenna Walker, 1865.

‡JSOCTENIA; Felder, 1875, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): Erklärung to pls 75-107: 6.

An incorrect subsequent spelling of *Isoctenia* Felder, 1874.

ISORROPUS Butler, 1880, Ann. Mag. nat. Hist. (5) 5: 342.
LITH

Type-species: *Isorropus tricolor* Butler, 1880, *ibidem* (5) 5: 343, by original designation. MADAGASCAR.

ISOSTOLA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.103, fig.15.

PERI

Type-species: *Isostola rhodobroncha* Felder, 1874, *ibidem* 2 (Abt.2): pl.103, fig.15, by monotypy. [Brazil]: Amazon R., cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 6 (as ‡*Jsostola*, an incorrect subsequent spelling).

Felder, 1874, in his script legends to his plates, did not distinguish between the capital letters I and J; the spelling *Isostola* has been adopted by authors.

‡JSOSTOLA; Felder, 1875, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): Erklärung to pls 75-107: 6.

An incorrect subsequent spelling of Isostola Felder, 1874.

ISOSTYLA Prout, 1918, Novit. zool. 25: 397 (key), 415.

Type-species: *Dioptis erycinoides* Felder, 1874, *in* Felder & Rogenhofer, *Reise öst. Fregatte Novara* (Zool.) 2 (Abt.2): pl.105, fig.4, by original designation. PANAMA: cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 4.

ITALAVIANA Kiriakoff, 1960, Revue fr. Ent. 27: 181. NOTO Type-species: Italaviana griveaudi Kiriakoff, 1960, ibidem
27: 182, fig.8, by original designation. MADAGASCAR: Italaviana, 6 km northwest of Fanovana.

ITYCA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1889.

Type-species: *Ityca humeralis* Walker, 1866, *ibidem* 35: 1890, by monotypy. [MALAYSIA]: [Malaya] Malacca.

I. humeralis is a junior subjective synonym of Teulisna chiloides Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 109.

See also Corcura Walker, 1862, and Diastrophia Felder, 1874.

IVELA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 388.

Type-species: Leucoma auripes Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 402, by original designation. JAPAN: Yokohama.

IXYLASIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 410. CTEN Type-species: Aclytia trogonoides Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 101, by original designation. Brazil.

JABAINA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 230.

Type-species: *Dasychira ania* Hering, 1926, *in* Seitz, *Gross-Schmett. Erde* 14: 177, pl.27, row c, by original designation. MADAGASCAR: Diego Suarez.

JACKSONIA Collenette, 1937. See Jacksoniana Nye nom.

JACKSONIANA Nye nom. n. for Jacksonia Collenette, 1937.

Type-species: Jacksonia striata Collenette, 1937, Ann. Mag. nat. Hist. (10) 20: 615, pl.13, fig.4, by original designation (for Jacksonia Collenette, 1937). UGANDA: Katera, Mababigambo.

JACKSONIA Collenette, 1937, Ann. Mag. nat. Hist. (10) 20: 614.

Type-species: *Jacksonia striata* Collenette, 1937, by original designation.

A junior homonym of *Jacksonia* Theobald, 1923, *Scott. Nat.* **1923**: 19, – Hemiptera. The objective replacement name is *Jacksoniana* Nye, nom. n.

JANTHINISCA Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 170.

Type-species: *Janthinisca lilacea* Kiriakoff, 1959, *ibidem* **95**: 171, fig.17, by original designation. [ZAIRE] Congo Belge: [Lubumbashi] Elisabethville.

‡JCHTHYOSOMA; misspelling. See Trichomaplata Wing, [1850].

‡JNOPSIS; misspelling. See Inopsis Felder, 1874. LITH

‡JONTHAS; misspelling. See Ionthas Hampson, 1914.

JORGENSENIA Schaus, 1924. See Geriojennsa Watson nom.

LITH

JOSIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 176.

Type-species: *Hypocrita ligula* Hübner, [1808] 1806, *Sanıml. exot. Schmett.* 1: pl.[180], by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 406. Type-locality: not stated, [Surinam].

JOSIODES Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.28.

Type-species: Josiodes batesii Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.28, by monotypy. Brazil: Amazonas, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 5.

Felder, 1874, in his script legends to his plates did not distinguish between the capital letters I and J; the spelling *Josiodes* has been adopted by authors.

‡JOSIOIDES; Butler, 1877, Trans. ent. Soc. Lond. 1877: 331. An incorrect subsequent spelling of Josiodes Felder, 1874.

‡JOSIOIDES; misspelling. See Josiodes Felder, 1874.

JOSIOMORPHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.104, fig.12. PERI

Type-species: Josiomorpha longivitta Felder, 1874, ibidem 2 (Abt.2): pl.104, fig.12, by monotypy. GUATEMALA, MEXICO: Orizaba. Both localities cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

Felder, 1874, in his script legends to his plates did not distinguish between the capital letters I and J; the spelling

Josiomorpha has been adopted by authors.

J. longivitta is a junior subjective synonym of Josia penetrata Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 134.

JOSIOMORPHIDES; misspelling. See Josiomorphoides Hering, 1925. PERI

JOSIOMORPHOIDES Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 434.

Type-species: Josia gigantea Druce, 1897, Biologia cent.-am. (Zool.) Lepid.-Heterocera 2: 406, pl.78, fig.25, by original designation. PANAMA: Volcan de Chiriqui.

The date of publication of the part of Seitz containing page 434 is cited on page 433 as 9.i.1925 in the German edition, and as 11.iii.1925 in the English edition.

‡JOSIOMORPHIDES; Bryk, 1931, in Strand, Lepid. Cat. 45:

An incorrect subsequent spelling of *Josiomorphoides* Hering, 1925.

JOZINIA Kiriakoff, 1970, Ann. Transv. Mus. 26 (9): 185.
NOTO

Type-species: *Jozinia leonina* Kiriakoff, 1970, *ibidem* **26** (9): 186, fig.4, by original designation. South Africa: Natal, Lebombo Mts, Jozini Dam.

‡*JSCHNOCAMPA*; misspelling. See *Ischnocampa* Felder, 1874.

‡JSCHNOGNATHA; misspelling. See Ischnognatha Felder, 1874.

‡JSOCTENIA; misspelling. See Isoctenia Felder, 1874.

‡JSOSTOLA; misspelling. See Isostola Felder, 1874. PERI

KAKOSTAUROPUS Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 632.

Type-species: Stauropus glaucoviridis Rothschild, 1917, Novit. zool. 24: 242, pl.3, fig.6, by monotypy. New Guinea: [Papua New Guinea], north eastern, Mambare, Biagi.

The date of publication of the part of Seitz containing page 632 is cited on page 625 as 11.iii.1930 in both the German and the English editions.

KALKOMA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 259 (key), 279.

Type-species: Symmerista alba Druce, 1898, Biologia cent.am. (Zool.) Lepid.-Heterocera 2: 460, pl.91, fig.3, by original designation. PANAMA: Volcan de Chiriqui. **KANCHIA** Moore, [1883] 1882-3, *Lepid. Ceylon* 2: 92.

Type-species: Leucoma subvitrea Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 344, by monotypy. [INDIA] Hindostan.

KARA Matsumura, 1925. See Parascotia Hübner, [1825].
NOCTUIDAE

KARSCHIOLA Ga'ede, 1926, in Seitz, Gross-Schmett. Erde 14: 112.

Type-species: Caryatis holoclera Karsch, 1894, Ent. Nachr. Berlin 20: 378, by monotypy. [Tanzania]: [Uzegua] Usegua.

The date of publication of the part of Seitz containing page 112 is cited on page 105 as 7.vii.1926 in both the German and English editions.

KASERIA Schaus, 1905, Proc. U.S. natn. Mus. 29: 282.

Type-species: *Kaseria pallida* Schaus, 1905, *ibidem* **29**: 282, by monotypy. French Guiana: Maroni River, St Jean.

KATHA Moore, 1878, Proc. zool. Soc. Lond. 1878: 16. LITH Type-species: Bombyx helvola Hübner, [1803] 1796, Samml. eur. Schmett. 3: pl.23, fig.95, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 130, 145, but cited as Noctua depressa Esper, 1787, a nominal species not originally included in Katha. Hampson, on page 145, placed depressa as the senior synonym of helvola, a nominal species originally included in Katha. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. Europe.

B. helvola is a junior subjective synonym of Noctua depressa

Esper, 1787, Die Schmett. 4 (1): 98, pl.93, fig.3.

KATMETEUGOA van Eecke, 1920, Zool. Meded. Leiden 5: 130.

Type-species: *Katmeteugoa hampsonia* van Eecke, 1920, *ibidem* 5: 130, text-fig.15, by monotypy. Java: Preanger.

KAWIELLA Roepke, 1943, Natuurh. Maandbl. 32 (9): 80. PYRALIDAE

Type-species: Kawiella testacea Roepke, 1943, ibidem 32 (9): 80, fig.10a; 1943, ibidem 32 (10): 88, fig.10b, by mon-

otypy. JAVA: W., Perbawattee.

Kawiella was established in the Notodontidae; it was transferred to the Lymantriidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2. Examination of the holotype of Kawiella testacea has shown it to be a junior subjective synonym of Sybrida ragonotalis Snellen, 1892, Tijdschr. Ent. 35: 153 in the Pyralidae, Pyralinae, syn. n. Kawiella is here transferred to the Pyralidae.

KERALA Moore, 1881, Proc. zool. Soc. Lond. 1881: 329.
NOCTUIDAE

Type-species: Kerala punctilineata Moore, 1881, ibidem 1881: 330, by monotypy.

Further information in Nye, 1975, Generic Names Moths World 1: 265.

‡KERASIA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 421.

An incorrect subsequent spelling of *Kerala* Moore, 1881. ‡*Kerasia* was cited by Neave, 1939, *Nomencl. zool.* 2: 824, as 'err. pro *Kaseria* Schaus, 1905' in the Notodontidae. However from the context of Pagenstecher's work it is almost certain that he was referring to *Kerala* Moore, 1881.

‡Kerasia Pagenstecher, in the Noctuidae, Chloephorinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

KERASIA; misspelling. See Kerala Moore, 1881.

KETTELIA Butler, 1879, Trans. Linn. Soc. Lond. (2) Zool. 1: 560.

Type-species: *Kettelia lowii* Butler, 1879, *ibidem* (2) Zool. 1: 560, by original designation. [BORNEO]: Sarawak.

KHASIDONTA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 175.

Type-species: *Notodonta picta* Hampson, 1900, *J. Bombay nat. Hist. Soc.* **13**: 42, by original designation. INDIA: Khasia [Hills].

‡KIKUCHIA; misspelling. See Kikuchiana Matsumura, 1927.

KIKUCHIANA Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19: 10.

Type-species: *Kikuchiana infuscata* Matsumura, 1927, *ibidem* 19: 10, pl.1, fig.9, by original designation. Talwan: Horisha, Baibara.

On the legend to plate 1, the generic name was published as ‡*Kikuchia*, an incorrect (of a multiple) original spelling.

‡KIKUCHIA; Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19: legend pl.1.

An incorrect (of a multiple) original spelling of *Kikuchiana* Matsumura, 1927.

KINTANA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 229.

Type-species: *Dasychira ocellatula* Hering, 1926, *in* Seitz, *Gross-Schmett. Erde* 14: 180, by original designation. MADAGASCAR: Central.

D. ocellatula is the objective replacement name for Dasychira ocellata Kenrick, 1914, Trans. ent. Soc. Lond. 1913: 594, pl.32, fig.27. D. ocellata Kenrick, and Laelia ocellata Holland, 1893, Psyche, Camb. 6: 432, pl.10, fig.8, were both placed in the genus Dasychira Hübner, by Hering, 1926.

KIRIAKOFFALIA Toulgoët, 1978, Nouv. Revue Ent. 8: 377.

Type-species: *Pericallia costimacula* Joicey & Talbot, 1924, *Bull. Hill Mus. Witley* 1: 553, by original designation (for *Kiriakoffia* Toulgoët, 1978). RWANDA: Rugege Forest, ZAIRE: E. Luvua valley.

Kiriakoffalia was established as an objective replacement name for Kiriakoffia Toulgoët, 1978.

KIRIAKOFFIA Toulgoët, 1978, Nouv. Revue Ent. 8: 221.

Type-species: *Pericallia costimacula* Joicey & Talbot, 1924, by original designation.

A junior homonym of *Kiriakoffia* Nakamura, 1974, *Trans. lepid. Soc. Japan* **25**: 117, – Lepid., Notodontidae. The objective replacement name is *Kiriakoffalia* Toulgoët, 1978.

KIRIAKOFFIA Nakamura, 1974, Trans. lepid. Soc. Japan 25: 117.

Type-species: *Allodonta longivitta* Gaede, 1930, *in* Seitz, *Gross-Schmett. Erde* **10**: 643, pl.80 row d, by original designation. [INDIA]: Sikkim.

Kiriakoffia was established as a subgenus of Allodonta Staudinger, 1887.

KIRIAKOFFIA Toulgoët, 1978. See Kiriakoffalia Toulgoët, 1978.

KNAPPETRA Nye nom. n. for Dulichia Walker, 1855. LYMA Type-species: Dulichia fasciata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 809, by monotypy (of Dulichia Walker, 1855). Sierra Leone.

This name is proposed in honour of Mrs Sheila D.M. Knapp in appreciation of her accurate clerical assistance during the compilation of this catalogue.

DULICHIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 809.

Type-species: *Dulichia fasciata* Walker, 1855, by monotypy. A junior homonym of *Dulichia* Kroyer, 1845, *Naturh. Tidsskr.* (2) 1: 512, 521, – Crustacea. The objective replacement name is *Knappetra* Nye nom. n.

KODIOSOMA Stretch, 1872, Illust. Zygaenidae Bombycidae N. Am. 1: 67.

Type-species: Kodiosoma fulva Stretch, 1872, ibidem 1: 67, pl.2, fig.7, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 244. U.S.A.: California, near San Francisco; San Mateo; and Saucelito.

CODIOSOMA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 244.

An unjustified emendation of *Kodiosoma* Stretch, 1872.

KOFFOLEANIA Laporte, 1977, in Rougeot, Mém. Mus. natn. Hist. nat. Paris (N.S.) A 105: 65.

NOCTUIDAE

Type-species: Koffoleania michaellae Laporte, 1977, ibidem 105: 65, pl.8, fig.91, by original designation. Holotype 3, ETHIOPIA: environs de Koffolé, (MNHN, Paris).

Koffoleania in the Noctuidae, Acronictinae is cited here as it was not included in Nye, 1975, Generic Names Moths World

‡KONOKARCHA; misspelling. See Konokareha Matsumura, [1928].

‡KONOKARECHA; misspelling. See Konokareha Matsumura, [1928].

KONOKAREHA Matsumura, [1928] 1927, Insecta matsum. 2: 120.

Type-species: Konokareha daisetsuzana Matsumura, [1928] 1927, ibidem 2: 120, fig.3, by original designation. [JAPAN]: Hokkaido, Mt Daisetsu.

K. daisetsuzana is a subspecies of Laria rossii Curtis, 1835, the type-species of Byrdia Schaus, 1927.

See also Byrdia Schaus, 1927.

‡KONOKARCHA; Gaede, 1932, in Seitz, Gross-Schmett. Erde 2 (Suppl.): 96.

An incorrect subsequent spelling of *Konokareha* Matsumura, [1928].

‡KONOKARECHA; Bryk, 1934, in Strand, Lepid. Cat. 62: 58, 393.

An incorrect subsequent spelling of *Konokareha* Matsumura, [1928].

KORAWA Moore, 1878, Proc. zool. Soc. Lond. 1878: 11.

Type-species: *Korawa pallida* Moore, 1878, *ibidem* **1878**: 12 (a secondary homonym; see below), by monotypy. INDIA: Darjeeling.

K. pallida Moore is a junior secondary homonym of Calligenia pallida Bremer, 1864, Mém. Acad. Sci. St Petersb. (7) 8: 97. The objective replacement name is Miltochrista perpallida Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 478, 495.

KUMATAIA Kiriakoff, 1967, Tijdschr. Ent. 110: 56. NOTO Type-species: Kumataia producta Kiriakoff, 1967, ibidem 110: 56, fig.10, by original designation. SUMATRA: south-western, western slopes Barisan Range.

KUNUSARA Nye nom. n. for Sankurua Collenette, 1960.

Type-species: Sankurua fontainei Collenette, 1960, Annls Mus. r. Congo Belge (8vo Sci. zool.) 80: 17, pl.A, fig.12, by

original designation (for Sankurua Collenette, 1960). [ZAIRE] Belgian Congo: Sankuru, Lusambo.

SANKURUA Collenette, 1960, Annls Mus. r. Congo Belge (8vo Sci. zool.) 80: 16.

Type-species: Sankurua fontainei Collenette, 1960, by orig-

inal designation.

A junior homonym of Sankurua Schouteden, 1951, Revue Zool. Bot. afr. 45: 149, - Insecta, Hemiptera. The objective replacement name is Kunusara Nye nom. n.

KUOHSINGIA Nakamura, 1974, Trans. lepid. Soc. Japan 25:

Type-species: Pydna nebulosa Wileman, 1914, Entomologist 47: 267, by original designation. [Talwan]: Arizan.

KURTIA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 263 (key),

Type-species: Kurtia modesta Schaus, 1901, ibidem 1901: 332, pl.12, fig.7, by original designation. VENEZUELA: Aroa.

LABORDEA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80:

Type-species: Calliteara prasina Butler, 1882, Cistula ent. 3: 16, by original designation. MADAGASCAR: Ankafana.

LACHANA Moore, 1888, Proc. zool. Soc. Lond. 1888: 397.

Type-species: Lachana ladakensis Moore, 1888, ibidem 1888: 398, by monotypy. [India/Pakistan]: [Kashmir], Ladak.

LACIDA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 801.

Type-species: Lacida antica Walker, 1855, ibidem 4: 802, by subsequent designation by Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 89. [SRI LANKA] Ceylon.

LACIDES Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 456.

Type-species: Noctua ficus Fabricius, 1775, Syst. Ent.: 595,

by monotypy. INDIA.

Lacides was established as a subgenus of Hypsa Hübner, [1819].

LACIPA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key), 790.

Type-species: Liparis picta Boisduval, 1847, in Delegorgue, Voyage Afr. australe 2: 599, by subsequent designation by Hampson, 1905, Ann. S. Afr. Mus. 3: 403. [RHODESIA]: [Matabeleland] Pays de Massilicatzi.

Massilicatzi probably refers to the first Matabele chief, also spelt as Mzilikazi, Msilikazi or Moselekatse. We are most grateful to Dr D. Kroon of Sasolburg, South Africa, for this

information.

MICROGYMNA Wallengren, 1865, K. svenska Vetensk Akad. Handl. (N.F.) 5 (4): 38.

Type-species: *Liparis picta* Boisduval, 1847, by monotypy. A junior objective synonym of Lacipa Walker, 1855.

LACYDES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 685.

Type-species: Eyprepia intercissa Freyer, 1842, Neuere Beitr. Schmett. 4: 118, pl.356 (as ‡incisa, an incorrect (of a multiple) original spelling), by monotypy. [U.S.S.R.] Russland: southern.

E. intercissa is a junior subjective synonym of Noctua spectabilis Tauscher, 1806, Mém. Soc. Nat. Univ. imp. Moscou 1: 212, fig.

See also Acymba Rambur, 1866.

LACYDOIDES Daniel, 1955, Bonn. zool. Beitr. 6: 135. LITH

Type-species: Lacydoides tibetensis Daniel, 1955, ibidem 6: 136, by original designation. CHINA: Tibet, Batang, Yangste.

LAELAPIA Butler, 1879, Ann. Mag. nat. Hist. (5) 4: 238.

Type-species: Laelapia notata Butler, 1879, ibidem (5) 4: 238, by monotypy. MADAGASCAR: Antananarivo.

Laelapia was established in the Liparidae, now Lymantriidae; it was transferred to the Arctiidae by Viette, 1949, Bull. mens. Soc. linn. Lyon 18: 170.

LAELAROA Hering, 1926, in Seitz, Gross-Schmett. Erde 14:

Type-species: Laelaroa flavimargo Hering, 1926, ibidem 14: 163, pl.28, row e, by original designation. [NIGERIA]: [R.]

Niger, Ogruga.

The date of publication of the part of Seitz containing page 163 is cited on page 161 as 5.xi.1926 in both the German and the English editions.

LAELIA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 62.

Type-species: Bombyx coenosa Hübner, [1808] 1796, Samml. eur. Schmett. 3: pl.51, fig.218; [1809], (Ziefer text): 120, by monotypy. [GERMANY (B.R.D.)]: Schwaben, Donauwörth (on page 120).

Laelia was proposed again by Stephens, 1829 [June], Nom. Br. Insects: 40; and 1829 [July], Syst. Cat. Br. Insects (2):

ANTHORA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (key), 801.

Type-species: Anthora subrosea Walker, 1855, ibidem 4: 801,

by monotypy. Sierra Leone.

A junior homonym of Anthora Doubleday, 1844, List Specimens lepid. Insects Colln Br. Mus. 1: 99, - Lepid., Nymphalidae. There is no objective replacement name, but A. subrosea is congeneric with Bombyx coenosa Hübner, [1808], the typespecies of Laelia Stephens, 1828. The latter is therefore available for use as a subjective replacement name.

LAELIOIDES Moore, [1883] 1882-3, Lepid. Ceylon 2: 83.

Type-species: Laelioides fasciata Moore, [1883] 1882-3, ibidem 2: 84, pl.110, fig.6, by monotypy. [Sri Lanka] Ceylon.

LAELIOLINA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 154.

Type-species: Laeliolina paetula Hering, 1926, ibidem 14: 154, pl.28, row e, by original designation. MADAGASCAR.

The date of publication of the part of Seitz containing page 154 is cited on page 153 as 5.xi.1926 in both the German and the English editions.

LAELIOPHILA Hering, 1932, Revue Zool. Bot. afr. 22: 105.

Type-species: Laeliophila seydeli Hering, 1932, ibidem 22: 105, pl.2, fig.23, by original designation. [ZAIRE]: [Katanga], [Lubumbashi] Elisabethville.

LAELIOPROCTIS Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 143.

Type-species: Laelioproctis taeniosoma Hering, 1926, ibidem 14: 143, pl.22, row a, by monotypy. Togo: Bismarckburg.

The date of publication of the part of Seitz containing page 143 is cited on page 137 as 1.xi.1926 in both the German and the English editions.

LAEMOCHARIS Herrich-Schäffer, [1854] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schniett. 1 (1): wrapper, pls 46-48, figs 238-258.

Type-species: Laemocharis pertyi Herrich-Schäffer, [1854]

1850-1858, *ibidem* 1 (1): wrapper, pl.47, fig.249 (but attributed to Boisduval, an incorrect authorship), by subsequent designation by Grote, 1873, *Bull. Buffalo Soc. nat. Sci.* 1: 32. [U.S.A.] Am.: [Georgia] Georg.

The name Laemocharis probably originated from Boisduval who attributed it to himself in Boisduval, 1870, Considérations

Lépid. Guatemala: 80.

HYALOPIS Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 101, 103.

Type-species: *Laemocharis pertyi* Herrich-Schäffer, [1854], by PRESENT DESIGNATION.

A junior objective synonym of *Laemocharis* Herrich-Schäffer, [1854].

Hyalopis was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3, under Dates of some important Publications.

LAFAJANA Dognin, 1891, Naturaliste 13: 257.

Type-species: Lafajana cupra Dognin, 1891, ibidem 13: 258, by monotypy (but cited as ‡Lafayana, an incorrect original spelling). [ECUADOR]: Loja.

Lafajana was the spelling adopted by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 926, and later authors from a multiple original spelling.

‡LAFAYANA; Dognin, 1891, Naturaliste 13: 258.

An incorrect (of a multiple) original spelling of Lafajana Dognin, 1891.

‡LAFAYANA; misspelling. See Lafajana Dognin, 1891.

LAGANDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 389.

BOMBYCIDAE

Type-species: *Laganda picaria* Walker, 1865, *ibidem* 32: 389, by monotypy. [NEW GUINEA]: [Irian Jaya], [Misoöl I.] Mysol.

Laganda was established in the Liparidae, now Lymantriidae; it was placed in the Bombycidae by Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 255.

LAGARIA Walker, 1854. See Rezia Kirby, 1892. CTEN

LAGELA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1086.

Type-species: Lagela fulva Walker, 1855, ibidem 5: 1086, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 620. [Brazil]: Rio [de] Janeiro.

LAMA Walker, 1869. See Ephestris Hübner, [1820]. PERI

LAMBULA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1890.

Type-species: *Lambula melaleuca* Walker, 1866, *ibidem* 35: 1890, by monotypy. [INDONESIA]: [Moluccas], Sula [I.].

LAMBULODES Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xix, 447.

Type-species: *Lambula sericea* Rothschild, 1912, *Novit. zool.* 19: 212, by original designation. New Guinea: [Irian Jaya], Mt Goliath.

LAMBULOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxii, 582.

Type-species: *Lambula aurantiaca* Rothschild, 1912, *Novit. zool.* 19: 214, by original designation. New GUINEA: [Irian Jaya], Mt Goliath.

LAMNIS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1072.

Type-species: *Lamnis divisa* Walker, 1855, *ibidem* 5: 1073, by monotypy. [Brazil]: Rio de Janeiro.

LAMORINA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 426.

Type-species: *Lamorina nitens* Gaede, 1928, *ibidem* 14: 427, pl.71 row a, by monotypy. EAST AFRICA.

The date of publication of the part of Seitz containing page 427 is cited on page 425 as 30.iii.1928 in both the German and

the English editions.

LAMORIODES Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 478.

Type-species: *Lamoriodes metaleuca* Hampson, 1910, *ibidem* (8) **5**: 479, by original designation. SIERRA LEONE.

LAMPROBALACRA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 72.

Type-species: *Balacra pulchra* Aurivillius, 1892, *Ent. Tidskr.* 13: 200, by original designation. CAMEROUN.

Lamprobalacra was established as a subgenus of Balacra Walker, 1856.

LAMPRONADATA Kiriakoff, 1967. See Rabtala Draeseke, 1926.

LAMPROSIA Hampson, 1900. See Lamprosiella Collins, 1962.

LAMPROSIELLA Collins, 1962, Ann. Mag. nat. Hist. (13) 5: 125.

Type-species: *Lithosia eborella* Boisduval, 1847, *in* Delegorgue, *Voyage Afr. australe* 2: 597, by original designation (for *Lamprosia* Hampson, 1900). [SOUTH AFRICA]: 'Pays de Amazoulous'.

Lamprosiella was established as the objective replacement name for Lamprosia Hampson, 1900.

LAMPROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 86, 217.

Type-species: Lithosia eborella Boisduval, 1847, by original designation.

A junior homonym of *Lamprosia* Hübner, [1821] 1806, *Samml. exot. Schmett.* 2: pl.[214], – Lepid., Noctuidae. The objective replacement name is *Lamprosiella* Collins, 1962.

LAMPROSOMA Grote, 1863. See Omoiala Grote, 1863.

LAMPROSTOLA Schaus, 1899, Jl N.Y. ent. Soc. 7: 216.

Type-species: Lamprostola molybdipera Schaus, 1899, ibidem 7: 216, by monotypy. MEXICO: Jalapa.

The name *Lamprostola* probably originated from Hampson, but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson, 1900.

LAMPROSTOLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 89, 360.

Type-species: Odozana olivacea Schaus, 1896, Jl N.Y. ent. Soc. 4: 149, by original designation. BRAZIL: São Paulo.

A junior homonym of *Lamprostola* Schaus, 1899. There is no objective replacement name but the type-species of *Lamprostola* Hampson is currently considered to be congeneric with

the type-species of Lamprostola Schaus. The latter is therefore available for use as a subjective replacement name.

LAMPROSTOLA Hampson, 1900. See Lamprostola Schaus,

LAMPRUNA Schaus, 1894, Proc. zool. Soc. Lond. 1894: 231.

Type-species: Lampruna rosea Schaus, 1894, ibidem 1894: 231, by monotypy. VENEZUELA: Aroa.

LANITRA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 229.

Type-species: Dasychira hexamitobalia Collenette, 1936, Novit. zool. 40: 163, by original designation. MADAGASCAR: E., Périnet (149km E. of Tananarivo).

LAORA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 6: 1482.

Type-species: Laora antennata Walker, 1855, ibidem 6: 1482, by monotypy. [Colombia]: Sante Fé de Bogota.

LARIA Schrank, 1802. See Leucoma Hübner, 1822.

LASIOCEROS Bethune-Baker, 1904, Novit. zool. 11: 380.

Type-species: Lasioceros aroa Bethune-Baker, 1904, ibidem 11: 381, by original designation. New Guinea: [Papua New Guinea], Upper Aroa River.

Lasioceros was established in the Notodontidae; it was transferred to the Hypsidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2; and is here transferred back to the Notodontidae, on the basis of tympanal and venational

LASIOPROCTA Wallengren, 1858, Ofvers. K. VetenskAkad. Förh. Stockh. 15: 135.

Type-species: Lasioprocta merra Wallengren, 1858, ibidem 15: 136, by original designation. [PERU]: Callao.

L. merra is a junior subjective synonym of Glaucopis subapicalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 190.

Lasioprocta was redescribed as new by Wallengren, 1861, K. svenska Fregatten Eugenies Resa omkring Jorden 2 (1): 358.

LASIOSCELES Hampson, 1909. See Scelilasia Hampson, 1914.

LAURON Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 333.

Type-species: Dioptis rica Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 31, pl.[92], figs 531, 532, by subsequent designation by Bryk, 1931, in Wagner, Lepid. Cat. 45: 41. CUBA.

Lauron Walker was established as a subgenus of Dioptis Hübner, 1818, in the Dioptidae; it was placed in the Arctiidae, Pericopinae by Hering, 1925, in Seitz, Gross-Schmett. Erde 6:

‡LAURONA; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1653.

An incorrect subsequent spelling of Lauron Walker, 1854.

‡LAURONA; misspelling. See Lauron Walker, 1854. PERI

LAVINIA Kiriakoff, 1962. See Eulavinia Fletcher nom. n.

LEBADIA Walker, 1869, Characters undescr. Lepid. Hetero-

Type-species: Lebadia cervina Walker, 1869, ibidem: 107, by monotypy. Type-locality: not stated.

The type has not been located and the type-species remains unrecognized.

LECHRIOLEPIS Butler, 1880, Ann. Mag. nat. Hist. (5) 5: 385. LASIOCAMPIDAE

Type-species: Lechriolepis anomala Butler, 1880, ibidem (5)

5: 386, by original designation. MADAGASCAR.

Lechriolepis was established in the Liparidae, now Lymantriidae; it was placed in the Bombycidae by Dewitz, 1881, Nova Acta Acad. Caesar. Leop. Carol. 42 (2): 78; it was retained in the Liparidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 497; and as ‡Lecriolepis, an incorrect subsequent spelling, was transferred to the Lasiocampidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498.

‡LECRIOLEPIS Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498.

An incorrect subsequent spelling of Lechriolepis Butler, 1880.

‡LECRIOLEPIS; misspelling. See Lechriolepis Butler, 1880. LASIOCAMPIDAE

LEIOCAMPA Stephens, 1828. See Pheosia Hübner, [1819].

‡LEIOCAMPHA; misspelling. See Pheosia Hübner, [1819].

LEMAIREGISA Thiaucourt, 1978, Lambillionea 77: 72. NOTO

Type-species: Lemairegisa limonensis Thiaucourt, 1978, ibidem 77: 73, figs 3, 4, by original designation. ECUADOR: Morona Santiago, 16km east of Limon.

‡LEMODONTA; misspelling. See Leucodonta Staudinger, 1892.

LEMPKEELLA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 76.

Type-species: Apisa dufranei Kiriakoff, 1952, Revue Zool. Bot. afr. 46: 402, figs 7, 8, by original designation. [ZAIRE]: [Katanga], [Lubumbashi] Elisabethville.

LEMURIANA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80:

Type-species: Dasychira mascarena Butler, 1878, Ann. Mag. nat. Hist. (5) 2: 294, by original designation. MADAGASCAR: Fianarantsoa.

LEMYRA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1690.

Type-species: Lemyra extensa Walker, 1856, ibidem 7: 1690, by monotypy. [SULAWESI] Celebes.

LEOPOLDINA Hering, 1934, Bull. Mus. r. Hist. nat. Belg. 10

Type-species: Leopoldina leopoldi Hering, 1934, ibidem 10 (21): 2, fig.1, by original designation. THAILAND.

LEPASTA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 694.

Type-species: Nystalea bractea Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.3, by monotypy. VENEZUELA: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 7.

LEPIDILEMA Aurivillius, 1910, in Sjöstedt, Wiss. Ergebn. schwed. zool. Exped. Kilimandjaro, Meru . . . 1905-1906 (9):

Type-species: Lepidilema unipectinata Aurivillius, 1910. ibidem (9): 26, fig.5, by monotypy. [Tanzania]: [Kilimanjaro]] Kilimandjaro, Kibonoto.

LEPIDOJULIA Orfila, 1952, Revta Soc. ent. argent. 15: 207.

Type-species: *Lepidojulia arnaui* Orfila, 1952, *ibidem* **15**: 300, fig., by original designation. ARGENTINA: Tucuman Province, Dept. of Trancas, San Pedro de Colalao.

LEPIDOKIRBYIA Travassos, 1943, Revta bras. Biol. 3: 463.

Type-species: Arara vittipes Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 642, by monotypy (of Arara Walker, 1855). BRAZIL.

Lepidokirbyia was established as the objective replacement name for Arara Walker, 1855.

ARARA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 641.

Type-species: Arara vittipes Walker, 1855, by monotypy.

A junior homonym of *Arara* Spix, 1844, *Avium Species novae Bras.* 1: index, 25, – Aves. The objective replacement name is *Lepidokirbyia* Travassos, 1943.

LEPIDOLACIPA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 141.

Type-species: *Lepidolacipa venosata* Hering, 1926, *ibidem* 14: 141, pl.28, row d, by original designation. [MALAWI] Nyassa-Land: Nehen.

The date of publication of the part of Seitz containing page 141 is cited on page 137 as 1.xi.1926 in both the German and the English editions.

LEPIDOLUTZIA Rego Barros, 1956, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 145: 8.

Type-species: Charidea baucis Dalman, 1823, Analecta Ent.: vi, 50, by original designation. BRAZIL.

LEPIDONEIVA Travassos, 1940, Revta Ent., Rio de J. 11: 480.

Type-species: Cosmosoma erubescens Butler, 1876, J. Linn. Soc. (Zool.) 12: 389, by original designation. BRAZIL.

LEPIDONEWTONIA Rego Barros, 1956, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 140: 1.

Type-species: *Halysidota acuta* Hampson, 1901, *Cat. Lepid. Phalaenae Br. Mus.* 3: 167, pl.41, fig.21, by original designation (but cited as ‡*Halisidota*, an incorrect subsequent spelling). BRAZIL: Sta Catarina.

LEPIDOPALPUS Janse, 1915, Ann. Transv. Mus. 5: 5 (key), 9.

Type-species: *Lepidopalpus hyalina* Janse, 1915, *ibidem* 5: 10, pl.1, fig.4, pl.3, fig.15, by original designation. South Africa: [Cape Province], Ngqeleni.

Lepidopalpus was established in the Lymantriidae; it was placed in the Pterothysanidae by Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 124; and was restored to the Lymantriidae by Collenette, 1955, Trans. R. ent. Soc. Lond. 107: 191.

LEPIDOZIKANIA Travassos, 1949, Revta bras. Biol. 9: 453.

Type-species: *Halysidota cinerascens* Walker, 1855, *List Specimens lepid. Insects Colln Br. Mus.* 3: 742, by original designation (but cited as ‡*Halesidota*, an incorrect subsequent spelling). [BRAZIL]: Rio [de] Janeiro.

LEPISTA Wallengren, 1863, Wien. ent. Monatschr. 7: 146.

Type-species: *Lithosia pandula* Boisduval, 1847, *in* Delegorgue, *Voyage Afr. australe* 2: 597, by original designation. [RHODESIA]: [Matabeleland] Pays de Massilicatzi.

Lepista was redescribed as new by Wallengren, 1865, K. svenska Vetensk Akad. Handl. 5 (4): 42.

Massilicatzi probably refers to the first Matabele chief, also

spelt as Mzilikazi, Msilikazi or Moselekatse. We are most grateful to Dr D. Kroon of Sasolburg, South Africa, for this information.

See also Dyphlebia Felder, 1874.

LEPTACTEA Prout, 1918, Novit. zool. 25: 397 (key), 420.

Type-species: Actea minuta Druce, 1885, Proc. zool. Soc. Lond. 1885: 526, by original designation. ECUADOR: Sarayacu.

LEPTARCTIA Stretch, 1872, Illust. Zygaenidae Bombycidae N. Am. 1: 118.

Type-species: *Lithosia decia* Boisduval, 1869, *Annls Soc. ent. Belg.* 12: 72, by subsequent designation by Barnes & Benjamin, 1927, *Bull. Sth. Calif. Acad. Sci.* 26: 46. U.S.A.: California.

L. decia is a junior subjective synonym of Nemeophila californiae Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 625.

Incorrect type-species: (1) Nemeophila californiae Walker, 1855, a nominal species not originally included in Leptarctia and not linked in synonymy with one of the originally included species when designated by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 276.

(2) Nemeophila californiae Walker, 1855, a nominal species not originally included in Leptarctia but placed as a senior subjective synonym of all three originally included nominal species when designated by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 220.

LEPTAROA Hampson, 1910, Proc. zool. Soc. Lond. 1910: 456.

LYMA

Type-species: *Leptaroa fulvicolora* Hampson, 1910, *ibidem* 1910: 456, pl.38, fig.5, by original designation. [ZAIRE] Congo: Katanga, Kambove.

LEPTEPILEPTA Collenette, 1929, Entomologist 62: 274.

Type-species: *Leptepilepta caudata* Collenette, 1929, *ibidem* **62**: 275, fig., by original designation. MADAGASCAR: central, Brickaville.

LEPTOCERYX Kiriakoff, 1953, Lambillionea 53: 52. CTEN Type-species: Leptoceryx pusilla Kiriakoff, 1953, ibidem 53: 53, by original designation. [ZAIRE]: Tshuapa R., Eala.

LEPTOCNERIA Butler, 1886, Trans. ent. Soc. Lond. 1886:

Type-species: *Leptocneria binotata* Butler, 1886, *ibidem* **1886**: 386, 441, pl.9, fig.3, by monotypy. Australia: [Queensland], Peak Downs.

LEPTOLEPIDA Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 559.

Type-species: Leptolepida malangae Bethune-Baker, 1911, ibidem (8) 7: 559, by original designation. [ANGOLA]: Malange.

‡LEPTOLIPIDIA; Neave, 1939, Nomencl. zool. 2: 913.

An incorrect subsequent spelling of *Leptolepida* Bethune-Baker, 1911.

‡LEPTOLIPIDIA; misspelling. See Leptolepida Bethune-Baker, 1911.

LEPTONADATA Aurivillius, 1904, Ark. Zool. 2 (4): 2. NOTO Type-species: Leptonadata sjostedti Aurivillius, 1904, ibidem 2 (4): 3, text-fig.10, by monotypy. CAMEROUN.

L. sjostedti is based on the Swedish name Sjöstedt, not on a German word and does not require more than the deletion of the diacritic mark under the Code, Article 32(c)(i).

LEPTONADATOIDES Kiriakoff, 1969, Bull. Inst. fond. Afr. noire (A) 30 (4): 1449.

Type-species: *Leptonadatoides incurvata* Kiriakoff, 1969, *ibidem* (A) **30** (4): 1450, fig.9, by original designation. IVORY COAST: Adiopodoumé.

The date of publication of *Leptonadatoides* as January 1969, is taken from the foot of page 1762 of part 4, not from the titlepage of that part which is dated October 1968.

LEPTOPEPLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 90, 520.

Type-species: *Leptopepla procridiformis* Hampson, 1900, *ibidem* 2: 521, fig.374, by original designation. BOLIVIA: Songo, PERU: E., Huayabamba.

LEPTOSOMA Boisduval, 1832. See *Nyctemera* Hübner, [1820].

‡LEPTOSOMUS; misspelling. See Nyctemera Hübner, [1820].

LEPTOSPHETTA Butler, 1878, Trans. ent. Soc. Lond. 1878: 67. PYRALIDAE

Type-species: Leptosphetta rabdina Butler, 1878, ibidem 1878: 67, by original designation. [BRAZIL]: Amazon region, Rio Jutahi, near Rio Curnem.

Leptosphetta was established in the Notodontidae; it was placed in the Pyralidae by Hampson, 1896, Trans. ent. Soc. Lond. 1896: 471.

LEPTOTHRIX Heylaerts, 1892. See *Chionaema* Herrich-Schäffer, 1855.

LEPYPIRANGA Rego Barros, 1966, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 255: 5.

Type-species: *Ypiranga nigroruba* Rego Barros, 1959, *ibidem* (Nov. Ser.) Zool. **192**: 3, by original designation (for *Ypiranga* Rego Barros, 1959). BRAZIL: São Paulo, Campos de Jordão, Lagoinha.

Lepypiranga was established as the objective replacement name for Ypiranga Rego Barros, 1959.

YPIRANGA Rego Barros, 1959, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 192: 1.

Type-species: *Ypiranga nigroruba* Rego Barros, 1959, by

original designation.

A junior homonym of *Ypiranga* Mello-Leitão, 1922, *Ann. Mag. nat. Hist.* (9) 9: 331, – Arachnida. The objective replacement name is *Lepypiranga* Rego Barros, 1966.

LERINA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 477.

Type-species: *Lerina incarnata* Walker, 1854, *ibidem* **2**: 477, by monotypy. MEXICO.

LERNA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 33: 805.

Type-species: Lerna nivosa Walker, 1865, ibidem 33: 805, by

monotypy. AUSTRALIA.

Lerna was established in the Noctuidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 431; and was transferred to the 'Lithosiidae' by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497.

LEUCALOA Butler, 1875, Cistula ent. 2: 44.

Type-species: Spilosoma eugraphica Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 292, by original designation. SOUTH AFRICA.

LEUCANIOCHROA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.95, fig.17.

Type-species: Leucaniochroa lignosa Felder, 1874, ibidem

2 (Abt.2): pl.95, fig.17, by monotypy. Type-locality: not known, cited as 'Patria?' by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 6.

The type-specimen of *L. lignosa* has not yet been found nor has its figure been matched in the BMNH. *Leucaniochroa* was not placed in a family when it was established; it was placed in the Notodontidae by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 612.

LEUCANOPSIS Rego Barros, 1956, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 132: 1.

Type-species: *Halysidota leucanina* Felder & Rogenhofer, 1874, *Reise öst. Fregatte Novara* (Zool.) **2** (Abt.2): pl.109, fig.1, by original designation (but cited as ‡*Halesidota*, an incorrect subsequent spelling). [COLOMBIA]: Bogota.

LEUCARCTIA Packard, 1864. See Estigmene Hübner, [1820].

LEUCODONTA Staudinger, 1892, in Romanoff, Mém. Lépid. 6: 349.

Type-species: *Bombyx bicoloria* [Denis & Schiffermüller], 1775, *Ankündung syst. Werkes Schmett. Wienergegend*: 49, by subsequent designation (for *Microdonta* Duponchel, [1845]) by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 603. [AUSTRIA]: Vienna district.

Leucodonta was proposed as the objective replacement name for *Microdonta* Duponchel, [1845].

See also Shironia Matsumura, 1925.

MICRODONTA Duponchel, [1845] 1844, Cat. méth. Lépid. Eur. (2): 93.

Type-species: *Bombyx bicoloria* [Denis & Schiffermüller], 1775, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 603.

A junior homonym of *Microdonta* Dejean, 1835, *Cat. Coléopt*. (Edn 2): 364, – Insecta, Coleoptera. The objective replacement name is *Leucodonta* Staudinger, 1892.

HIEROPHANTA Meyrick, 1894, Entomologist's mon. Mag. 30: 230.

Proposed unnecessarily as the objective replacement name for *Microdonta* Duponchel, [1845]; the name *Leucodonta* Staudinger, 1892, had already been established.

‡LEMODONTA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 428.

An incorrect subsequent spelling of *Leucodonta* Staudinger, 1892.

LEUCOLOPHA Hampson, 1896, *Fauna Br. India* (Moths) 4: 460.

Type-species: *Leucolopha undulifera* Hampson, 1896, *ibidem* **4**: 460, fig.245, by original designation. INDIA: Khasis.

‡LEUCOMA Hübner, [1806]. See Euproctis Hübner, [1819].

LEUCOMA Hübner, 1822, Syst.-alphab. Verz.: 14-16, 18, 19.

Type-species: *Phalaena salicis* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 502, by subsequent designation by Westwood, 1840, *Introd. mod. Classif. Insects* 2 (Synopsis Genera Br. Insects): 92. [EUROPE].

Incorrect type-species designation: Bombyx v-nigrum Fabricius, 1775, cited by Boisduval, 1836, Hist. nat. Insectes (Spec. gén. Lépid.) 1: 143. Boisduval in his lengthy 154 page Introduction, reviewed earlier classifications and designated up to three different type-species for each generic name. In his 'Exposé de notre méthode' from page 155-690, no type-species designations were made for the genera he himself used. Under the Code, Article 69(a)(iii), the type-designation of an author is eligible for consideration if he states that it is the type '...

and if it is clear that he himself accepts it as the typespecies.' Boisduval's type-designations although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not in the past been accepted by Hemming or other authors.

LARIA Schrank, 1802, Fauna Boica 2 (2): 150.

Type-species: Phalaena salicis Linnaeus, 1758, by subsequent designation by Kozhanchikov, 1950, Fauna SSSR (N.S.) 42 (Lepid. 12): 340.

A junior homonym of Laria Scopoli, 1763, Ent. Carniolica: 21, - Coleoptera. The objective replacement name is Leucoma Hübner, 1822.

STILPNOTIA Westwood, 1843, in Humphreys & Westwood, Br. Moths Transformations 1: 90.

Type-species: Phalaena salicis Linnaeus, 1758, by monotypy. A junior objective synonym of Leucoma Hübner, 1822.

LEUCOSIA Rambur, 1866, Cat. syst. Lépid. Andalousie (2):

Type-species: *Phalaena salicis* Linnaeus, 1758, by monotypy. Leucosia is dated from the wrapper of livraison 2, and not from the title-page of the work, 1858.

A junior homonym of Leucosia Weber, 1795, Nomencl. ent. ... Fabricii: 92, - Crustacea. The objective replacement name is Leucoma Hübner, 1822.

Leucosia Rambur, 1866, under the ruling of Opinion 712 was placed on the Official Index of Rejected and Invalid Names in Zoology: Name No.1733.

NYMPHYXIS Grote, 1895, Mitt. Roemermus. Hildesh. 1: [4]. Type-species: *Phalaena salicis* Linnaeus, 1758, by monotypy. A junior objective synonym of Leucoma Hübner, 1822.

LEUCOPARDUS Hampson, 1894, Fauna Br. India (Moths) 2: vi, 2, 31.

Type-species: Leucopardus tigrinus Hampson, 1894, ibidem 2: 31, fig. 12, by original designation. [INDIA]: [Assam], Khasia Hills.

LEUCOPERINA Aurivillius, 1909, Ark. Zool. 5 (5): 8. LYMA Type-species: Leucoperina atroguttata Aurivillius, 1909, ibidem (5) 5: 8, by monotypy. [ZAIRE] Congo State: Bakoko.

‡SYNTAXIS Hering, 1926, in Seitz, Gross-Schmett. Erde 14:

An unavailable name under the Code, Article 11(d).

‡Syntaxis was first published as 'Syntaxis umbrinorma Tessm.' in the synonymy of Leucoperina atroguttata Aurivillius, and has not subsequently been treated as an available name.

LEUCOPHALERA Janse, 1920, Ann. Transv. Mus. 7: 199.

Type-species: Leucophalera elegans Janse, 1920, ibidem 7: 199, pl.5, fig.8, pl.6, figs 22-25, pl.14, fig.4, by original designation. [ZIMBABWE-]RHODESIA: Umtali.

L. elegans is a junior subjective synonym of Phalera latipennis Butler, 1897, Proc. zool. Soc. Lond. 1896: 844, pl.42, fig.3.

LEUCOPLEURA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xviii, 27, 423.

Type-species: Cosmosoma cucadma Druce, 1894, Ann. Mag. nat. Hist. (6) 13: 171, by original designation. PANAMA: Bugaba.

‡LEUCOPSUMIA; misspelling. See Leucopsumis Hübner, [1820].

LEUCOPSUMIS Hübner, [1820] 1816, Verz. bekannter Schmett.: 177.

Type-species: Phalaena collaris Drury, 1782, Illust. nat. Hist. exot. Insects 3: index, 27, pl.21, fig.2, by subsequent designation by Bryk, 1931, in Wagner, Lepid. Cat. 45: 13. BRAZII.

P. collaris is a junior subjective synonym of Phalaena amica Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 158, pl.370, fig.H. P. amica is dated from the wrapper of the part and not from the title-page of the volume, 1782.

See also Calodesma Hübner, [1820].

‡LEUCOPSUMIA; Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 288.

An incorrect subsequent spelling of Leucopsumis Hübner, [1820].

LEUCORHODIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 85, 354.

Type-species: Lithosia ragua Druce, 1897, Biologia cent.-am. (Zool.) Lepid.-Heterocera 2: 400, pl.78, fig.12, by original designation. GUATEMALA: Pantaleon.

LEUCOSIA Rambur, 1866. See Leucoma Hübner, 1822.

LEUCOTMEMIS Butler, 1876, J. Linn. Soc. (Zool.) 12: 391.

Type-species: Glaucopis latilinea Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 167, by original designation. BRAZIL.

PHACUSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 175.

Type-species: Glaucopis varipes Walker, 1854, ibidem 1: 175, by PRESENT DESIGNATION. [Brazil]: Para.

A junior homonym of *Phacusa* Walker, 1854, *ibidem* 1: 150, Lepid., Zygaenidae. There is no objective replacement name but the type-species of Phacusa Walker, 1854, ibidem 1: 175 is currently considered to be congeneric with the type-species of Leucotmemis Butler, 1876. The latter is therefore available for use as a subjective replacement name.

Phacusa Walker, 1854, ibidem 1: 175 was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; its type-species G. varipes was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 220.

See also Mystrocneme Herrich-Schäffer, 1855.

LEUCULA Guenée, 1857, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 9: 134.

Type-species: Leucula empusaria Guenée, 1857, ibidem 9: 134; 1858, ibidem (Atlas): pl.17, fig.3, by subsequent designation by Hulst, 1896, Trans. Am. ent. Soc. 23: 317. Brazil. Leucula has been used as the original generic name for

species now placed in the Arctiidae.

LEUCULODES Dyar, [1903] 1902, Bull. U.S. natn. Mus. 52:

Type-species: Leucula lacteolaria Hulst, 1887, Entomologica

am. 2: 51, 120, by monotypy. U.S.A.: Arizona.

Leuculodes was established in the Liparidae, now Lymantriidae; it was transferred to the Pericopidae, now Arctiidae, Pericopinae, by Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 559.

LEXIS Wallengren, 1863, Wien. ent. Monatschr. 7: 145, 146.

Type-species: Lithosia bipunctigera Wallengren, 1860, ibidem 4: 45, by original designation. [South Africa]: Caffraria, eastern.

Lexis was redescribed as new by Wallengren, 1865, K. svenska VetenskAkad. Handl. 5 (4): 41.

LIBIDO Bryk, 1950, Ark. Zool. 42 (A) 19: 25. NOTO Type-species: Libido voluptuosa Bryk, 1950, ibidem 42 (A) 19: 26, pl.3, fig.11, by original designation. Burma: Kambaiti. LIBYSSA Dognin, 1911. See Eudognina Fletcher nom. n. NOTO

LICCANA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 152 (key), 177.

Type-species: Bireta terminicana Kiriakoff, 1962 [July 31], ibidem 98 (10): 178; 1962 October 20, Bonn. zool. Beitr. 13: 226, abb.1, fig.10, pl.1, fig.9 (as Bireta terminicana sp.nov.), by original designation. CHINA: Chekiang, [east Tien-mu-Shan]. Additional data cited by Kiriakoff, 1962 October 20, ibidem 13: 229.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr. to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper B. terminicana was described as a new species. As the dates show, the Belgian paper appeared first.

In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

Liccana was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 45.

LICHENIA Sodovsky, 1837. See Lithosia Fabricius, 1798.

LICNOPTERA Meyrick, 1889, Trans. ent. Soc. Lond. 1889: 460.

Type-species: *Licnoptera crocodora* Meyrick, 1889, *ibidem* 1889: 461, by monotypy. New Guinea.

MONOTROPIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 307.

An unnecessary replacement name for *Licnoptera* Meyrick, 1889, which Kirby probably considered to be a junior homonym of *Lichnoptera* Herrich-Schäffer, 1855, *Syst. Bearbeitung Schmett. Eur.* 6: 94, 96, – Lepid., Noctuidae.

‡LIMBINA; misspelling. See Cimbina Walker, 1865.

LIOCAMPA Agassiz, 1847. See Pheosia Hübner, [1819].

LIPARIS Ochsenheimer, 1810. See Penthophera Germar, 1812.

LIPARODONTA Hering, 1927, in Seitz, Gross-Schmett. Erde 14: 197.

Type-species: *Liparodonta convexa* Hering, 1927, *ibidem* 14: 197, by original designation. [CAMEROUN] Kamerun: Lolodorf.

The date of publication of the part of Seitz containing page 197 is cited on page 193 as 4.i.1927 in both the German and the English editions.

LIPAROPSIS Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 125 (key), 154.

Type-species: *Liparopsis postalbida* Hampson, [1893] 1892, *ibidem* 1: 154, fig.93, by original designation. INDIA: Naga Hills.

LIPARUS Harris, 1841. See Penthophera Germar, 1812.

LIRIMIRIS Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 468.

Type-species: *Lirimiris lignitecta* Walker, 1865, *ibidem* 32: 469, by monotypy. Type-locality: not stated. The species

is now known to be widely distributed in CENTRAL and SOUTH AMERICA from Mexico to Brazil.

LIRIOCHROA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 67.

Type-species: *Desmeocraera divergens* Kiriakoff, 1954, *Revue Zool. Bot. afr.* **49**: 303, pl.1, fig.7, pl.2, figs 2, 3, by original designation. [ZAIRE]: [Lubumbashi] Elisabethville.

LISTOCA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 1021. LASIOCAMPIDAE

Type-species: *Listoca lignaria* Walker, 1855, *ibidem* 5: 1021, by monotypy. [Australia]: [New South Wales], Sydney.

Listoca was established in the Notodontidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 841.

LITHOPROCRIS Dognin, 1899, Annls Soc. ent. Belg. 43: 133.

Type-species: *Lithoprocris jason* Dognin, 1899, *ibidem* **43**: 133, by monotypy. [ECUADOR]: Zamora valley.

This name probably originated from Hampson but was used and made nomenclaturally available by Dognin prior to its proposal as a new generic name by Hampson, 1900.

LITHOPROCRIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 86, 342.

Type-species: Lithoprocris methyalea Hampson, 1900, ibidem 2: 342, fig.246, by original designation. BOLIVIA.

A junior homonym of *Lithoprocris* Dognin, 1899. There is no objective replacement name but the type-species of *Lithoprocris* Hampson, 1900, is currently considered to be congeneric with the type-species of *Lithoprocris* Dognin, 1899. The latter is therefore available for use as a subjective replacement name.

LITHOPROCRIS Hampson, 1900. See Lithoprocris Dognin, 1899.

‡LITHOSA; misspelling. See Lithosia Fabricius, 1798.

LITHOSARCTIA Daniel, 1954, Bonn. zool. Beitr. 5: 138.

Type-species: *Lithosarctia hoenei* Daniel, 1954, *ibidem* 5: 138, pl.3, fig.77, by original designation (but cited as ‡hönei, an incorrect original spelling). CHINA: N. Yunnan, Li-kiang.

LITHOSIA Fabricius, 1798, Suppl. Ent. Syst.: 418, 459.

Type-species: *Phalaena quadra* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 511, by subsequent designation by Latreille, 1810, *Considérations générales sur l'Ordre naturel des Animaux*: 441 (but attributed to Fabricius, an incorrect authorship). [EUROPE].

Latreille, 1810, listed three nominal species but of these only quadra was an originally included name. Acceptance of such type designations by Latreille was approved by the International Commission on Zoological Nomenclature, 1939, Opin. Decl. int. Commn zool. Nomencl. 2 (Opinion 136): 15.

‡LITHOSA; Lewin, 1805, Prodromus Ent., nat. Hist. lepid. Insects N.S.W.: 16.

An incorrect subsequent spelling of *Lithosia* Fabricius, 1798. *LITHOSIS* Billberg, 1820, *Enumeratio Insect. Mus. G.J. Billberg*: 91.

An unjustified emendation of Lithosia Fabricius, 1798.

LICHENIA Sodovsky, 1837, Bull. Soc. imp. Nat. Moscou 1837 (6): 85.

An unnecessary replacement name for *Lithosia* Fabricius, 1798, which Sodovsky considered to be an inappropriate name.

LITHOSIS Billberg, 1820. See Lithosia Fabricius, 1798.

JITH

LITODONTA Harvey, 1876, Can. Ent. 8: 5. NOTO Type-species: Litodonta hydromeli Harvey, 1876, ibidem 8: 5, by original designation. [U.S.A.]: Texas.

‡LOBERA; misspelling. See Lopera Walker, 1855. LYMA

LOBEZA Herrich-Schäffer, [1854] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.18, fig.85; [1856] ibidem 1: 11.

Type-species: *Lobeza aglone* Herrich-Schäffer, [1854] 1850-1858, *ibidem* 1 (1): wrapper, pl.18, fig.85 (but attributed to Boisduval, an incorrect authorship), by monotypy. BRAZIL.

The second species included in *Lobeza* on page 11 of the text was originally published in 1854 as *Anurocampa mingens*, so that at the time *Lobeza* was established it was a monotypic genus.

‡LOPEZA; Neave, 1939, Nomencl. zool. 2: 990.

An incorrect subsequent spelling of *Lobeza* Herrich-Schäffer, [1854].

LOBILEMA Aurivillius, 1910, in Sjöstedt, Wiss. Ergebn. schwed. zool. Exped. Kilimandjaro, Meru . . . 1905-1906 (9): 25.

Type-species: *Lobilema conspersa* Aurivillius, 1910, *ibidem* (9): 25, fig.3, by monotypy. [Tanzania]: [Kilimanjaro] Kilimandjaro, Kibonoto.

LOBOBASIS Hampson, 1896, Fauna Br. India (Moths) 4: xxv, 498.

Type-species: Lobobasis niveimaculata Hampson, 1896, ibidem 4: 498, fig., by original designation. INDIA: Bhutan.

LOBOGONA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.98, fig.4.

Type-species: Lobogona hapygia Felder, 1874, ibidem 2 (Abt.2): pl.98, fig.4, by monotypy. Surinam: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

L. hapygia is a junior subjective synonym of Corymbia simplex Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 33: 766.

LOCHARNA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 53.

Type-species: *Locharna strigipennis* Moore, 1879, *ibidem* (1): 53, pl.3, fig.11, by monotypy. INDIA: [Assam], Khasia Hills.

LOCHMAEUS Doubleday, 1841, Entomologist 1: 57. NOTO Type-species: Lochmaeus manteo Doubleday, 1841, ibidem
1: 58, by monotypy. U.S.A.: [New York], Trenton Falls. See also Tadana Walker, 1855.

LODA Kiriakoff, 1970, Tijdschr. Ent. 113: 116. NOTO Type-species: Loda celebensis Kiriakoff, 1970, ibidem 113: 116, fig.19, by original designation. [SULAWESI] Celebes: western, Paloe district, Loda.

LOMADONTA Holland, 1893, Psyche, Camb. 6: 417. LYMA Type-species: Lomadonta erythrina Holland, 1893, ibidem 6: 417, 567, pl.10, fig.1, by monotypy. [GABON]: [Ogooué] Ogové R. valley (see page 373).

LOMASPILIS Felder, 1874. See Argina Hübner, [1819].

LOMATOSTICHA Möschler, 1872, Stettin. ent. Ztg 33: 358.

Type-species: Lomatosticha nigrostriata Möschler, 1872, ibidem 33: 359, by monotypy. [AUSTRALIA]: Melbourne. Lomatosticha was established in the Notodontidae; its type-

species, L. nigrostriata, was placed in the Cnethocampinae, now Thaumetopoeidae, by Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 374, as a junior subjective synonym of Oenosandra boisduvalii Newman, 1856, Trans. ent. Soc. Lond. (2) 3 (8): 286.

See also Oenosandra Walker, 1856.

LOMELA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 33.

Type-species: *Lomela pudica* Kíriakoff, 1962, *ibidem* 114: 33, fig.41, by original designation. [ZAIRE]: Tshuapa, Likete s[ur R.] Lomela.

LOMUNA Field, 1952, Proc. U.S. natn. Mus. 102: 228. LITH Type-species: Mulona nigripuncta Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 387, pl.29, fig.6, by original designation. [PUERTO RICO]. Field, 1952, ibidem 102: 229, stated that this species was known only from Puerto Rico and that the locality, Colombia, cited for the holotype by Hampson was probably an error.

LOPERA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 99.

Type-species: *Lopera squamosa* Walker, 1855, *ibidem* **4**: 920, by monotypy. South Africa.

‡LOBERA; Spitz, 1930, Ent. Rdsch. 47: 41.

An incorrect subsequent spelling of Lopera Walker, 1855.

‡LOPEZA; misspelling. See Lobeza Herrich-Schäffer, [1854].

LOPHILEMA Aurivillius, 1910, in Sjöstedt, Wiss. Ergebn. schwed. zool. Exped. Kilimandjaro, Meru . . . 1905-1906 (9): 25.

Type-species: Lophilema sordida Aurivillius, 1910, ibidem (9): 26, fig.4, by monotypy. [TANZANIA]: [Kilimanjaro] Kilimandjaro, lower Kibonoto.

L. sordida is a junior subjective synonym of Macrosia polioplaga Hampson, 1901, Ann. Mag. nat. Hist. (7) 8: 180.

LOPHOCAMPA Harris, 1841, Rep. Insects Mass. injurious to Vegn: 258.

Type-species: Lophocampa caryae Harris, 1841, ibidem: 258, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 208. U.S.A.: Massachusetts.

LOPHOCOSMA Staudinger, 1887, in Romanoff, Mém. Lépid. 3: 220.

Type-species: *Notodonta atriplaga* Staudinger, 1887, *ibidem* 3: 220, pl.12, fig.8, by monotypy. [U.S.S.R.]: Vladivostok, Askold [Island].

Lophocosma was established as a subgenus of Notodonta Ochsenheimer, 1810.

LOPHODONTA Packard, 1864, Proc. ent. Soc. Philad. 3: 357.

Type-species: Lophodonta ferruginea Packard, 1864, ibidem 3: 357, by subsequent designation by Grote, 1895, Abh. naturw. Ver. Bremen 14: 48. [U.S.A.]: Massachusetts.

LOPHONEURA Hampson, 1894, Fauna Br. India (Moths) 2: vii, 41, 78.

Type-species: *Lophoneura uniformis* Hampson, 1894, *ibidem* 2: 78, fig.45, by original designation. [BURMA]: Tenasserim Valley.

LOPHONTOSIA Staudinger, 1892, in Romanoff, Mém. Lépid. > 6: 361.

Type-species: *Odontosia cuculus* Staudinger, 1887, *ibidem* 3: 226, pl.13 [recte 17], fig.5, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Sidemi.

Lophontosia was established as a subgenus of Lophopteryx Stephens, 1828.

LOPHOPTERYX Stephens, 1828. See Ptilodon Hübner, 1822.

LOPIENA Kiriakoff, 1961, Bull. Annls Soc. r. ent. Belg. 97: 115.

Type-species: Polienus ochracea Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 558, by original designation. [ANGOLA]: Malange.

LOUDONTA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 151 (key), 164.

Type-species: Loudonta dispar Kiriakoff, 1962 [July 31], ibidem 98 (10): 165; 1962 October 20, Bonn. zool. Beitr. 13: 220, abb.1, figs 2, 3 pl.1, fig.2 (as Pydna dispar sp.nov.), by original designation. CHINA: [Chekiang, east Tien-mu-Shan]. Additional data cited by Kiriakoff, 1962 October 20, ibidem 13: 221.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr. to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper L. dispar was described as a new species in the combination Pydna dispar, with a footnote stating that the species required a new genus which had been described in a revision of the Pydna -group; the revision, however, which was published in the Belgian journal, appeared first.

In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

LOXOBIRETA Nakamura, 1976, Ent. Rev. Japan 29: 48.

Type-species: Pydna albifusa Wileman, 1910, Entomologist 43: 345, by original designation. [TAIWAN]: Kanshirei.

LOXOMIMA Draudt, 1915, in Seitz, Gross-Schmett. Erde 6: 146.

Type-species: Loxomima imitans Draudt, 1915, ibidem 6: 146, by monotypy. [BRAZIL]: Sta Catarina, Joinville.

The date of publication of the part of Seitz containing page 146 was cited on page 153 as 30.iv.1915 in both the German and English editions.

LOXOPHLEBIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 381.

Type-species: Poecilosoma vesparis Butler, 1873, Ann. Mag. nat. Hist. (4) 12: 227, by original designation. Peru.

LOXOZONA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xviii, 29, 416.

Type-species: Josia lanceolata Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 307, by original designation. [BRAZIL]: Para.

LUCA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 256. NOTO Type-species: Luca herbida Walker, 1862, ibidem (3) 1: 256, by monotypy. [BRAZIL]: Rio de Janeiro.

LUSURA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1067.

Type-species: Lusura discalis Walker, 1855, ibidem 5: 1067, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 597. [BRAZIL]: Para.

LYCES Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 366.

Type-species: Lyces angulosa Walker, 1854, ibidem 2: 367, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 400. BRAZIL.

LYCLENE Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 300. LITH

Type-species: Cyllene humilis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 544, by subsequent designation (for Cyllene Walker, 1854) by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 426. [Burma] E. Indies: [Moulmein], cited on a data label of the holotype.

Lyclene Moore was established as the objective replacement

name for Cyllene Walker, 1854.

CYLLENE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 543.

Type-species: Cyllene humilis Walker, 1854, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus.

A junior homonym of Cyllene Griffith & Pidgeon, 1834, The animal Kingdom arranged . . . by Baron Cuvier . . . 12: 597 (but attributed to Gray, an incorrect authorship), - Mollusca. The objective replacement name is Lyclene Moore, 1858.

LYCOMORPHA Harris, 1839, Am. J. Sci. 36: 317. Type-species: Sphinx pholus Drury, 1773, Illust. nat. Hist. exot. Insects 2: index, 49, pl.28, fig.3, by monotypy. U.S.A.: New England; New York; Maryland; Virginia; and 'Carolina'.

Lycomorpha was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 522; and was placed in the Lithosiidae by Forbes, 1960, Lepid. New York and neighboring States 4: 50.

PREPODES Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 100, 101.

Type-species: Sphinx pholus Drury, 1773, by monotypy.

A junior homonym of Prepodes Schoenherr, 1823, Isis, Jena 1823: 1140, - Insecta, Coleoptera. The objective replacement name is Lycomorpha Harris, 1839.

Prepodes was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

LYCOMORPHODES Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 84, 378.

Type-species: Trichromia strigosa Butler, 1877, Trans. ent. Soc. Lond. 1877: 328, by original designation. [BRAZIL]: [Espirito Santo] Espiritu Sancto.

LYCOREA Walker, 1854. See Mallodeta Butler, 1876. CTEN

LYGROTES Butler, 1878, Trans. ent. Soc. Lond. 1878: 57.

Type-species: Lygrotes arctipennis Butler, 1878, ibidem 1878: 57, by original designation. [BRAZIL]: Amazon region. near Lago Macaco.

Lygrotes was established in the Lithosiidae, now Arctiidae.

but is now included in the Notodontidae. No published reference to the transfer has been found.

LYMANTICA Collenette, 1936, Novit. zool. 40: 153. LYMA Type-species: Lymantica epelytes Collenette, 1936, ibidem 40: 154, pl.12, fig.24, by original designation. MADAGASCAR: E., 149km E. of Tananarivo, Périnet.

LYMANTRALEX Collenette, 1938, J. fed. Malay St. Mus. 18: 263.

Type-species: *Lymantralex heteroides* Collenette, 1938, *ibidem* **18**: 264, by original designation. [MALAYSIA] Malaya: Pahang, Fraser's Hill.

LYMANTRIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 160.

Type-species: *Phalaena monacha* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 501, by subsequent designation by Moore, [1883] 1882-83, *Lepid. Ceylon* 2: 99. EUROPE.

HYPOGYMNA Billberg, 1820, Enumeratio Insect. Mus. G.J. Billberg: 84.

Type-species: *Phalaena monacha* Linnaeus, 1758, by subsequent designation by Curtis, 1839, *Br. Ent.* 16: 767.

A junior objective synonym of *Lymantria* Hübner, [1819]. Neave, 1939, *Nomencl. zool.* 2:738, cited Kirby & Spence, 1826, *Introd. Ent.* 3: 102 as the original reference for *Hypogymna*. It was however a correct subsequent usage of the name.

‡HYPOGYMNIA; [Hübner], [1825], Cat. Lépid. composent Colln feu Mr Franck: 88.

An incorrect subsequent spelling of *Hypogymna* Billberg, 1820.

PSILURA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 57. Type-species: Phalaena monacha Linnaeus, 1758, by monotypy.

A junior objective synonym of *Lymantria* Hübner, [1819]. *Psilura* was proposed again by Stephens, 1829 [June], *Nom. Br. Insects*: 40; and 1829 [July], *Syst. Cat. Br. Insects* (2): 50.

‡HYPOGYMA; Hemming, 1937 Hübner 2: 211.

An incorrect subsequent spelling of *Hypogymna* Billberg, 1820.

LYMANTRIADES Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 551.

Type-species: *Lymantriades obliqualinea* Bethune-Baker, 1911, *ibidem* (8) 7: 552, by original designation. [ANGOLA]: Gunnal.

LYMANTRIOPSIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xviii, 84, 465.

Type-species: *Nudaria lacteata* Holland, 1893, *Psyche, Camb.* 6: 412, by original designation. [GABON]: [Ogooué] Ogové R. valley.

LYMIRE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 489.

Type-species: *Lymire melanocephala* Walker, 1854, *ibidem* **2**: 490, by monotypy. Jamaica.

‡SYMIRE; Gaede, 1926, Dt. ent. Z. 1926: 130.

An incorrect subsequent spelling of Lymire Walker, 1854.

LYMPHORTA Walker, [1858] 1857, List Specimens lepid. Insects Colln Br. Mus. 12: 967 (key), 977. NOTO

Type-species: *Lymphorta unilinea* Walker, [1858] 1857, *ibidem* 12: 977, by monotypy. [Brazil]: Valley of the Amazon.

Lymphorta was established in the Hemiceridae, considered by Walker to be part of the Noctuidae, but now included in the Notodontidae; for that reason Lymphorta is here transferred

to the Notodontidae. The type-specimen of Lymphorta unilinea has not yet been found.

LYSANA Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 347.

Type-species: *Lysana plexa* Möschler, 1883, *ibidem* 32 (Abh.): 348, pl.18, fig.36, by monotypy. Surinam.

LYSCEIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 541.

Type-species: Lysceia bigutta Walker, 1854, ibidem 2: 541,

by monotypy. South Africa.

MACADUMA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1704.

Type-species: *Macaduma tortricella* Walker, 1866, *ibidem* 35: 1705, by monotypy. JAVA.

MACADUMOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxii, 571.

Type-species: *Garudinia excisa* Rothschild, 1912, *Novit. zool.* 19: 234, by original designation. New Guinea: [Irian Jaya], Snow Mts, near Oetakwa River.

MACHADOIA Rego Barros, 1956, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 134: 24.

Type-species: *Hemihyalea xanthosticta* Hampson, 1901, *Cat. Lepid. Phalaenae Br. Mus.* 3: 132, fig., by original designation. [Brazil]: N[ova] Friburgo.

MACHAERAPTENUS Schaus, 1894, Proc. zool. Soc. Lond. 1894: 228.

Type-species: Machaeraptenus ventralis Schaus, 1894, ibidem 1894: 229, by monotypy. VENEZUELA: Aroa.

‡MACHAEROPTENUS; Strand, 1919, in Wagner, Lepid. Cat. 22: 14.

An incorrect subsequent spelling of *Machaeraptenus* Schaus, 1894.

MACHAEROPHORA Hampson, 1900. See Machairophora Hampson, 1893.

‡MACHAEROPTENUS; misspelling. See Machaeraptenus Schaus, 1894.

MACHAIROPHORA Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 14, 86.

Type-species: *Machairophora fulvipuncta* Hampson, 1893, *ibidem* 9: 14, 86, pl.159, fig.10, by monotypy. [SRI LANKA] Ceylon: Maskelyia.

MACHAEROPHORA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 89, 411.

An unjustified emendation of *Machairophora* Hampson, 1893.

MACOTASA Moore, 1878, Proc. zool. Soc. Lond. 1878: 24.

Type-species: *Teulisna biplagella* Butler, 1877, *Trans. ent. Soc. Lond.* 1877: 355, by monotypy. [BORNEO]: Sarawak.

MACROBROCHIS Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 95 (key), 97. Available but without included species until [1856].

Type-species: *Macrobrochis interstitialis* Herrich-Schäffer, [1856] 1850-1858, *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1(1): wrapper, 15, 17, pl.93, fig.531, by subsequent monotypy. [INDIA] 'Ind. or'.

M. interstitialis is a junior subjective synonym of Lithosia gigas Walker, 1854, List Specimens lepid. Insects Colln Br.

1us. 2: 494.

Macrobrochis was included by Herrich-Schäffer in his 'Syn-

opsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

‡ISARES Herrich-Schäffer, [1858] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 72, 84.

The name *Isares* Herrich-Schäffer was attributed to Boisduval; it was published in the synonymy of *Macrobrochis* Herrich-Schäffer, [1856]; it was a nomen nudum, and was not subsequently treated as an available name. Under the *Code*, Article 11(d), ‡*Isares* Herrich-Schäffer is therefore not nomenclaturally available.

MACROCNEME Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 15.

Type-species: Zygaena maja Fabricius, 1787, Mantissa Insect. 2: 106, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 128. AMERICA: [tropical].

MACROCRAMBUS Kirby, 1892. See Crambomorpha Felder, 1874.

MACROHOPLITIS Roepke, 1943, Natuurh. Maandbl. 32: 79.

Type-species: *Macrohoplitis kalisi* Roepke, 1943, *ibidem* 32: 79, fig.8, by monotypy. Java: eastern, Djunggo-Ardjuno.

M. kalisi is a junior subjective synonym of Damata microsticta Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 157. See also Damatoides Matsumura, 1927.

MACROLAELIA Aurivillius, 1925, Ergebn. zweiten dt. zent.-Afr. Exped. 1910-1911 (Zool.) 1 (18): 1285. LYMA

Type-species: *Macrolaelia corticina* Aurivillius, 1925, *ibidem* (Zool.) 1: 1286, fig.69, by monotypy. [CAMEROUN] Kamerun: S., Kungulu-Urwald.

MACRONADATA Möschler, 1887, Abh. senckenb. natur-forsch. Ges. 15: 82.

Type-species: *Macronadata collaris* Möschler, 1887, *ibidem* **15**: 83, pl., fig.7, by monotypy. [Ghana] Goldküste: Aburi.

MACRONEURODES Warren, 1900, Novit. zool. 7: 127.

Type-species: *Macroneurodes albimacula* Warren, 1900, *ibidem* 7: 127, by original designation. Colombia: Bogota.

M. albimacula is a junior subjective synonym of Devara frigida Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 185.

Macroneurodes was established in the Cyllopodinae, now Geometridae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 413.

See also Euchontha Walker, 1865, and Gnatholophia Felder, 1874.

MACRONOLA Kirby, 1892. See Cyana Walker, 1854. LITH

MACRONYX Felder, 1874. See Teracotona Butler, 1878.

MACROPTILA Dognin, 1894, Annls Soc. ent. Belg. 38: 129.

Type-species: *Macroptila crinada* Dognin, 1894, *ibidem* 38: 129, by monotypy. ECUADOR: Loja.

MACROSENTA Holland, 1893, Psyche, Camb. 6: 565. NOTO Type-species: Macrosenta longicauda Holland, 1893, ibidem 6: 566, fig., by original designation. [GABON]: [Ogové] Ogooué R. valley.

Macrosenta was established in the Noctuidae; it was placed in the Notodontidae by Strand, 1912, Fauna exot. 2: 31.

MACROSHACHIA Matsumura, 1925, Zool. Mag. Tokyo 37: 395.

Type-species: *Macroshachia lineata* Matsumura, 1925, *ibidem* 37: 395, text-fig., by original designation. TAIWAN.

M. lineata is a subspecies of Notodonta nodyna Swinhoe, 1907, Ann. Mag. nat. Hist. (7) 19: 206.

See also Angustiala Bryk, 1950.

MACROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 81, 129.

Type-species: Lithosia fumeola Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 508, by original designation. [SOUTH AFRICA]: Cape Province.

MACRUROCAMPA Dyar, 1893, Ent. News 4: 34. NOTO Type-species: Phalaena marthesia Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 127, 175, pl.264, fig.B, by original designation. [U.S.A.]: Virginia.

P. marthesia is dated from the wrapper of part 22, not from

the title-page of volume 3 which is dated 1782.

MACULONACLIA Griveaud, 1964, Faune Madagascar 17: 33.

Type-species: Syntomis muscella Mabille, 1884, Annls Soc. ent. Belg. 1884 (Bull.): clxxxviii, by original designation. MADAGASCAR.

MADEMA Griveaud, 1977, Faune Madagascar 43: 250.

Type-species: Crorema viettei Collenette, 1959, Annls Soc. ent. Fr. 128: 106, pl.1, fig.7, by original designation. MADAGASCAR: E., route d'Anosibe, Km.26, foret de Sandrangato.

MAENAS Hübner, [1819]. See Paralacydes Aurivillius, 1899.

MAENOLENEURA Butler, 1878, Trans. ent. Soc. Lond. 1878: 56.

Type-species: Maenoleneura anomala Butler, 1878, ibidem 1878: 57, by original designation. [BRAZIL]: Rio Purus, Gapatiny; and Araras.

M. anomala is a junior subjective synonym of Tricypha furcata Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 655.

See also Tricypha Möschler, 1878.

MAEPHA Walker, 1864, List Specimens lepid. Insects Colln Br. Mus. 30: 1013.

Type-species: Maepha opulentana Walker, 1864, ibidem 30: 1014, by monotypy. [Brazil]: [R. Amazon], [Tefé] Ega.

‡MALPHA; Strand, 1920, in Wagner, Lepid. Cat. 24: 714. An incorrect subsequent spelling of Maepha Walker, 1864.

MAFANA Viette, 1979, Bull. mens. Soc. linn. Lyon 48: 14.

Type-species: *Mafana lajonquierei* Viette, 1979, *ibidem* 48: 15, figs 1, 19, by original designation. Holotype &, MADAGASCAR: E., environs de Périnet, foret d'Analamazaotra, (MNHN, Pario)

Mafana in the Noctuidae, Ophiderinae is cited here as it was not included in Nye, 1975, Generic Names Morhs World 1.

MAGAVA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 503.

Type-species: Magava multilinea Walker, 1865, ibidem 32:

503, by monotypy. [Honduras]: Limas.

Magava was established in the Limacodidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 584.

MAGUILA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 204.

Type-species: *Fentonia maguila* Schaus, 1928, *Proc. U.S. natn. Mus.* 73 (19): 80, by original designation. Philippines: Mindanao, Surigao.

MAHAVIRA Moore, 1878, Proc. zool. Soc. Lond. 1878: 11.

Type-species: *Mahavira flavicollis* Moore, 1878, *ibidem* 1878: 11, pl.1, fig.3, by monotypy. INDIA: Darjeeling.

MAHENSIA Fryer, 1912, Trans. Linn. Soc. Lond. (2) Zool. 15: 5.

Type-species: *Mahensia seychellarum* Fryer, 1912, *ibidem* (2) Zool. **15**: 5, pl.1, figs 2, 3, by original designation. Seychelles: Silhouette Island.

MAHOBA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 52.

Type-species: Cyclidia plagidotata Walker, [1863] 1862, List Specimens lepid. Insects Colln Br. Mus. 26: 1483, (but included by Moore as ‡plagiodotata, an incorrect subsequent spelling), by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 482. INDIA: [W. Bengal, Darjeeling], Darjiling.

MAIMAIA Matsumura, 1933, Insecta matsum. 7: 140. LYMA Type-species: Ocneria furva Leech, [1889] 1888, Proc. zool. Soc. Lond. 1888: 631, pl.31, fig.10, by original designation. JAPAN: Oiwake.

MAINIELLA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 44.

Type-species: *Mainiella subterminalis* Kiriakoff, 1962, *ibidem* **114**: 45, fig.57, by original designation. [NIGERIA]: Warri.

MALABUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 716.

Type-species: *Malabus incertus* Walker, 1855, *ibidem* 3: 716, by monotypy. [Brazil]: Para.

MALACHITIS Hampson, 1895, Trans. ent. Soc. Lond. 1895: 291.

Type-species: *Malachitis melanochlora* Hampson, 1895, *ibidem* 1895: 291, fig., by original designation. INDIA: Bhutan.

MALESIA van Eecke, 1920, Zool. Meded. Leiden 5: 132.

Type-species: *Malesia eugoana* van Eecke, 1920, *ibidem* 5: 132, fig., by monotypy. JAVA: W., Preanger and unstated locality in E. Java.

MALGADONTA Kiriakoff, 1962, Tijdschr. Ent. 105: 255.

Type-species: *Dinotodonta nigripunctella* Viette, 1955, *Bull. mens. Soc. linn. Lyon* **24**: 226, by original designation. MADAGASCAR: eastern, Perinet district, [Analamazaotra] Analamazoatra forest.

MALLOCEPHALA Blanchard, 1852, in Gay, Hist. Chile (Zool.) 7: 68.

Type-species: Mallocephala rubripes Blanchard, 1852, ibi-

dem 7: 68; 1854, Atlas Hist. Chile 2 (Ent., Lepid.): pl.4, fig.7, by monotypy. CHILE: Coquimbo.

MALLODETA Butler, 1876, J. Linn. Soc. (Zool.) 12: 398.

Type-species: Glaucopis clavata Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 192, by monotypy (of Lycorea Walker, 1854). BRAZIL.

Mallodeta was established as the objective replacement name for Lycorea Walker, 1854. Lycorea was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was retained in the Zygaenidae by Butler, 1876, ibidem 12: 398; and was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 193.

LYCOREA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 192.

Type-species: Glaucopis clavata Walker, 1854, by monotypy. A junior homonym of Lycorea Doubleday, [1847], Genera diurnal Lepid. 1: 105, – Lepid., Nymphalidae. The objective replacement name is Mallodeta Butler, 1876.

Lycorea Walker was established as a subgenus of Glaucopis

Fabricius, 1807.

MALLOSTETHUS Butler, 1876, J. Linn. Soc. (Zool.) 12: 408.

Type-species: Glaucopis metamelas Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 145, by original designation. [Brazil]: Para.

MALOCAMPA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 261 (key), 307.

Type-species: *Phalaena punctata* Stoll, 1780, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* **4**: 36, pl.307, fig.F, by original designation. SURINAM.

The type-species is dated from the wrapper of part 26, not from the title-page of volume 4 which is dated 1782.

from the title page of volume 4 which is dated 1702.

‡MALPHA; misspelling. See Maepha Walker, 1864. LITH

MALUPA Schaus, 1905, Proc. U.S. natn. Mus. 29: 245. NOTO Type-species: Malupa elongata Schaus, 1905, ibidem 29: 246, by monotypy. [GUYANA] British Guiana: Omai.

MAMALA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 472.

Type-species: *Mamala instructalis* Walker, 1865, *ibidem* 32: 472, by monotypy. [Dominican Republic]: St Domingo.

MANNINA Dyar, 1916, Proc. U.S. natn. Mus. 51: 7. ARCT Type-species: Mannina hagnoleuca Dyar, 1916, ibidem 51: 8, by original designation. MEXICO: Hidalgo, Guerrero Mill.

MANOBA Walker, [1863] 1864, J. Proc. Linn. Soc. (Zool.) 7: 62.

Type-species: *Manoba implens* Walker, [1863] 1864, *ibidem* 7: 62, by monotypy. [BORNEO]: Sarawak.

MANTALA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 108.

Type-species: *Mantala tineoides* Walker, 1862, *ibidem* **6**: 109, by monotypy. Borneo: Sarawak.

MANTRUDA Schaus, 1906, Proc. U.S. natn. Mus. 30: 94.

Type-species: *Mantruda erratica* Schaus, 1906, *ibidem* 30: 95, by monotypy. VENEZUELA: Aroa.

MANULEA Wallengren, 1863, Wien. ent. Monatschr. 7: 145, 146.

Type-species: Lithosia gilveola Ochsenheimer, 1810,

Schmett. Eur. 3: 137, by subsequent designation by Moore, 1878, Proc. zool. Soc. Lond. 1878: 18. GERMANY.

Manulea was redescribed as new by Wallengren, 1865, K.

svenska VetenskAkad. Handl. 5 (4): 43.

L. gilveola is a junior subjective synonym of Noctua unita [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 68.

MARANE Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 397.

Type-species: *Marane subargentea* Walker, 1865, *ibidem* 32: 397, by monotypy. AUSTRALIA: northern.

M. subargentea is a junior subjective synonym of Teara

argentata Walker, 1865, ibidem 32: 355.

Marane was established in the Liparidae, now Lymantriidae; it was placed in the Eupterotidae by Turner, 1903, Trans. R. Soc. S. Aust. 27: 23, and by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497; Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 368, placed it in the Notodontidae, subfamily Cnethocampinae, now Thaumetopoeidae, and its placing in this family was confirmed by Kiriakoff, 1969, Bull. Annls Soc. r. ent. Belg. 105: 279.

MARBLA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 436.

Type-species: *Eloria divisa* Walker, 1855, *List Specimens lepid. Insects Colln Br. Mus.* 4: 815, by original designation. Type-locality: not known, [West Africa].

MARBLEPSIS Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 158.

Type-species: *Marblepsis pirgula* Hering, 1926, *ibidem* 14: 158, pl.22, row h, by subsequent designation by Bryk, 1934, *in* Strand, *Lepid. Cat.* 62: 303. WEST AFRICA: Kwai.

The date of publication of the part of Seitz containing page 158 was cited on page 153 as 5.xi.1926 in both the German and

the English editions.

MARBLOIDES Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 157.

Type-species: *Marbloides paradoxa* Hering, 1926, *ibidem* 14: 157, pl.22, row i, by original designation. [IVORY COAST] Elfenbeinküste.

The date of publication of the part of Seitz containing page 157 was cited on page 153 as 5.xi.1926 in both the German and the English editions.

MARCELLA Dognin, 1914, Hétérocères nouv. Am. Sud 7: 28.

Type-species: *Marcella volcancita* Dognin, 1914, *ibidem* 7: 28, by original designation. Colombia: Quindin, Volcancito.

MARCIPA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 807.

NOCTUIDAE

Type-species: Marcipa inscripta Walker, 1855, ibidem 4:

808, by monotypy.

Marcipa was established in the Liparidae, now Lymantridae; it was transferred from the Liparidae, but was not placed elsewhere, by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497; and was placed in the Noctuidae, 'Noctuinae' [sensu Hampson] now Ophiderinae, by Hampson, 1926, Descr. new Genera Species Lepid. Phalaenae Subfamily Noctuinae Br. Mus.: 161.

MARCIPALINA Pelletier, 1978, Mém. Mus. Hist. nat. Paris (N.S.) A 110: 10.

NOCTUIDAE

Type-species: *Marcipa violacea* Pelletier, 1974, *Lambillionea* 74: 53, figs, by original designation. Holotype &, GABON: Belinga, camp central, 700m, (MNHN, Paris).

Marcipalina in the Noctuidae, Ophiderinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

MARDARA Walker, 1865. See Ramadra Nye nom. n. LYMA

MARECIDIA Schaus, 1901, Jl N.Y. ent. Soc. 9: 43. CTEN Type-species: Marecidia sanguipuncta Schaus, 1901, ibidem 9: 43, by monotypy. BRAZIL: Sta Catarina.

MARGAROPSECAS Kiriakoff, 1964, Bonn. zool. Beitr. 14: 289.

Type-species: *Margaropsecas margarethae* Kiriakoff, 1964, *ibidem* 14: 290, abb.6, fig.61, pl.5, fig.54, by original designation. China: north Yunnan, Li-Kiang.

MARISSA Walker, 1854. See Eunomia Hübner, 1818.

MARMAX Rafinesque, 1815, Analyse Nat.: 128. ZYGAENIDAE Type-species: Zygaena argynnis Fabricius, 1781, Species Insect. 2: 161, by subsequent designation (for Charidea Dalman, 1816) by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 169.

Z. argynnis is a junior subjective synonym of Sphinx hypparchus Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 7, pl.197, fig.C.

Marmax was established as the objective replacement name for Glaucopis Fabricius, 1807.

GLAUCOPIS Fabricius, 1807, Magazin Insektenk. (Illiger) 6: 289.

Type-species: Zygaena argynnis Fabricius, 1781, by subsequent designation (for Charidea Dalman, 1816) by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 169.

A junior homonym of *Glaucopis* Gmelin, 1788, *in* Linnaeus, *Syst. Nat.* (Edn 13) 1: 363, – Aves. The objective replacement name is *Marmax* Rafinesque, 1815.

Glaucopis Fabricius, 1807, was placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1843, by the International Commission on Zoological Nomenclature, 1966, Bull. zool. Nom. 23 (Opinion 789): 214.

CHARIDEA Dalman, 1816, K. svenska VetenskAkad. Handl. 1816: 225.

Type-species: Zygaena argynnis Fabricius, 1781, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 169, but cited as ‡hyparchus, an incorrect subsequent spelling of Sphinx hypparchus Cramer, 1779, a nominal species not originally included in Glaucopis Fabricus, 1807. Kirby, however, placed hypparchus as the senior synonym of Z. argynnis, a nominal species which was originally included in Glaucopis Fabricius, 1807. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species.

Charidea was established, unnecessarily, as an objective replacement name for Glaucopis Fabricius, 1807, and is a junior objective synonym of Marmax Rafinesque, 1815.

Charidea has been used in the Ctenuchidae following Butler, 1876, J. Linn. Soc. Lond. (Zool.) 12: 415, who placed in Charidea several nominal species which with one exception are currently placed in Cyanopepla Clemens, 1861.

MARSYPOPHORA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.28.

Type-species: Marsypophora erycinoides Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.28, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

MARTHANTIA Schaus, 1923, Ann. Carnegie Mus. 15: 83.

Type-species: *Marthantia stelligera* Schaus, 1923, *ibidem* 15: 83, pl.4, fig.6, by original designation. French Guiana: Oyapok River, Pied Saut.

MARTHULA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 9: 154 (key), 164.

Type-species: Marthula quadrata Walker, 1856, ibidem 9:

164, by monotypy. [Brazil]: Para.

Marthula was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 275.

MARUSHACHIA Matsumura, 1934, Insecta matsum. 8: 153, 154.

Type-species: *Marushachia rotundata* Matsumura, 1934, *ibidem* 8: 153, fig.2, by original designation. TAIWAN: Baibara.

M. rotundata is a junior subjective synonym of Cleapa latifascia formosae Strand, 1920, Arch. Naturgesch. 84 (A) 12: 186.

See also Cleapa Walker, 1855.

MASCHANE Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 27: 2.

Type-species: *Maschane erratipennis* Walker, 1863, *ibidem* 27: 3, by subsequent designation by Schaus, 1901, *Trans. ent. Soc. Lond.* 1901: 317. [BRAZIL]: Para.

Maschane was established in the Crambites, now Pyralidae; it was placed in the Notodontidae by Schaus, 1901, ibidem 1901: 317.

MASOANDRO Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 228.

Type-species: *Mardara peculiaris* Butler, 1879, *Ann. Mag. nat. Hist.* (5) **4**: 240, by original designation. MADAGASCAR: Antananarivo.

MASSICYTA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1712.

Type-species: *Massicyta fusca* Walker, 1856, *ibidem* 7: 1712, by monotypy. Brazil.

MASTIGOCERA Harris, 1839. See Horama Hübner, [1819].

MATARUM Viette, [1976] 1975, Bull. Soc. ent. Fr. 80: 215.
NOCTUIDAE

Type-species: *Matarum etiennei* Viette, [1976] 1975, *ibidem* **80**: 216, figs 1, 2, by original designation. Holotype & Réunion: foret de Bébour, (MNHN, Paris).

Matarum in the Noctuidae, Acronictinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

MAXIA Kirby, 1892. See Phryganopteryx Saalmüller, 1884.

MAZAERAS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 632. ARCT

Type-species: *Mazaeras conferta* Walker, 1855, *ibidem* 3: 633, by monotypy. BRAZIL.

MECISTORHABDIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 29. THYRET

Type-species: *Metarctia haematoessa* Holland, 1893, *Psyche, Camb.* 6: 396, by original designation. [GABON]: [Ogooué] Ogové R. valley.

MEDAMA Matsumura, 1931, 6000 Illust. Insects Japan-Empire: 717.

Type-species: *Medama basimaculatum* Matsumura, 1931, *ibidem*: 717, fig., by monotypy. [TAIWAN] Formosa: Niitaka, Tamaho.

M. basimaculatum is a junior subjective synonym of Euproctis diplaga Hampson, 1910, J. Bombay nat. Hist. Soc. 20: 113.

Medama was again proposed as a new genus by Matsumura,

1933, Insecta matsum. 7: 140, and Euproctis diplaga Hampson, 1910, cited as the type-species.

MEDANELLA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 401.

Type-species: *Medanella subterminalis* Kiriakoff, 1974, *ibidem* 17: 401, text-fig.20, pl.4, fig.2, by original designation. SUMATRA: northern, Dolok Merangir.

MEDLERANA Laporte, 1973, Annls Fac. Sci. Univ. Cameroun 14: 110.

NOCTUIDAE

Type-species: *Medlerana nigeriensis* Laporte, 1973, *ibidem* 14: 111, fig.1, by original designation. Holotype & NIGERIA: Obudu, (MNHN, Paris).

MEGACERAMIS Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 126 (key), 167.

Type-species: *Megaceramis lamprolepis* Hampson, [1893] 1892, *ibidem* 1: 167, fig.103, by original designation. [INDIA]: Sikkim.

MEGALOBOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxii, 587.

Type-species: *Garudinodes recurviloba* Rothschild, 1912, *Novit. zool.* 19: 232, by original designation. New Guinea: [Papua New Guinea], Mambare R., Biagi.

MEGANACLIA Aurivillius, 1892, Ent. Tidskr. 13: 190.

THYRET

Type-species: *Naclia sippia* Plötz, 1880, *Stettin. ent. Ztg* **41**: 78, by monotypy. [CAMEROUN]: [mountains] Cameroons-Gebirge.

MEGAPISA Aurivillius, 1904, Ark. Zool. 2 (4): 29. THYRET Type-species: Megapisa nigripennis Aurivillius, 1904, ibidem 2 (4): 30, by monotypy. [AFRICA] Centralafrika.

Megapisa was established in the Syntomidae, now Ctenuchidae; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 9.

MEGASHACHIA Matsumura, 1929, Insecta matsum. 4: 36, 37.

Type-species: *Megashachia takamukuana* Matsumura, 1929, *ibidem* 4: 36, pl.1, fig.2, by original designation. Talwan: Horisha, Kyuhabon near Horisha.

M. takamukuana is a subspecies of Crino fulgurifera Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 14: 1347.

MEKRANIA Brandt, 1941, Mitt. münch. ent. Ges. 31: 856.

NOCTUIDA

Type-species: *Mekrania punctalis* Brandt, 1941, *ibidem* 31: 856, pl.27, fig.55, by monotypy. Syntypes 18, 19, IRAN: [Port Chah Bahar] Bender Tchahbahar.

Mekrania in the Noctuidae, Hypeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

MELAGONA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 635.

Type-species: *Melagona dentatus* Gaede, 1930, *ibidem* 10: 635, pl.84 row e (as pl.84 row c in error in the English edition), by monotypy. [Sulawesi] Celebes: northern.

The date of publication of the part of Seitz containing page 635 was cited on page 633 as 21.ii.1930 in the German edition, and as 17.v.1930 in the English edition.

MELAGONINA Gaede, 1933, in Seitz, Gross-Schmett. Erde 2 (Suppl.): 182.

Type-species: *Melagonina hoenei* Gaede, 1933, *ibidem* 2 (Suppl.): 182, pl.14 row f (but published as ‡*hoehni*, an incorrect original spelling, in the legend), by monotypy. [CHINA]: Kiangsu, Nanking.

The date of publication of the part of Seitz containing page 182 was cited on page 177 as 22.iii.1933 in both the German and English editions.

‡MELAINA Zikan, 1968, in Zikan & Zikan, Pesquisa Agropec. bras. 3: 95.

‡Melaina is unavailable under the Code, Article 13(a)(b); it was neither described nor accompanied by the fixation of a type-species. One of the two originally included species (jonesi Rothschild) is available but its identity is uncertain. Rothschild described two species of Ctenuchidae named jonesi: Aclytia jonesi Rothschild, 1912, Novit. zool. 19: 159, and Ctenucha jonesi Rothschild, 1912, ibidem 19: 184.

‡MELALOPHA Hübner, [1806]. See Clostera Samouelle, 1819.

MELALOPHA Hübner, 1822. See Clostera Samouelle, 1819.

MELANAEMA Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 397. LITH

Type-species: *Melanaema venata* Butler, 1877, *ibidem* (4) **20**: 397, by monotypy. JAPAN: Yokohama.

MELANARCTIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 33.

Type-species: *Automolis ockendeni* Rothschild, 1910, *Novit. zool.* **16**: 40, pl.5, fig.41, by original designation. PERU: Inambari River, La Oroya.

MELANAREAS Butler, 1889, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 7: 4, 29. ARCT

Type-species: *Euprepia imperialis* Kollar, [1844] 1848, *in* Hügel, *Kaschmir und das Reich der Siek* 4: 466, pl.21, fig.1, by monotypy. [INDIA]: [Mussooree] Masuri.

E. imperialis was established in the nominal genus Euprepia Agassiz, 1846, an unjustified emendation of Eyprepia Ochsenheimer, 1810.

nemier, 1010.

MELANIA Wallengren, 1863. See Siccia Walker, 1854.

MELANONACLIA Griveaud, 1964, Faune Madagascar 17: 66.

Type-species: *Naclia luctosa* Oberthür, 1911, *Annls Soc. ent.* Fr. 79: [467], by original designation. MADAGASCAR: Fénérive.

MELASTROTA Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 439.

Type-species: *Pseudoblabes nigrisquamata* Swinhoe, 1901, *ibidem* (7) 7: 467, by original designation. [AUSTRALIA]: Queensland, Coomoo.

MELEBAEAS Distant, 1903, Insecta Transv. 1: 93. NOTO Type-species: Hoplitis gigas Distant, 1899, Ann. Mag. nat. Hist. (7) 3: 463, by original designation. [SOUTH AFRICA]: Transvaal, Lydenburg district.

H. gigas is a junior subjective synonym of Amyops ingens Karsch, 1895, Ent. Nachr. Berlin 21: 362, pl.3, fig.4.

See also Amyops Karsch, 1895.

‡MELEBAEUS; Gaede, 1934, in Strand, Lepid. Cat. 59: 137. An incorrect subsequent spelling of Melebaeas Distant, 1903.

‡MELEBAEUS; misspelling. See Melebaeas Distant, 1903.

MELESE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 491. ARCT

Type-species: *Melese postica* Walker, 1854, *ibidem* 2: 492, by monotypy. Brazil: Para.

‡MELESSE; Rothschild, 1910, Novit. zool. 16: 48. An incorrect subsequent spelling of Melese Walker, 1854.

‡MELISE; Rothschild, 1922, Ann. Mag. nat. Hist. (9) 9: 478. An incorrect subsequent spelling of Melese Walker, 1854.

‡HETEROMELESE Zikan, 1968, in Zikan & Zikan, Pesquisa Agropec. bras. 3: 90.

Heteromelese is undescribed and unavailable under the Code, Article 13(a). The only included nominal species, Neritos babosa Dognin, 1894, is currently considered to be congeneric with the type-species of Melese Walker, 1854.

‡MELESSE; misspelling. See Melese Walker, 1854. ARCT

MELGONA Nye nom. n. for Coniortodes Collenette, 1955.

Type-species: Coniortodes conia Collenette, 1955, Ann. Mag. nat. Hist. (12) 8: 644, pl.16, fig.13, by original designation (for Coniortodes Collenette, 1955). KENYA: Mt Elgon.

CONIORTODES Collenette, 1955, Ann. Mag. nat. Hist. (12) 8: 643.

Type-species: Coniortodes conia Collenette, 1955, by original designation.

A junior homonym of *Coniortodes* Wagner, 1952, in Dahl, *Tierwelt Dtl.* 41: 172, 187, – Hemiptera. The objective replacement name is *Melgona* Nye nom. n.

MELIA Walker, 1855. See Ilema Moore, [1860]. LYMA

MELIA Neumoegen, 1892. See Eumelia Neumoegen, 1893.

MELISA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 264.

THYRET

Type-species: *Euchromia diptera* Walker, 1854, *ibidem* 1: 265, by monotypy. Congo.

Melisa was established as a subgenus of Euchromia Hübner, [1819], in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 152; and placed in the Thyretidae by Kiriakoff, 1960, Genera Insect. 214 (E): 5.

‡MELISE; misspelling. See Melese Walker, 1854. ARCT

MELISOIDES Strand, 1912, Arch. Naturgesch. 78 (A) 6: 192.

Type-species: *Melisoides lobata* Strand, 1912, *ibidem* 78 (A) 6: 193, by original designation. [Cameroun or Equatorial Guinea] Süd-Kamerun or Spanisch Guinea.

MELLAMASTUS Rego Barros, 1959, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 190: 1.

Type-species: *Amastus nero* Weymer, 1907, *Dt. ent. Z. Iris* **20**: 27, pl.2, fig.3, by original designation. BRAZIL: mountains near Joinville.

MELLONA van Eecke, 1926, Zool. Meded. Leiden 9: 260 (key). Nomenclaturally available but without included species until van Eecke, 1927, ibidem 10: 94.

Type-species: *Mellona blanda* van Eecke, 1927, *ibidem* 10: 95; 1926, *ibidem* 9: pl.4, fig.7 (without legend), by original designation. Sumatra: Fort de Kock.

MELORA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 773.

Type-species: *Melora amygdaloides* Walker, 1855, *ibidem* 3: 773, by monotypy. [Brazil]: Rio [de] Janeiro.

The type material of *M. amygdaloides* Walker is lost. Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 204, has been followed in his family placement of this species.

MENAPIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 461.

Type-species: *Menapia xanthophila* Walker, 1865, *ibidem* 32: 462, by monotypy. [INDIA]: North Hindostan.

M. xanthophila is a junior subjective synonym of Bireta longivitta Walker, 1856, ibidem 7: 1754.

See also Bireta Walker, 1856.

MENECLIA Grünberg, 1910, Denkschr. med.-naturw. Ges. Jena 16: 122.

Type-species: *Meneclia pallidula* Grünberg, 1910, *ibidem* 16: 123, by monotypy. [Namibia] South [West] Africa: [Windhoek] Windhuk.

MENEGITES Kiriakoff, 1954, Revue Zool. Bot. afr. 50: 182.

ARCT

Type-species: *Menegites nivea* Kiriakoff, 1954, *ibidem* **50**: 182, pl.2, fig.19, by original designation. [ZAIRE]: [Katanga], [Lubumbashi] Elizabethville.

‡MENINGOCERA; misspelling. See Meringocera Felder, 1874.

MENNIS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 379. GEOMETRIDAE

Type-species: *Mennis orislinea* Walker, 1854, *ibidem* 2: 379, by subsequent designation by Fletcher, 1954, *Zoologica, N.Y.* 39: 153. [COLOMBIA]: Santa Fé de Bogota.

Mennis was established in the Lithosiidae, now Arctiidae; it was transferred to the Geometridae by Druce, 1901, Ann. Mag. nat. Hist. (7) 7: 75, but cited as ‡Menis, an incorrect subsequent spelling.

MERAGISA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 262 (key), 314.

Type-species: Orthosoma valdiviesoi Dognin, 1890, Naturaliste 12: 193, by original designation. [ECUADOR]: San Francisco, près Loja.

MERINGOCERA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.101, fig.11. ARCT

Type-species: *Meringocera plutonica* Felder, 1874, *ibidem* 2 (Abt.2): pl.101, fig.11, by monotypy. [INDONESIA]: Molucc[as], cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 6.

‡MENINGOCERA; Pagenstecher, 1888, Jb. nassau. Ver. Naturk. 41: 118.

An incorrect subsequent spelling of Meringocera Felder, 1874.

MESAESCHRA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 273.
NOTO

Type-species: *Mesaeschra senescens* Kiriakoff, 1964, *ibidem* 14: 273, abb.3, fig.30, abb.4, fig.31 pl.3, fig.31, by original designation. China: Fukien, Kuatun.

‡MESENOCHROA Boisduval, 1870. See Mesenochroa Felder, 1874. PERI

MESENOCHROA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.30.

Type-species: *Mesenochroa guatemalteca* Felder, 1874, *ibidem* **2** (Abt.2): pl.105, fig.30, by monotypy. GUATEMALA: cited by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 6.

Felder & Rogenhofer in their Erklärung attributed *Mesenochroa* to Boisduval, 1870 (an incorrect authorship).

‡MESENOCHROA Boisduval, 1870, Considérations Lépid. Guatemala: 94 (but attributed to Felder, an incorrect authorship).

A nomen nudum first published in the synonymy of Pyralopsis

Boisduval, 1870, *ibidem*: 94 (but attributed to Felder, an incorrect authorship) and unavailable under the *Code*, Article 11(d); it was without a description and has not subsequently been treated as an available name dating from Boisduval, 1870.

The originally included nominal species ‡guatemalteca was not described by Boisduval, and was attributed to Felder, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.30; but this plate was not published until November 1874. It is probable that Boisduval had obtained a proof copy of Felder & Rogenhofer's plate and legend prior to its publication in Vienna (see Fletcher, 1977, in Nye, Generic Names Moths World 3).

MESOCEREA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xv, 302.

Type-species: *Teucer apicalis* Rothschild, 1911, *Novit. zool.* **18**: 42, by original designation. Surinam: Maroewym Valley, Aroewarwa Creek.

MESODONTA Matsumura, 1920. See Peridea Stephens, 1828.

MESODREPTA Turner, 1924, Proc. Linn. Soc. N.S.W. 49: 401.

Type-species: *Mesodrepta harpotoma* Turner, 1924, *ibidem* 49: 401, by monotypy. AUSTRALIA: Queensland, Toowoomba.

MESOLASIA Hampson, 1898. See Aethria Hübner, [1819].

MESONACLIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 20, 21. THYRET

Type-species: *Meganaclia minor* Hampson, 1914, *Cat. Lepid. Phalaenae Br. Mus.* (Suppl.) 1: 61, by original designation. UGANDA: Entebbe.

MESONADATA Kiriakoff, 1960, Revue Zool. Bot. afr. 61: 63.

Type-species: *Mesonadata quinquemaculata* Kiriakoff, 1960, *ibidem* 61: 63, fig.16, by original designation. [ZAIRE]: Uele, Paulis.

MESOPHALERA Matsumura, 1920, Zool. Mag. Tokyo 32: 150.

Type-species: *Phalera sigmata* Butler, 1877, *Ann. Mag. nat. Hist.* (4) 20: 473, by monotypy. JAPAN: Yokohama, Hakodate.

MESOSTAUROPUS Nakamura, 1974, Trans. lepid. Soc. Japan 25: 115.

Type-species: Stauropus sikkimensis Moore, 1866, Proc. zool. Soc. Lond. 1865: 811, pl.43, fig.5, by original designation. [INDIA]: Darjeeling.

Mesostauropus was established as a subgenus of Stauropus Germar, 1812.

MESOTHEN Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 403.

Type-species: *Mesothen ignea* Druce, 1898 May, *ibidem* (7) 1: 40, by monotypy. [VENEZUELA]: Mérida.

Mesothen probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson, 1898 [December], as the name for a new genus.

MESOTHEN Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xii, 25, 212.

Type-species: *Dycladia pyrrha* Schaus, 1889, *Entomologica am.* 5: 89, by original designation. MEXICO: Vera Cruz State, Paso de San Juan.

A junior homonym of *Mesothen* Druce, 1898 May. There is no objective replacement name but the type-species of *Mesothen* Hampson, 1898 [December] is currently considered

to be congeneric with the type-species of *Mesothen* Druce, 1898 May. The latter is therefore available for use as a subjective replacement name.

MESOTHEN Hampson, 1898 [December]. See Mesothen Druce, 1898 May.

METACRIAS Meyrick, 1887, Proc. Linn. Soc. N.S.W. (2) 1: 93, 749.

Type-species: Metacrias erichrysa Meyrick, 1887, ibidem (2) 1: 749, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 263. New Zealand: Mount Arthur.

METACRISIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xv, 7, 431.

Type-species: *Purius courregesi* Dognin, 1891, *Naturaliste* 13: 257, by original designation. [ECUADOR]: near Loja.

METACRISIODES Dyar, 1916, Proc. U.S. natn. Mus. 51: 8.

Type-species: *Metacrisiodes pua* Dyar, 1916, *ibidem* 51: 8, by original designation. MEXICO: Hidalgo, Guerrero Mill.

METACROCEA Dyar, 1905, Proc. U.S. natn. Mus. 29: 173.

Type-species: *Metacrocea postflava* Schaus, 1905, *ibidem* 29: 193, by original designation (in anticipation of Schaus's description of *postflava*). BRAZIL: Rio [de] Janeiro.

Metacrocea was available but was without an available included species between its date of publication 1905 October 3rd and the date of publication of its type-species by Schaus, 1905 October 11th.

METAGYLLA Hampson, 1907, Ann. Mag. nat. Hist. (7) 19: 231.

Type-species: *Metagylla miroides* Hampson, 1907, *ibidem* (7) 19: 231, by original designation. FRENCH GUIANA: St Jean Maroni.

METALLOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xx, 86, 555.

Type-species: *Metallosia chrysotis* Hampson, 1900, *ibidem* 2: 555, fig.408, by original designation. BRAZIL: [Espirito Santo] Espiritu Santo; and Rio [de] Janeiro.

METALOBA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xv, 26, 283.

Type-species: Calonotos nana Druce, 1897, Ann. Mag. nat. Hist. (6) 20: 300, by original designation (but cited as ‡Calanotos, an incorrect subsequent spelling by Hampson). Colombia: San Martin, L[l]anos of the Rio Meta.

METALOBOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 86, 373.

Type-species: Odozana cuprea Schaus, 1896, Jl N.Y. ent. Soc. 4: 150, by original designation. Brazil: São Paulo.

METAMICROPTERA Hulstaert, 1923, Revue zool. afr. 11: 408.

Type-species: *Metamicroptera rotundata* Hulstaert, 1923, *ibidem* 11: 409, by original designation. [ZAIRE]: [Katanga], [Lubumbashi] Elizabethville.

Metamicroptera when established was not placed in a family; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 8.

See also Neobalacra Kiriakoff, 1952.

‡MICROMETAPTERA; Kiriakoff, 1960, Genera Insect. 214 (E): 3, 5, 62.

An incorrect subsequent spelling of *Metamicroptera* Hulstaert, 1923.

METAMYA Travassos, 1946, Papéis Dep. Zool. S. Paulo 7: 260.

Type-species: *Paramya picta* Druce, 1898, *Ann. Mag. nat. Hist.* (7) 1: 405, by monotypy (of *Paramya* Druce, 1898 May). French Guiana: Cayenne.

Metamya was established as the objective replacement name for Paramya Druce, 1898 May.

PARAMYA Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 405. Type-species: Paramya picta Druce, 1898, by monotypy.

A junior homonym of *Paramya* Conrad, 1860, *Proc. Acad. nat. Sci. Philad.* **1860**: 232, – Mollusca. The objective replacement name is *Metamya* Travassos, 1946.

The name *Paramya* probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson, 1898 [December], as the name for a new genus.

PARAMYA Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xii, 25, 164.

Type-species: *Paramya intersecta* Hampson, 1898, *ibidem* 1: 164, by original designation. [Brazil]: Para, Amazon region.

A junior homonym of *Paramya* Conrad, 1860, and of *Paramya* Druce, 1898 May. There is no objective replacement name but the type-species of *Paramya* Hampson, 1898 [December], is currently considered to be congeneric with the type-species of *Metamya* Travassos, 1946. The latter is therefore available for use as a subjective replacement name.

METAPICONOMA Rothschild, 1910, Novit. zool. 17: 444.

THYRET

Type-species: *Metapiconoma rattrayi* Rothschild, 1910, *ibidem* 17: 444, by PRESENT DESIGNATION. UGANDA: Entebbe.

Metapiconoma was established in the Syntomidae, now Ctenuchidae; its type-species was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 14.

METARCTIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 769.

Type-species: *Metarctia rufescens* Walker, 1855, *ibidem* 3: 769, by monotypy. [South Africa]: Caffraria; and [Durban] Port Natal.

Metarctia was established in the Arctiidae; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 30, 35.

‡METARETIA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling of Metarctia Walker, 1855.

METARCTINA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 417.

Type-species: *Metarctina ochricostata* Gaede, 1928, *ibidem* 14: 417, pl.69 row i, by monotypy. [Tanzania]: Tanganyika, western.

The date of publication of the part of Seitz containing page 417 was cited on that page as 29.iii.1928 in both the German and the English editions.

#METARETIA; misspelling. See Metarctia Walker, 1855.

METAREVA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 86, 216.

Type-species: *Metareva aenescens* Hampson, 1900, *ibidem* 2: 216, fig.133, by original designation. BOLIVIA: Chaco.

METARHODIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 36.

THYRET

Type-species: *Metarctia rubripuncta* Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 147, pl.5, fig.4. by original designation. [GABON] Gaboon.

Metarhodia was established as a subgenus of Metarctia Walker, 1855.

METASCHALIS Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 125 (key), 158.

Type-species: Celeia disrupta Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 62, by original designation. INDIA: Darjeeling.

METASCRANCIA Kiriakoff, 1970, Ann. Transv. Mus. 26 (9): 191.

Type-species: *Metascrancia jansei* Kiriakoff, 1970, *ibidem* **26** (9): 191, fig.11, by original designation. South Africa: [Natal], Durban.

METASOMERA Matsumura, 1925, Zool. Mag. Tokyo 37: 396.

Type-species: *Metasomera plagifera* Matsumura, 1925, *ibidem* 37: 396, pl.6, fig.2, by original designation. TAIWAN.

M. plagifera is a subspecies of Chadisra bipars Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 82.

See also Chadisra Walker, 1862.

METASTATIA Butler, 1876, Cistula ent. 2: 115. CTEN Type-species: Hyelosia pyrrhorhoea Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 12, pl.[8], figs 43, 44, by original designation. BRAZIL.

‡HYPOCRITA Hübner, 1808, Erste Zuträge Samml. exot. Schmett.: 4.

Included in a work rejected for nomenclatural purposes by Int. Commn Zool. Nom., 1966, Bull. zool. Nom. 23 (Opinion 789): 213. Placed on the Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1860.

Of the two originally included species ‡Hypocrita chrysorrhoea Hübner, 1808, ibidem: 4, was subsequently established

as Hyelosia pyrrhorhoea Hübner, 1818.

The name *Hypocrita* was established a year earlier by Hübner, [1807], *Samml. exot. Schmett.* 1: pl.[187], – Arctiidae, Pericopinae.

METAXANTHIA Druce, 1899, Ann. Mag. nat. Hist. (7) 3: 465.

Type-species: *Metaxanthia vespiformis* Druce, 1899, *ibidem* (7) 3: 465, by monotypy. [Brazil]: [Amazonas], Villa Nova.

Metaxanthia probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson, 1901, as a name for a new genus.

METAXANTHIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: x, 5, 172.

Type-species: Metaxanthia vespiformis Druce, 1899, by original designation.

A junior homonym of *Metaxanthia* Druce, 1899. The objective replacement name is *Metaxanthia* Druce, 1899.

‡METOXANTHIA; Druce, 1905, Ann. Mag. nat. Hist. (7) 15: 467.

An incorrect subsequent spelling of *Metaxanthia* Druce, 1899.

METAXANTHIA Hampson, 1901. See Metaxanthia Druce, 1899.

METERYTHROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 85, 242.

Type-species: *Tuina sangala* Druce, 1885, *Biologia cent.-am*. (Zool.) Lepid.-Heterocera 1: 122, pl.12, fig.12, by original designation. Guatemala: Panima, Panama: Chiriqui volcano.

METEUGOA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 91, 335.

Type-species: Eugoa ochrivena Hampson, 1898, J. Bombay

nat. Hist. Soc. 11: 439, by original designation. [INDIA]: Khasia Hills.

METEUPROCTIS Matsumura, 1927, J. Coll. Agric. Hok-kaido imp. Univ. 19: 44.

Type-species: *Meteuproctis tomponis* Matsumura, 1927, *ibidem* 19: 44, by original designation. [TAIWAN] Formosa: near Niitaka, Tompo.

METEURA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 81, 123.

Type-species: Scoliacma cervina Lucas, 1890, Proc. Linn. Soc. N.S.W. (2) 4: 1068, by original designation. AUSTRALIA: Brisbane.

METEXILISIA Toulgoët, 1958, Mém. Inst. scient. Madagas-car (E) 9: 197.

Type-species: *Metexilisia citrago* Toulgoët, 1958, *ibidem* (E) **9**: 197, text-fig.16, pl.2, fig.13, by original designation. MADAGASCAR: E., Perinet, Analamazaotra forest.

METHYPSA Butler, 1875, Trans. ent. Soc. Lond. 1875: 324. IMMIDAE

Type-species: *Hypsa saturata* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 217, by original designation. Type-locality: not stated by Walker. [Malaysia].

METHYSIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 397.

Type-species: Glaucopis notabilis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 144, by original designation. [Brazil]: Para.

METOPOLOPHOTA Bethune-Baker, 1909, Ann. Mag. nat. Hist. (8) 3: 424.

Type-species: *Metopolophota epinephela* Bethune-Baker, 1909, *ibidem* (8) 3: 424, by original designation. [ZAIRE]: Ituri Forest, between Beni and Makala.

METOPTERYX Fletcher nom. n. for Taeniopteryx Janse, 1920.

Type-species: *Taeniopteryx cinerea* Janse, 1920, *Ann. Transv. Mus.* 7: 212, pl.8, figs 14-19, pl.14, fig.9, by original designation (for *Taeniopteryx* Janse, 1920). South Africa: [Natal], New Hanover.

TAENIOPTERYX Janse, 1920, Ann. Transv. Mus. 7: 212.

Type-species: *Taeniopteryx cinerea* Janse, 1920, by original designation.

A junior homonym of *Taeniopteryx* Pictet, 1842, *Hist. nat. Insectes Névroptères* (Perlides): 343, — Insecta, Plecoptera. The objective replacement name is *Metopteryx* Fletcher nom. n.

‡METOXANTHIA; misspelling. See Metaxanthia Druce, 1899.

METRIAESCHRA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 270.

Type-species: *Metriaeschra apatela* Kiriakoff, 1964, *ibidem* **14**: 270, abb.3, fig.26, pl.3, fig.26, by original designation. CHINA: north Yunnan, Li-Kiang.

METRIOPHYLA Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: x, 43. CTEN

Type-species: *Charidea apicalis* Herrich-Schäffer, [1854 December 31] 1850-1858, *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.45, fig.236, by original designation. VENEZUELA.

C. apicalis is a junior subjective synonym of Euchromia albiplaga Walker, 1854 [February 11], List Specimens Lepid. Insects Colln Br. Mus. 1: 218.

METULA Kiriakoff, 1958. See Epimetula Kiriakoff, 1964.

MEVANIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 442.

Type-species: *Mevania quadricolor* Walker, 1854, *ibidem* 2: 443, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 163. VENEZUELA.

MICCOTROGIA Toxopeus, 1948, Treubia 19: 475. LYMA Type-species: Euproctis tamsi Collenette, 1932, Novit. zool. 38: 68, pl.2, fig.11 (but cited in text as fig.38), by original designation. [MALAYSIA]: Selangor, Bukit Kutu.

MICHELLIANA Laporte, 1976, Bull. mens. Soc. linn. Lyon 45: 135.

Type-species: *Michelliana afroalpina* Laporte, 1976, *ibidem* **45**: 135, fig. genitalia 1, by monotypy. Holotype & ETHIOPIA: environs du Mont Batu, (MNHN, Paris).

Michelliana in the Noctuidae, Hadeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡MICRAGA; misspelling. See Micragrella Watson nom. n. CTEN

MICRAGRA Hampson, 1898. See Micragrella Watson nom.

MICRAGRELLA Watson nom. n. for Micragra Hampson, 1898.

Type-species: *Micragra sanguiceps* Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 466, fig. 252, by original designation (for *Micragra* Hampson, 1898). BRAZIL: Iguarasu.

MICRAGRA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xix, 29, 466.

Type-species: *Micragra sanguiceps* Hampson, 1898, by original designation.

A junior homonym of *Micragra* Chandoir, 1872, *Annls Soc.* ent. Belg. 15: 155, – Insecta, Coleoptera. The objective replacement name is *Micragrella* Watson nom. n.

‡MICRAGA; Hampson, 1901, Ann. Mag. nat. Hist. (7) 8: 174.
An incorrect subsequent spelling of Micragra Hampson, 1898.

MICRAGYRTA Butler, 1876, Cistula ent. 2: 114. CTEN Type-species: Agyrta gavisa Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 147, by original designation. [COLOMBIA]: Bogota.

A. gavisa is a junior subjective synonym of Dioptis diminuta Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 328.

MICRARCTIA Seitz, 1910, Gross-Schmett. Erde 2: 83. ARCT Type-species: Nyctemera trigona Leech, 1899, Trans. ent. Soc. Lond. 1899: 170, by subsequent designation by Hampson, 1920, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 2: 342. China: western.

The date of publication of that part of Seitz containing page 83 was cited on page 81 as 20.iv.1910 in both the German and English editions.

MICRAROA Hampson, 1905, Ann. S. Afr. Mus. 3: 404.

Type-species: *Micraroa rufescens* Hampson, 1905, *ibidem* 3: 405, by monotypy. South Africa: Natal.

MICRILEMA Hampson, 1903, Ann. Mag. nat. Hist. (7) 11: 343.

Type-species: *Micrilema craushayi* Hampson, 1903, *ibidem* (7) **11**: 343, by monotypy. [Lesotho] Basutoland: Machacha.

MICROBERGERIA Kiriakoff, 1972, Lambillionea 71: 102.

Type-species: *Microbergeria luctuosa* Kiriakoff, 1972, *ibidem* 71: 102-105, fig., by original designation. CAMEROUN: Lomié.

MICROCHADISRA Kiriakoff, 1960, Revue Zool. Bot. afr. 61: 58.

Type-species: *Microchadisra angustipennis* Kiriakoff, 1960, *ibidem* 61: 58, fig.10, by original designation. [ZAIRE]: Tshuapa, Bokuma.

‡MICRODATA; misspelling. See Trocodima Watson nom.

MICRODONTA Duponchel, [1845]. See Leucodonta Staudinger, 1892.

‡MICRODONTA; misspelling. See Mimodonta Matsumura, 1920.

MICRODONTELLA Strand, 1934. See Mimodonta Matsumura, 1920.

MICRODOTA Dognin, 1906. See Trocodima Watson nom.

MICROGITON Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.103, figs 17, 18.

Type-species: *Microgiton selene* Felder, 1874, *ibidem* 2 (Abt.2): pl.103, fig.17, by subsequent designation by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 467, but cited as *Virbia submacula* Walker, 1854, a nominal species not originally included in *Microgiton*. Hampson, however, placed *submacula* as the senior synonym of *selene*, a nominal species originally included in *Microgiton*. Under the *Code*, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. VENEZUELA: cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 6.

M. selene is a junior subjective synonym of Virbia submacula Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 472

See also Teracotona Butler, 1878.

MICROGYMNA Wallengren, 1865. See Lacipa Walker, 1855.

MICROHOPLITIS Marumo, 1920, J. Coll. Agric. imp. Univ. Tokyo 6: 281 (key), 296.

Type-species: *Drymonia circumscripta* Butler, 1885, *Cistula ent.* 3: 125, by original designation. JAPAN: Nikko.

See also Shachia Matsumura, 1919.

MICROHYLE Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 437.

Type-species: *Hylemera fadella* Mabille, 1882, *Naturaliste* 2: 100, by original designation. MADAGASCAR.

MICROLITHOSIA Daniel, 1954, Bonn. zool. Beitr. 5: 135.

Type-species: *Microlithosia shaowuica* Daniel, 1954, *ibidem* 5: 135, text-fig.97, by monotypy. CHINA: Fukien, Shaowu.

MICROMELALOPHA Nagano, 1916, Bull. Nawa ent. Lab. 1: 10.

Type-species: *Pygaera troglodyta* Graeser, 1890, *Berl. ent.* Z. **35**: 72, by monotypy. [U.S.S.R.]: Sidemi Island, south of Vladivostok.

‡MICROMELOLAPHA; Matsumura, 1929, Insecta matsum. 4: 45.

An incorrect subsequent spelling of *Micromelalopha* Nagano, 1916.

‡MICROMELIS; misspelling. See Micronaclia Hampson, 1898.

‡MICROMELOLAPHA; misspelling. See Micromelalopha Nagano, 1916.

‡MICROMETAPTERA; misspelling. See Metamicroptera Hulstaert, 1923.

MICROMMIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.32. LITH Type-species: Micrommia jugifera Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.32, by monotypy. [Brazil]: Amazon region, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

MICROMORPHE Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.98, fig.18.

Type-species: *Micromorphe choerotricha* Felder, 1874, *ibidem* 2 (Abt.2): pl.98, fig.18, by monotypy. [INDONESIA]: [Moluccas] ins. molucc., cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 6.

MICRONACLIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xi, 22, 128.

Type-species: *Pseudonaclia simplex* Butler, 1879, *Ann. Mag. nat. Hist.* (5) 4: 236, by original designation. MADAGASCAR: Fianarantsoa.

‡MICROMELIS; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling of *Micronaclia* Hampson, 1898, teste Neave, 1940, *Nomencl. 2001*. 3: 159.

MICRONYGMIA Toxopeus, 1948, Treubia 19: 476. LYMA Type-species: Artaxa linta Moore, [1860] 1858-9, in Horsfield Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 351, by original designation, but cited as 'Euproctis linta Moore'. Java.

MICROPHALERA Butler, 1885, Cistula ent. 3: 119. NOTO Type-species: Microphalera grisea Butler, 1885, ibidem 3: 120, by monotypy. JAPAN: Yezo.

MICROPTEROGYNA Rambur, 1866. See Orgyia Ochsenheimer, 1810.

MICRORGYIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.99, fig.12. LYMA Type-species: Microrgyia amazonum Felder, 1874, ibidem 2 (Abt.2): pl.99, fig.12, by monotypy. [BRAZIL]: [R. Amazon] fl. Amazonas, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

MICROSTOLA Lower, 1920, Trans. Proc. R. Soc. S. Aust. 44: 68.

Type-species: *Microstola ammoscia* Lower, 1920, *ibidem* 44: 68, by monotypy. Australia: North Queensland, near Cairns, Kuranda.

Microstola was established in the Xyloryctidae; it is here transferred to the Arctiidae, Lithosiinae on the advice of and with acknowledgements to Dr I.F.B. Common, C.S.I.R.O., Canberra, Australia.

MICROTANE Hampson, 1901, Ann. Mag. nat. Hist. (7) 8: 183.

Type-species: *Microtane fusca* Hampson, 1901, *ibidem* (7) 8: 183, by original designation. [SRI LANKA] Ceylon: Matelé.

MIDARA Schaus, 1928, Proc. ent. Soc. Wash. 30: 53. CTEN Type-species: Midara bengueta Schaus, 1928, ibidem 30: 54,

by original designation. Philippines: Benguet province, Baguio.

MIDASIA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 152 (key), 208.

Type-species: Niganda aurata Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 64, by original designation. INDIA: Darjeeling, Khasia Hills.

Midasia was established as a subgenus of Bireta Walker, 1856; it was placed as a junior subjective synonym of Eushachia Matsumura, 1925, by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 85.

See also Eushachia Matsumura, 1925.

MIEZA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 527. YPONOMEUTIDAE

Type-species: *Mieza igninix* Walker, 1854, *ibidem* 2: 527, by subsequent designation by Busck, 1913, *Insecutor Inscit. menstr.* 1: 71. U.S.A.: E. Florida, St John's Bluff.

Mieza has been used as the original generic name for species now placed in the Arctiidae.

MIGOPLASTIS Felder, 1868, Verh. zool.-bot. Ges. Wien 18 (Abh.): 285.

Type-species: Migoplastis ceylanica Felder, 1868, ibidem 18 (Abh.): 285, by monotypy. [SRI LANKA] Ceylon: Peradenia, near Kandy.

M. ceylanica is a junior subjective synonym of Hypsa correcta Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 217.

See also Pachyphilona Butler, 1875.

MILODORA Boisduval, 1870. See Phavaraea Walker, 1854.

MILTASURA Roepke, 1946, Tijdschr. Ent. 87: 85. LITH Type-species: Miltasura celebensis Roepke, 1946, ibidem 87: 85, text-fig.9, by monotypy. [Sulawesi] Celebes: Todjambu; and Para Salamakki, G. Lampobattang.

MILTOCHRISTA Hübner, [1819] 1816, Verz. bekannter Schmett.: 166.

Type-species: *Noctua rubicunda* [Denis & Schiffermüller], 1775, *Ankündung syst. Werkes Schmett. Wienergegend*: 68, by monotypy. [AUSTRIA]: Vienna district.

N. rubicunda is a junior subjective synonym of Phalaena miniata Forster, 1771, Nova Species Insectorum: 75.

See also Calligenia Duponchel, [1845].

‡MILTOCHRYSTA; Humphreys & Westwood, 1843, Br. Moths 1: 98.

An incorrect subsequent spelling of *Miltochrista* Hübner, [1819].

‡MITTOCHISTA; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1685.

An incorrect subsequent spelling of *Miltochrista* Hübner, [1819].

‡MILTOCHRYSTA; misspelling. See Miltochrista Hübner, [1819].

MIMAGYRTA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xiii, 25, 206.

Type-species: *Metastatia pampa* Druce, 1893, *Proc. zool. Soc. Lond.* **1893**: 284, by original designation. ECUADOR: Sarayacu.

MIMARCTIA Neumoegen & Dyar, 1894, Jl N.Y. ent. Soc. 2: 9.

Type-species: *Phalaena arge* Drury, 1773, *Illust. nat. Hist.* exot. *Insects* 2: index to vol.1; 1770, *ibidem* 1: 35, pl.18, fig.3

(but without an available name until 1773), by monotypy. [U.S.A.]: New York.

MIMESISOMERA Bryk, 1950, Ark. Zool. 42 (A) 19: 27.

Type-species: Mimesisomera aureobrunnea Bryk, 1950, ibidem 42 (A) 19: 28, pl.2, fig.11, by original designation. BURMA: Kambaiti.

MIMICA Oberthür, 1881, Etudes d'Entomologie 6: 33. CTEN Type-species: Mimica lycoides Oberthür, 1881, ibidem 6:

33, pl.10, fig.9, by monotypy. PERU: Punamarca.

M. lycoides Oberthür, 1881, and Euchromia lycoides Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 256, were both placed in Correbia Herrich-Schäffer, [1855], by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 514. M. lycoides Oberthür thus became a junior secondary homonym for which the objective replacement name Correbia oberthuri [sic] Hampson, 1898, ibidem 1: 515, was established.

MIMODONTA Matsumura, 1920, Zool. Mag. Tokyo 32: 144.

Type-species: Mimodonta albicosta Matsumura, 1920, ibidem 32: 144, by monotypy. JAPAN: Honshu, Nikko, Chichibu.

Mimodonta was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 794.

‡MICRODONTA; Strand, 1934, in Gaede, in Strand, Lepid. Cat. 59: 111.

An incorrect subsequent spelling of Mimodonta Matsumura,

MICRODONTELLA Strand, 1934, in Gaede, in Strand, Lepid. Cat. 59: 111.

An unnecessary replacement name for Mimodonta Matsumura, 1920, which Strand, as editor, added to Gaede's work but spelt incorrectly as ‡Microdonta, a preoccupied name.

MIMOPYDNA Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 37.

Type-species: Bireta pallida Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 473, by original designation (but cited as 'pallida Walker', an incorrect authorship). JAPAN: Yokohama.

MIMULOSIA Toulgoët, 1958, Mém. Inst. scient. Madagascar (E) 9: 200.

Type-species: Philenora pseudotortrix Toulgoët, 1955, Revue fr. Ent. 22: 196, text-fig.5, pl.6, fig.16, by original designation. MADAGASCAR: central, massif de l'Ankaratra, Manjakatompo, Ambahona forest.

MINARA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1711.

Type-species: Minara pardalina Walker, 1856, ibidem 7:

1712, by monotypy. BRAZIL.

M. pardalina is a junior subjective synonym of Notodonta histrionica Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1(1): wrapper, pl.67, fig.382.

MINTOPOLA Schaus, 1899, Jl N.Y. ent. Soc. 7: 217. LITH Type-species: Mintopola braziliensis Schaus, 1899, ibidem 7: 217, by monotypy. Brazil: Nova Friburgo.

Mintopola probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson, 1900, as a name for a new genus.

MINTOPOLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 87, 189.

Type-species: Mintopola braziliensis Schaus, 1899, by original designation.

A junior homonym of Mintopola Schaus, 1899. The objective replacement name is Mintopola Schaus, 1899.

MINTOPOLA Hampson, 1900. See Mintopola Schaus, 1899.

MIOSTAUROPUS Kiriakoff, 1964, Bonn. zool. Beitr. 14: 273.

Type-species: Stauropus mioides Hampson, 1904, J. Bombay nat. Hist. Soc. 16: 150, by original designation. INDIA: Assam, Khasis.

MIRACIDION Kiriakoff, 1953, Lambillionea 53: 93. Type-species: Amata caudatula Kiriakoff, 1953, ibidem 53: 93, pl.1, fig.6, by original designation. [ZAIRE] Congo Belge: Lualaba, Kapanga.

Miracidion was established as a subgenus of Amata Fabricius, 1807.

MIRANDISCA Travassos, 1955, Archos Mus. nac. Rio de J. 42: 669.

Type-species: Cosmosoma harpalyce Schaus, 1892 August, Proc. zool. Soc. Lond. 1892: 275, by original designation. Brazil: Petropolis.

C. harpalyce was redescribed by Schaus, 1892 November, Am. Lepid.: 4.

MIROBRIGA Walker, 1864. See Birgorima Watson nom. n.

MISMIA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 151 (key), 159.

Type-species: Mismia impunctibasis Kiriakoff, 1962, ibidem 98: 159, fig.5, pl.1, col.2, by original designation. [BURMA]: Mishmi Hills, Pange.

MISOGADA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 449. NOTO

Type-species: Misogada sobria Walker, 1865, ibidem 32: 450, by monotypy. North America.

M. sobria is a junior subjective synonym of Lochmaeus unicolor Packard, 1864, Proc. ent. Soc. Philad. 3: 373.

See also Stematocampa Packard, 1895.

MITHUNA Moore, 1878, Proc. zool. Soc. Lond. 1878: 21.

Type-species: Mithuna quadriplaga Moore, 1878, ibidem 1878: 21, pl.2, fig.9, by monotypy. [INDIA]: Darjeeling.

MITRADAEMON Butler, 1878, Trans. ent. Soc. Lond. 1878: 60.

Type-species: Mitradaemon velutinum Butler, 1878, ibidem 1878: 60, by original designation. [Brazil]: Tabatinga, Barreiras de Jutahi.

ACTEA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 307.

Type-species: Hipocrita monilis Hübner, [1811] 1806, Samml. exot. Schmett. 1: pl.[183], by monotypy. Type-local-ITY: not stated, [ECUADOR], [BRAZIL].

A junior homonym of Actea Germar, 1842, in Münster, Beitr. zur Petrefacten-Kunde 5: 85, - Insecta, Hemiptera. There is no objective replacement name, but H. monilis is currently treated as being congeneric with Mitradaemon velutinum Butler, 1878, the type-species of Mitradaemon Butler, 1878; the latter is thus available for use as a subjective replacement name.

‡ACTAEA; Druce, 1885, Proc. zool. Soc. Lond. 1885: 526. An incorrect subsequent spelling of Actea Walker, 1854.

‡MITTOCHISTA; misspelling. See Miltochrista Hübner, [1819].

MOCHLOPTERA Butler, 1876, J. Linn. Soc. (Zool.) 12: 386.

Type-species: Glaucopis acroxantha Perty, [1833] 1830-

1834, in Martius, Delectus Anim. articulatorum Bras.: 156, pl.31, fig.4, by original designation. BRAZIL: equatorial.

G. acroxantha is a junior subjective synonym of Isanthrene chalciope Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 20, pl.[81], figs 469, 470.

See also Nyridela Lucas, 1857.

MOMONIPTA Warren, 1897, Novit. zool. 4: 421. Type-species: Momonipta albiplaga Warren, 1897, ibidem 4: 421, by original designation. COLOMBIA: R. Dagua.

Momonipta was established in the Geometridae; placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 412.

MONARCTIA Talbot, 1929, Bull. Hill Mus. Witley 3: 126.

Type-species: Monarctia lactea Talbot, 1929, ibidem 3: 126, pl.3, fig.9, by original designation. Angola: S. central, Upper Cubango-Cunene watershed.

MONOCREAGRA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.8.

Type-species: Monocreagra pheloides Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.8, by monotypy. [COLOMBIA]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

See also Tanaostyla Warren, 1897.

MONOSYNTAXIS Swinhoe, 1901, Ann. Mag. nat. Hist. (7) 7: 466.

Type-species: Monotaxis trimaculata Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 182, by original designation (for Monotaxis Hampson, 1900). Borneo: [Sabah], Kina Balu.

Monosyntaxis was established as the objective replacement name for Monotaxis Hampson, 1900.

MONOTAXIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 83, 181.

Type-species: Monotaxis trimaculata Hampson, 1900, by original designation.

A junior homonym of Monotaxis [Bennett], 1830, in Raffles, Mem. T.S. Raffles: 688, - Pisces. The objective replacement name is Monosyntaxis Swinhoe, 1901.

MONOTAXIS Hampson, 1900. See Monosyntaxis Swinhoe, 1901.

MONOTROPIA Kirby, 1892. See Licnoptera Meyrick, 1889. LITH

MOOREA Hampson, 1894. See Sebastia Kirby, 1892. ARCT

MORASA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 859. LYMA

Type-species: Morasa modesta Walker, 1855, ibidem 4: 859, by monotypy. South Africa.

Morasa when established contained a second nominal species, but this was doubtfully included and, under the Code Article 68(c), is not eligible for fixation as type-species.

See also Sarothropyga Felder, 1874.

MORESA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1159 (key), 1170.

Type-species: Moresa costalis Walker, 1855, ibidem 5: 1171, by monotypy. BRAZIL.

Moresa was established in the Drepanidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 580.

‡MOSADA; misspelling. See Mosoda Walker, 1866. LITH

MOSARA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1032. NOCTUIDAE

Type-species: Mosara apicalis Walker, 1855, ibidem 5: 1032, by monotypy. INDIA: northern.

Mosara was established in the Notodontidae: it was placed in the Noctuidae by Moore, [1885] 1884-7, Lepid. Ceylon 3: 118.

MOSODA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1899.

Type-species: Mosoda anartoides Walker, 1866, ibidem 35: 1900, by monotypy. AUSTRALIA.

M. anartoides is a junior subjective synonym of Setina sinuata Wallengren, 1860, Wien. ent. Monatschr. 4: 46.

‡MOSADA; Turner, 1940, Proc. R. Soc. Qd 51: 82. An incorrect subsequent spelling of Mosoda Walker, 1866.

MOTADA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 924.

Type-species: Motada lateralis Walker, 1855, ibidem 4: 924, by monotypy. [BRAZIL]: and Rio [de] Janeiro.

MPANJAKA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80:

Type-species: Mardara viola Butler, 1879, Ann. Mag. nat. Hist. (5) 4: 240, by original designation. MADAGASCAR: Antananarivo.

MULONA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1896.

Type-species: Mulona lapidaria Walker, 1866, ibidem 35: 1896, by monotypy. [HISPANIOLA] St Domingo.

MUNONA Schaus, 1894, Proc. zool. Soc. Lond. 1894: 233.

Type-species: Munona iridescens Schaus, 1894, ibidem 1894: 233, by monotypy. VENEZUELA: Aroa.

MUNYCHIA Wallengren, 1858, Ofvers. K. VetenskAkad. Förh. Stockh. 15: 213.

Type-species: Munychia callipyga Wallengren, 1858, ibidem 15: 214, by original designation. [South Africa] Caffraria.

M. callipyga is a junior subjective synonym of Phalaena amasis Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 23, pl.206, fig.B.

See also Rhodogastria Hübner, [1819].

MUXTA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 58.

Type-species: Lepista xanthopa Holland, 1893, Psyche, Camb. 6: 411, by original designation. [GABON]: [Ogooué] Ogové R. valley.

MYCLELA Watson nom. n. for Cymella Felder, 1874. LITH Type-species: Cymella congerens Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.14, by monotypy (of Cymella Felder, 1874). Type-LOCALITY: not known. There are no specimens in the BMNH which match the type but closely related nominal species occur in INDIA.

CYMELLA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.14.

Type-species: Cymella congerens Felder, 1874, by monotypy. A junior homonym of Cymella Meek, 1864, Smithson. misc. Collns 7 (177): 14, 34, - Mollusca. The objective replacement name is Myclela Watson nom. n.

MYDROMERA Butler, 1876, Cistula ent. 2: 112. CTEN Type-species: Euagra isthmia Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.24, by monotypy. [Brazil]: Amazon R., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 4.

MYDROPASTEA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xvi, 25, 326.

Type-species: *Eupyra albicincta* Schaus, 1896, *Jl N.Y. ent. Soc.* 4: 131, by original designation. ECUADOR: Chimbo.

MYLANTRIA Aurivillius, 1904, Ark. Zool. 2 (4): 54. LYMA Type-species: Aroa xanthospila Plötz, 1880, Stettin. ent. Ztg 41: 84, by monotypy. [Ghana]: Aburi.

MYONIA Walker, 1854. See Erbessa Walker, 1854. DIOP

MYOPSYCHE Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: [ix], 22, 48.

Type-species: *Syntomis ochsenheimeri* Boisduval, [1828] 1829, *Essai Monogr. Zygénides*: 116, pl.7, fig.4, by original designation. [INDIA]: Bengal, JAVA.

MYRMECOPSIS Newman, 1850, Zoologist 8 (Appendix): exxii.

Type-species: Myrmecopsis eumenides Newman, 1850, ibidem 8 (Appendix): cxxii, by monotypy. [BRAZIL]: [R. Amazon], [Tefé] Ega.

M. eumenides is a junior subjective synonym of Pseudosphex polistes Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 12, pl. [7], figs 39, 40.

MYSTROCNEME Herrich-Schäffer, 1855, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.73, fig.419.

Type-species: *Mystrocneme geminata* Herrich-Schäffer, 1855, *ibidem* 1(1): wrapper, pl.73, fig.419, by monotypy. [BRAZIL]: Cameta.

M. geminata is a junior subjective synonym of Euchromia varipes Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 235.

See also Leucotmemis Butler, 1876.

NACLIA Boisduval, 1840. See Dysauxes Hübner, [1819].

NACLIODES Strand, 1912, Arch. Naturgesch. 78 (A) 6: 183.

Type-species: Meganaclia microsippia Strand, 1912, ibidem 78 (A) 6: 183, by monotypy. [CAMEROUN or EQUATORIAL GUINEA] Süd-Kamerun or Spanisch Guinea: 'Alen'; 'Makonam nach Mabungo'.

Nacliodes was established as a subgenus of Meganaclia Aurivillius, 1892.

NADATA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1062.

Type-species: *Phalaena gibbosa* Smith, 1797, *in* Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 163, pl.82, by monotypy. [U.S.A.]: Georgia.

ALASTOR Boisduval, 1869, Annls Soc. ent. Belg. 12: 87.

Type-species: *Phalaena gibbosa* Smith, 1797, by monotypy. A junior homonym of *Alastor* Lepeletier, 1841, *Hist. nat. Insectes* (Hyménoptères) 2: 668, – Insecta, Hymenoptera. The objective replacement name is *Nadata* Walker, 1855.

NADIASA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 1014.

LASIOCAMPIDAE

Type-species: Nadiasa concolor Walker, 1855, ibidem 5: 1015, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 842. SOUTH AFRICA.

Nadiasa was established in the Notodontidae; it was placed in the Lasiocampidae by Kirby, 1892, ibidem 1: 842.

NADUNA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 260 (key), 291.

NOTO
Type-species: Naduna lignea Schaus, 1901, ibidem 1901:

291, pl.11, fig.8, by original designation. BRAZIL: south-east-ern, São Paulo.

NAENIA Stephens, 1827, in Anonymous, Retrospective Review (2) 1: 243.

Type-species: *Phalaena typica* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 518, by monotypy (but included as *typicoides* Donovan, a misidentification; see below). Type-locality: not stated, [Europe].

P. typicoides Donovan sensu Stephens, 1827, was a misidentification of Phalaena typicoides Donovan, 1811, Nat. Hist. Br. Insects 15: 1, pl.505. In Stephens, 1829, Illust. Br. Ent. (Haustellata) 2: 165 the error was corrected.

Under the *Code*, Article 70(a), the case of a misidentified type-species should be referred to the Commission.

PHALAENA Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 495.

Type-species: *Phalaena typica* Linnaeus, 1758, by indication, under the *Code*, Articles 67(b) and 68(b).

Phalaena was suppressed for the purposes of the Law of Priority, but not for those of the Law of Homonymy, by the International Commission on Zoological Nomenclature, 1957, Opin. Decl. int. Commn zool. Nom. 15 (Opinion 450): 253. Placed on the Official Index of Rejected and Invalid Generic Names in Zoology with Name No. 850.

Phalaena was used by Linnaeus to include all Heterocera except the Sphingidae, and has been used as the original generic name for species placed in the families included in this catalogue.

NAGANDOPSIS Matsumura, 1934, Insecta matsum. 8: 152, 153.

Type-species: *Nagandopsis kawachiensis* Matsumura, 1934, *ibidem* 8: 152, fig.1, by original designation. JAPAN: Honshu, Mitake district, Kawachi near Tokyo.

N. kawachiensis is a junior subjective synonym of Asteroscopus atrovittatus Bremer, 1861, Bull. Acad. imp. Sci. St. Pétersb. 3: 483.

See also Shaka Matsumura, 1920.

NAGANOEA Matsumura, 1920, Zool. Mag. Tokyo 32: 142.

Type-species: *Drymonia manleyi* Leech, 1889, *Proc. zool.* Soc. Lond. **1888**: 639, pl.32, fig.2, by monotypy. JAPAN: Yokohama.

Naganoea was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 775.

NAGIDUSA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 257.

Type-species: Nagidusa xylocampoides Walker, 1862, ibidem (3) 1: 257, by monotypy. [Brazil]: Rio de Janeiro.

‡DASUNAGA; Holland, 1893, Psyche, Camb. 6: 551.

An incorrect subsequent spelling of Nagidusa Walker, 1862. In the original description of Dinotodonta Holland, 1893, ibidem 6: 551, Holland stated 'Allied to Dasunaga, Moore'. Neave, 1939, Nomencl. zool. 2: 16, cited 'Dasunaga (err. pro Nagidusa Walker, 1862)'. No other published reference to the name ‡Dasunaga has been found.

NAGUNDA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 53.

Type-species: Alope semicincta Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 620, by original designation. INDIA: North.

NANNA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) Zool. 1: 35, figs.

Type-species: *Eilema diplisticta* Bethune-Baker, 1911. *Ann. Mag. nat. Hist.* (8) 7: 535, by original designation. WEST AFRICA: Gunnal.

NANNOCERYX Roepke, 1937, Ent. Z., Frankf.a.M. 50: 490.

Type-species: *Eressa myiella* van Eecke, 1920, *Zool. Meded. Leiden* 5: 116, fig., by monotypy. Java: Wonosobo.

NANNODOTA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 407.

Type-species: *Halysidota minuta* Rothschild, 1910, *Novit. zool.* 17: 67 (but cited as ‡*Halisidota*, an incorrect subsequent spelling), by original designation (but cited as ‡*Halisidota*). [PERU]: Carabaya, La Oroya.

NAPATA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 231.

Type-species: *Euchromia terminalis* Walker, 1854, *ibidem* 1: 231, by monotypy. Brazil: unstated locality; and Pernambuco, HONDURAS.

Napata was established as a subgenus of Euchromia Hübner, [1819].

‡NEPETA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling of Napata Walker, 1854, teste Neave, 1940. Nomencl. zool. 4: 314.

NAPREPA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1046.

Type-species: *Naprepa camelinerdes* Walker, 1855, *ibidem* 5: 1046, by monotypy. BRAZIL.

‡NARASODES; misspelling. See Narosodes Moore, [1887].

NAROMA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1744.

Type-species: *Naroma signifera* Walker, 1856, *ibidem* 7: 1744, by monotypy. SIERRA LEONE.

See also Zarfa Walker, 1869.

NAROSODES Moore, [1887] 1884-7, Lepid. Ceylon 3: 535.

Type-species: Tospitis punctana Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 431, by monotypy. [SRI LANKA] Ceylon.

‡NARASODES; Rothschild, 1913, Novit. zool. 20: 224.

An incorrect subsequent spelling of Narosodes Moore, [1887], repeated by Hampson, 1926, Descr. new Genera Species Lepid. Phalaenae Subfamily Noctuinae Br. Mus.: 192.

NAVARCOSTES Schaus, 1905, Proc. U.S. natn. Mus. 29: 280.

Type-species: *Navarcostes limnatis* Schaus, 1905, *ibidem* **29**: 280, by monotypy. French Guiana: St Jean.

NAYACA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 43.

Type-species: Arctia imbuta Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 614, by original designation. INDIA: northern; and Himalayas.

NEACEREA Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1:

Type-species: Neacerea albiventus Druce, 1898 May, ibidem (7) 1: 406, by subsequent designation by Collins, 1962, Ann. Mag. nat. Hist. (13) 5: 123. BRAZIL: Minas [Gerais] Geraes.

Neacerea probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson, 1898 [December], as the name for a new genus.

NEACEREA Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xx, 27, 478.

Type-species: Aclytia rufiventris Schaus, 1894, Proc. zool. Soc. Lond. 1894: 228, by original designation. MEXICO: Jalapa.

A junior homonym of Neacerea Druce, 1898 May. There is no objective replacement name but the type-species of Neacerea Hampson, 1898 [December], is currently considered to be congeneric with the type-species of Neacerea Druce, 1898 May. The latter is therefore available for use as a subjective replacement name.

NEACEREA Hampson, 1898 [December]. See Neacerea Druce, 1898 May.

NEAGYLLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 91, 213.

Type-species: Areva nevosa Dognin, 1894, Annls Soc. ent. Belg. 38: 128, by original designation. [ECUADOR]: San Francisco, near Loja.

NEARDONAEA Gaede, 1925, Mitt. zool. Mus. Berl. 11: 240.

Type-species: Neardonaea metallica Gaede, 1925, ibidem 11: 240, by original designation. VENEZUELA: Mérida.

NEASURA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvii, 83, 422.

Type-species: *Neasura hypophaeola* Hampson, 1900, *ibidem* 2: 423, fig.335, by original designation. [Indonesia]: [Sangihe] Sangir.

NEASUROIDES Matsumura, 1927, J. Coll. Agric. Hokkaido imp. Univ. 19 (1): 72.

Type-species: *Neasuroides simplicior* Matsumura, 1927, *ibidem* **19** (1): 72, by original designation. [TAIWAN] Formosa: Horisha.

NEAXIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 2, 30. ARCT

Type-species: Amaxia theon Druce, 1900, Ann. Mag. nat. Hist. (7) 6: 66, by original designation. Venezuela: Mérida.

NEBRARCTIA Watson nom. n. for Spilarctia Staudinger, 1891.

Type-species: Arctia semiramis Staudinger, 1891, Dt. ent. Z. Iris 4: 161, by monotypy (of Spilarctia Staudinger, 1891). [Turkey] Kleinasien.

SPILARCTIA Staudinger, 1891, Dt. ent. Z. Iris 4: 161, 162. Type-species: Arctia semiramis Staudinger, 1891, by monotypy.

A junior homonym of *Spilarctia* Butler, 1875, *Cistula ent*. 2: 39, – Lepid., Arctiidae. The objective replacement name is *Nebrarctia* Watson nom. n.

Spilarctia was established as a subgenus of Arctia Schrank, 1802.

NEERESSA Hampson, 1905, Ann. Mág. nat. Hist. (7) 15: 426.

Type-species: *Trichaeta sagada* Semper, 1898, *Reisen Archipel Philipp. Semper* (2) **6**: 423, by original designation. Philippines: north Luzon, Cagayan.

NEEUGOA Wileman & South, 1916, Entomologist 49: 178.

Type-species: *Neeugoa kanshireiensis* Wileman & South, 1916, *ibidem* 49: 179, by original designation. [TAIWAN] Formosa: Kanshirei.

NEIDALIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 3, 13. ARCT

Type-species: *Idalus villacresi* Dognin, 1894, *Annls Soc. ent. Belg.* **38**: 238, by original designation. [ECUADOR]: Zamora, near Loja.

NELOPA Billberg, 1820. See Nudaria Haworth, [1809].

NELPHE Herrich-Schäffer, [1858] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 74 (but attributed to Boisduval, an incorrect authorship).

Type-species: Charidea confinis Herrich-Schäffer, [1855] 1850-1858, ibidem 1 (1): pl.51, fig.277 (but attributed to Boisduval, an incorrect authorship), by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 172.

Nelphe was first published in the synonymy of Charidea Dalman, 1816, but was subsequently treated by Kirby, 1892, ibidem 1: 172, as a valid name with Herrich-Schäffer as author and is therefore nomenclaturally available under the Code, Article 11(d).

NEMEOPHILA Stephens, 1828. See Parasemia Hübner, [1820].

NEMETA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 968.

Type-species: *Nemeta bifacies* Walker, 1855, *ibidem* 4: 968, by monotypy. BRAZIL.

NEMETA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1083.

Type-species: Nemeta basifusca Walker, 1855, ibidem 5:

1083, by monotypy. [BRAZIL]: Rio de Janeiro.

A junior homonym of *Nemeta* Walker, 1855, *ibidem* 4: 968, – Lepid., Limacodidae. There is no objective replacement name, but *Nemeta basifusca* is conspecific with *Nemeta bifacies* Walker, 1855, the type-species of *Nemeta* Walker, 1855, *ibidem* 4: 968; the latter is thus available for use as a subjective replacement name.

Nemeta Walker, 1855, ibidem 5: 1083 was established in the Notodontidae; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1 placed it in the Limacodidae on page 525 and in the Notodontidae on page 620. It is currently included in the Limacodidae in the synonymy of Phobetron Hübner, [1825].

NEMETA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1083. See Nemeta Walker, 1855, ibidem 4: 968.

LIMACODIDAE

NEOARCTIA Neumoegen & Dyar, 1893, Ent. News 4: 141.

Type-species: Antarctia beanii Neumoegen, 1891, Can. Ent. 23: 123, by original designation. Canada: Alberta, Laggan.

NEOBALACRA Kiriakoff, 1952, Biol. Jaarb. 19: 79. THYRET Type-species: Balacra paradoxa Hering, 1932, Revue Zool. Bot. afr. 22: 107, pl.2, fig.30, by original designation. [ZAIRE]: [Katanga], Lubumbashi.

B. paradoxa is a junior subjective synonym of Metamicroptera rotundata Hulstaert, 1923, the type-species of Metamicroptera Hulstaert, 1923.

See also Metamicroptera Hulstaert, 1923.

NEOBLACODES Fletcher nom. n. for Blacodes Kiriakoff, 1959.

Type-species: Blacodes friedae Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 182, fig.27, by original designation (for Blacodes Kiriakoff, 1959). ZAIRE: Sankuru, Lusambo.

The objective replacement name was suggested by Prof. Dr S.G. Kiriakoff in a personal communication

BLACODES Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 182.

Type-species: *Blacodes friedae* Kiriakoff, 1959, by original designation.

A junior homonym of Blacodes Dejean, 1834, Cat. Coléop.

(Edn 2): 190, – Insecta, Coleoptera. The objective replacement name is *Neoblacodes* Fletcher nom. n.

NEOBLAVIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 81, 105.

Type-species: *Neoblavia scoteola* Hampson, 1900, *ibidem* 2: 105, fig.45, by original designation. [INDIA]: Sikkim.

‡NEOBOURQUINIA Köhler, 1943. See Neobourquinia Köhler gen. n.

NEOBOURQUINIA Köhler gen. n. NOTO Type-species: Neobourquinia bifasciata Köhler, 1943, Revta Soc. ent. argent. 12: 27, pl.6, figs 3, 4, by PRESENT DESIGNATION. [ARGENTINA]: Cordoba, Sierra de la Ventana.

Neobourquinia Köhler is here established with the agreement of Dr P. Köhler (personal communication to D.S. Fletcher of 25.i.1978) for the concept of ‡Neobourquinia Köhler, 1943, ibidem 12: 27. When the name was originally proposed in 1943 for two species, it was accompanied by a statement 'that purports to give characters differentiating the taxon' (Article 13(a)(i)) but was not 'accompanied by the definite fixation of a type-species' (Article 13(b)).

The term 'gen.n., sp.n. or its exact equivalent' published as 'Neobourquinia bifasciata, gen. et spec. nov.' in the legend to plate 6, figure 3, accompanying the original description of the type-species in 1943, cannot be interpreted as the original designation of a type-species of a new nominal genus after 1930 (Article 68(a)(i)).

Neobourquinia is unavailable from 1943, being a generic name published after 1930 that does not satisfy the provisions of the Code, Article 13 (b). The type-species Neobourquinia bifasciata is, however, available from 1943; under the Code, Article 17(3), a species-group name remains available even though the genus-group name with which it was first combined is invalid or unavailable.

‡NEOBOURQUINIA Köhler, 1943, Revta Soc. ent. argent. 12: 27.

‡Neobourquinia was proposed for two included nominal species, but was not originally accompanied by the definite fixation of a type-species and is therefore unavailable under the Code, Article 13(b).

NEOBROCHA Meyrick, 1886, *Proc. Linn. Soc. N.S.W.* (2) 1: 692, 707.

Type-species: *Neobrocha phaeocyma* Meyrick, 1886, *ibidem* (2) 1: 708, by subsequent designation by Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* 2: 498. [Australia]: [Queensland], Thursday Island.

NEOCERURA Matsumura, 1929, Insecta matsum. 4: 88 (key), 89.

Type-species: Cerura liturata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 988, by original designation. [BANGLADESH]: Sylhet, [INDIA]: Madras.

NEOCHADISRA Kiriakoff, 1969, Faune Madagascar 29: 53.

Type-species: *Neochadisra protea* Kiriakoff, 1969, *ibidem* 29: 54, text-figs 32, 33, pl.2, figs 16, 18, by original designation. MADAGASCAR: eastern, Perinet district, Analamazaotra forest.

NEOCHELONIA Draeseke, 1926. See Euleechia Dyar, 1900.

NEOCHERA Hübner, [1819] 1816, Verz. bekannter Schmett.: 173.

Type-species: *Phalaena dominia* Cramer, 1780, *Uitlandsche Kapellen (Papillons exot.)* 3: 123, 174, pl.263, figs A, B, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 389. [INDIA]: Coromandel.

P. dominia is dated from the wrapper of the part and not from the title-page of the volume, 1782.

NEOCLOSTERA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 254.

Type-species: *Neoclostera insignior* Kiriakoff, 1964, *ibidem* 14: 254, abb.1, fig.8, pl.1, fig.6, by original designation. CHINA: north Yunnan, A-tun-tse.

NEODRYMONIA Matsumura, 1920, Zool. Mag. Tokyo 32: 143.

Type-species: Drymonia delia Leech, 1889, Proc. zool. Soc. Lond. 1888: 640, pl.32, fig.3, by monotypy. Japan: Oiwake. Neodrymonia was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 780.

NEODUMA Hampson, 1918, Novit. zool. 25: 101. LITH Type-species: Neoduma ectozona Hampson, 1918, ibidem 25: 102, by original designation. Philippines: Luzon, Los Baños.

NEOFENTONIA Matsumura, 1929, Insecta matsum. 4: 40.

Type-species: *Neofentonia acuminata* Matsumura, 1929, *ibidem* 4: 40, pl.1, fig.8, by original designation. Taiwan: Horisha.

NEOHARPYIA Daniel, 1965, Z. wien. ent. Ges. 50: 44. NOTO Type-species: Cossus verbasci Fabricius, 1798, Ent. Syst. (Suppl.): 441, by original designation. [FRANCE]: Montpellier.

NEOLA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1033.

Type-species: *Neola semiaurata* Walker, 1855, *ibidem* 5: 1033, by monotypy. [AUSTRALIA]: Tasmania, New South Wales, Sydney.

NEOLAURONA Druce, 1893, Proc. zool. Soc. Lond. 1893: 292.

Type-species: *Pericopis hazara* Butler, 1871, *Ann. Mag. nat. Hist.* (4) 8: 287, by original designation. [BRAZIL]: [Para], Vila Nova, ECUADOR.

NEOLIPARINA Schultze, 1934, Dt. ent. Z. Iris 48: 107.

Type-species: *Neoliparina paradoxa* Schultze, 1934, *ibidem* **48**: 107, pl.1, fig.2, by monotypy. [Central African Empire]: Ubangi, Mongoumba.

NEOMARDARA Hering, 1926, in Seitz, Gross-Schmett. Erde

Type-species: *Lepasta africana* Holland, 1893, *Ent. News* 4: 343, pl.15, fig.11, by original designation. [GABON]: [Ogooué R. valley], Kangwé.

The date of publication of the part of Seitz containing page 155 was stated on page 153 as 5.xi.1926 in both the German and the English editions.

‡NEOMELESE Zikan, 1968, in Zikan & Zikan, Pesquisa Agropec. bras. 3: 91.

Neomelese is undescribed and unavailable under the Code, Article 13(a). The only included nominal species, ‡Neomelese mimica Zikan, 1968, ibidem 3: 91 is also undescribed and unavailable under Article 13.

NEOMULONA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxv, 700.

Type-species: Clemensia torniplaga Jones, 1914, Trans. ent. Soc. Lond. 1914: 9, pl.1, fig.24, by original designation. BRAZIL: São Paulo.

NEONERITA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 2, 37. ARCT

Type-species: *Neonerita dorsipuncta* Hampson, 1901, *ibidem* 3: 37, fig.30, by original designation. BRAZIL: Rio Grande do Sul.

NEOPHEMULA Kiriakoff, 1957, Revue Zool. Bot. afr. 55: 282.

Type-species: *Pseudapiconoma vitrina* Oberthür, 1909, *Etud. Lépid. comp.* 3: 95, 99, pl.10, fig.5, by original designation. [Cameroun] Kamerun.

‡NEOPHEOGIA; misspelling. See Neopheosia Matsumura, 1920.

NEOPHEOSIA Matsumura, 1920, Zool. Mag. Tokyo 32: 147.
NOTO

Type-species: *Pheosia fasciata* Moore, 1888, *Proc. zool. Soc. Lond.* **1888**: 401, by monotypy. [INDIA]: Dharmsala.

In the original combination of the type-species, the generic name was published as ‡*Pheosa*, an incorrect subsequent spelling.

Neopheosia was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 783, 784; on page 784 the name was incorrectly spelt as ‡Neopheogia.

‡NEOPHEOGIA; Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 784.

An incorrect subsequent spelling, repeated by Schaus, 1928, *Proc. U.S. natn. Mus.* 73 (19): 81; the Schaus misspelling was listed by Neave, 1940, *Nomencl. zool.* 4: 748.

NEOPHYTA Bryk, 1950, Ark. Zool. 42 (A) 19: 36. NOTO Type-species: Celeia sikkima Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 63, by original designation. INDIA: Darjeeling.

NEOPLYNES Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 90, 250.

Type-species: *Lithosia cytheraea* Druce, 1894, *Ann. Mag. nat. Hist.* (6) 13: 177, by original designation. MEXICO: near Durango city.

NEORGYIA Bethune-Baker, 1908, Novit. zool. 15: 188.

Type-species: Neorgyia ochracea Bethune-Baker, 1908, ibidem 15: 189, by original designation. New Guinea: [Papua New Guinea], [Papua], Mambaré River, Biagi.

NEOSCAPTIA Hampson, 1900, Cat. Lepid.Phalaenae Br. Mus. 2: xiii, 84, 282.

Type-species: *Neoscaptia collateralis* Hampson, 1900, *ibidem* **2**: 282, by original designation. New Guinea: [Irian Jaya], [Kapaor] Kapaur.

NEOSELCA Hampson, 1898, Trans. ent. Soc. Lond. 1898: 244.

Type-species: *Nola minuta* Druce, 1885, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 1: 140, pl.13, fig.17, by original designation. Costa Rica: Irazu.

NEOSHACHIA Matsumura, 1925, Zool. Mag. Tokyo 37: 400.

Type-species: *Neoshachia parabolica* Matsumura, 1925, *ibidem* 37: 400, pl.6, fig.6, by original designation. Talwan.

N. parabolica is a junior subjective synonym of Harpyia ocypete Bremer, 1861, Bull. Acad. imp. Sci. St.-Pétersb. 3: 481.

See also Fentonia Butler, 1881, and Urocampa Staudinger, 1892.

NEOSICCIA van Eecke, 1926, Zool. Meded. Leiden 9: 260 (key). Nomenclaturally available but without included species until van Eecke, 1927, ibidem 10: 95.

Type-species: *Neosiccia accurata* van Eecke, 1927, *ibidem* **10**: 95; 1926, *ibidem* **9**: pl.4, fig.8 (without legend), by original designation. Sumatra: Fort de Kock.

NEOSTAUROPUS Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 89.

Type-species: Stauropus basalis Moore, 1877, Ann. Mag. nat. Hist. (4) 20: 90, by original designation. [CHINA]: Shanghai.

NEOTALARA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxv, 693.

Type-species: *Neotalara metamelaena* Hampson, 1914, *ibidem* 1: 693, fig.221, by monotypy. ECUADOR: Quevedo.

NEOTHYONE Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxiv, 647.

Type-species: *Talara schistaceoplagiata* Rothschild, 1913, *Novit. zool.* 20: 201, by original designation. PERU: Carabaya, Oconeque.

NEOTRICHURA Druce, 1896, Ann. Mag. nat. Hist. (6) 18: 33.

Type-species: Neotrichura penates Druce, 1896, ibidem (6) 18: 33, by original designation. PANAMA: Chiriqui.

N. penates is a junior subjective synonym of Laemocharis nigripes Heylaerts, 1890, Annls Soc. ent. Belg. 34 (Bull.): xiv.

NEOTUERTA Kiriakoff, 1976, Tierreich 99: 63. AGARISTIDAE Type-species: Agarista sabulosa Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.107, fig.11, by original designation. Type(s) &, [U.S.A.]: California, San Francisco, cited by Felder, 1875, ibidem 2 (Abt.2): Erklärung to pls 75-107: 1, (BMNH).

Neotuerta in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

NEOZANA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxv, 681.

Type-species: *Odozana germana* Rothschild, 1913, *Novit. zool.* 20: 197, by original designation. PERU: Carabaya, Rio Huacamayo, La Union.

NEOZATREPHES Druce, 1893, Proc. zool. Soc. Lond. 1893: 286.

Type-species: *Neozatrephes telesilla* Druce, 1893, *ibidem* **1893**: 287, pl.19, fig.5, by original designation. ECUADOR: Sarayacu.

NEPALIA Nakamura, 1974, Trans. lepid. Soc. Japan 25: 121.

Type-species: *Nepalia vesperalis* Nakamura, 1974, *ibidem* 25: 121, figs 12, 71-75, by original designation. Nepal: between Lhawn Pokn and Chitrei.

NEPE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 336.

Type-species: *Dioptis fenestra* Walker, 1854, *ibidem* 2: 337, by monotypy. Type-locality: not stated, [Brazil].

Nepe was established as a subgenus of Dioptis Hübner, 1818.

‡NEPETA; misspelling. See Napata Walker, 1854. CTEN

NEPHELIPHORA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 343.

Type-species: *Chadisra nubifera* Hampson, 1910, *Ann. Mag. nat. Hist.* (8) **5**: 460, by original designation. [Kenya]: Taveta.

NEPHELOMILTA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xviii, 87, 468.

Type-species: Lyclene suffusa Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 5, 49, pl.139,

fig.18, by original designation. [INDIA]: Nilgiri Mts, western slopes.

NEPHELOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 84, 214.

Type-species: *Lithosia caecina* Druce, 1897, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 2: 400, pl.78, fig.7, by original designation. Guatemala: San Isidro.

NEPITA Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 302. LITH

Type-species: *Pitane conferta* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 2: 533, by subsequent designation by Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* 2: 426 (for *Pitane* Walker, 1854). [SRI LANKA] Ceylon.

Nepita was established as the objective replacement name

for Pitane Walker, 1854.

PITANE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 531.

Type-species: *Pitane conferta* Walker, 1854, by subsequent designation by Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* 2: 426.

A junior homonym of *Pitane* Walker, 1854, *ibidem* 2: 462, – Lepid., Arctiidae. The objective replacement name is *Nepita* Moore, 1858.

‡NERICA; misspelling. See Nerice Walker, 1855. NOTO

NERICE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1076.

Type-species: Nerice bidentata Walker, 1855, ibidem 5: 1076, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 487. [U.S.A.]: New York, Trenton Falls.

‡NERICA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.:

An incorrect subsequent spelling of Nerice Walker, 1855.

NERICOIDES Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 35.

Type-species: Nerice bipartita Butler, 1885, Cistula ent. 3: 119, by original designation. JAPAN: Sapporo, Nikko, Yezo.

NERITONACLIA Strand, 1921, Arch. Naturgesch. 86 (A) 7: 113.

Type-species: Neritonaclia argenteogutta Strand, 1921, ibidem 86 (A) 7: 114, by original designation. GUYANA.

NERITOS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 711. ARCT

Type-species: Neritos repanda Walker, 1855, ibidem 3: 711, by monotypy. [Brazil]: Rio [de] Janeiro. See also Antiloba Weymer, 1895.

NESAEGOCERA Kiriakoff, 1974, in Kiriakoff & Viette, Faune Madagascar 39: 6 (key), 102.

AGARISTIDAE

Type-species: Aegocera comorana Jordan, 1926, Novit. zool. 33: 374, by original designation. Holotype ♀, MADAGASCAR: Comoro Is, (BMNH).

Nesaegocera in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

NESANAPHE Kiriakoff, 1969, Faune Madagascar 29: 217.

Type-species: Anaphe mirabilis Viette, 1955, Bull. mens. Soc. linn. Lyon 24: 228, by original designation. MADAGASCAR: eastern, Perinet district, [Analamazaotra] Analamazoatra forest.

NESEBRA Schaus, 1905, Proc. U.S. natn. Mus. 29: 275.

NOTO

Type-species: *Nesebra norema* Schaus, 1905, *ibidem* **29**: 275, by monotypy. French Guiana: Maroni River.

NESIOTICA Turner, 1917, Trans. Proc. R. Soc. S. Aust. 41: 54.

Type-species: *Nesiotica cladara* Turner, 1917, *ibidem* **41**: 54, by monotypy. Norfolk Island.

NESOCHADISRA Kiriakoff, 1969, Faune Madagascar 29: 13 (key), 53.

Type-species: Nesochadisra protea Kiriakoff, 1969, ibidem 29: 54, text-figs 32, 33, pl.2, figs 16, 18, by original designation. MADAGASCAR: eastern, Perinet district, Analamazaotra forest.

NESOPTILURA Kiriakoff, 1961, Lambillionea 60: 90. NOTO Type-species: Nesoptilura malgassica Kiriakoff, 1961, ibidem 60: 91, fig., by original designation. MADAGASCAR: northern, Diego Suarez district.

NESOTROPHA Turner, 1926, Pap. Proc. R. Soc. Tasm. 1925: 110. GLYPHIPTERIGIDAE

Type-species: Nesotropha pygmaeodes Turner, 1926, ibidem 1925: 110, by monotypy. Australia: Tasmania, Cradle Mt.

Nesotropha was established in the Arctiidae but is here transferred to the Glyphipterigidae (with acknowledgements to Dr I.F.B. Common who has kindly examined the type-material).

NETRIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 6: 1388 (key), 1504.

Type-species: *Netria viridescens* Walker, 1855, *ibidem* 6: 1504, by monotypy. JAVA.

NEUROXENA Kirby, 1896, Ann. Mag. nat. Hist. (6) 18: 382.

Type-species: *Neuroxena ansorgei* Kirby, 1896, *ibidem* (6) **18**: 383, pl.19, fig.2, by monotypy. UGANDA: [Entebbe] Mtebe.

NEZULA Schaus, 1896, Jl N.Y. ent. Soc. 4: 137. ARCT Type-species: Nezula grisea Schaus, 1896, ibidem 3: 137, by monotypy. VENEZUELA: Aroa.

NIASANA Roepke, 1937, Ent. Z., Frankf.a.M. 51: 194. ARCT Type-species: Arctia dehanna Pagenstecher, 1885, Jb. nassau. Ver. Naturk. 38: 14, pl.1, fig.10, by original designation. [INDONESIA]: Sumatra, Nias [I.].

NICAEA Hampson, 1901. See Nikaea Moore, 1879. ARCT

‡NICHTHEMERA; misspelling. See Nyctemera Hübner, [1820].

NIGANDA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 63.

Type-species: Niganda strigifascia Moore, 1879, ibidem (1): 63, pl.3, fig.15, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 580. INDIA: Darjeeling.

NIKAEA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 11.

Type-species: Hypercompa longipennis Walker, 1855, Lis

Type-species: *Hypercompa longipennis* Walker, 1855, *List Specimens lepid. Insects Colln Br. Mus.* 3: 655, by monotypy. [BANGLADESH]: [Sylhet] Silhet.

NICAEA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 5, 218.

An unjustified emendation of Nikaea Moore, 1879.

A junior homonym of *Nicaea* Robineau-Desvoidy, 1863, – Insecta, Diptera. The objective replacement name is *Nikaea* Moore, 1879.

NILGIRICOLA Hampson, 1918, Novit. zool. 25: 104. LITH Type-species: Nilgiricola sicciana Hampson, 1918, ibidem 25: 105, by original designation. [INDIA]: Madras, Nilgiris, Pykara.

NIODA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1069.

Type-species: *Nioda fusiformis* Walker, 1855, *ibidem* 5: 1070, by monotypy. [SRI LANKA] Ceylon, INDIA: North.

Nioda was established in the Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 488.

NIPPONASURA Inoue, 1965, Kontyû 33: 241. LITH Type-species: Nipponasura sanguinea Inoue, 1965, ibidem 33: 242, figs 1, 2, by original designation. JAPAN: Ryukyu Islands, Ishigaki Is, Ka-arayama.

NIPPONOTENSHA Nakamura, 1972, Ent. Rev. Japan 24: 54, 55.

Type-species: *Pseudofentonia amamiana* Nakamura, 1972, *ibidem* **24**: 54, fig.1, by original designation. Japan: Amami-Oshima Island, Koniya.

Nipponotensha was established as a subgenus of Pseudofentonia Strand, 1912.

NISHADA Moore, 1878, Proc. zool. Soc. Lond. 1878: 23.

Type-species: *Nishada flabrifera* Moore, 1878, *ibidem* **1878**: 23, by monotypy. [INDIA]: Calcutta district.

NOCTHADENA Laporte, 1976, Bull. mens. Soc. linn. Lyon 45: 136. NOCTUIDAE

Type-species: *Nocthadena griseoviridis* Laporte, 1976, *ibidem* 45: 136, figs 5, genitalia 2, by monotypy. Holotype 3, ETHIOPIA: environs du Mont Batu, (MNHN, Paris).

Nocthadena in the Noctuidae, Hadeninae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

NOCTULODES Schaus, 1937, Proc. U.S. natn. Mus. 84: 572.

NOTO

Type-species: *Noctulodes porpara* Schaus, 1937, *ibidem* 84: 573, by original designation. BRAZIL: Itatiaya.

NODOZA Schaus, 1896, Jl N.Y. ent. Soc. 4: 150. CTEN Type-species: Nodoza tristis Schaus, 1896, ibidem 4: 150, by monotypy. Brazil: São Paulo.

N. tristis is a junior subjective synonym of Delphyre hebes Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2:

See also Delphyre Walker, 1854.

NODOZANA Druce, 1899, Ann. Mag. nat. Hist. (7) 3: 467.

Type-species: *Nodozana xanthomela* Druce, 1899, *ibidem* (7) 3: 467, by monotypy. [BRAZIL]: Amazon valley.

The name *Nodozana* probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson, 1900, as the name for a new genus.

NODOZANA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 88, 375.

Type-species: *Nodozana thricophora* Hampson, 1900, *ibidem* 2: 375, fig.286, by original designation. Panama: Chiriqui, Caldera

A junior homonym of *Nodozana* Druce, 1899. There is no objective replacement name but the type-species of *Nodozana* Hampson, 1900, is currently considered to be congeneric with the type-species of *Nodozana* Druce, 1899. The latter is therefore available for use as a subjective replacement name.

NODOZANA Hampson, 1900. See Nodozana Druce, 1899.

NOLA Leach, [1815] 1830, in Brewster, Edinburgh Encycl. 9: 135.

Type-species: *Noctua palliola* [Denis & Schiffermüller], 1775, *Ankündung syst. Werkes Schmett. Wienergegend*: 69 (but cited by Leach as ‡*palliolatis*, an incorrect subsequent spelling, by monotypy. [AUSTRIA]: Vienna district.

N. palliola is a junior subjective synonym of Phalaena cucul-

latella Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 537.

Nola has been used as the original generic name for species now placed in the Arctiidae.

NOLECA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1079.

Type-species: *Noleca basigutta* Walker, 1855, *ibidem* 5: 1080, by monotypy. Type-locality: not known, [NIGERIA].

Noleca was established in the Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 489.

NOLINOPHANES Hulstaert, 1924, Ann. Mag. nat. Hist. (9) 13: 132.

Type-species: *Nolinophanes sicciaecolor* Hulstaert, 1924, *ibidem* (9) **13**: 133, by original designation. [INDONESIA]: [Tanimbar Is] Tenimber, Olilit.

NOLIPROCTIS Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 152.

Type-species: *Dasychira pumila* Butler, 1882, *Cistula ent.* 3: 16 (a junior primary homonym, see below), by monotypy. MADAGASCAR: Ankafana.

D. pumila Butler, 1882, is a junior primary homonym of Dasychira pumila Staudinger, 1881, Stettin. ent. Ztg 42: 405. There is no replacement name available so Noliproctis milupa Nye nom. n. is here established for it.

The date of publication of the part of Seitz containing page 152 was stated on page 145 as 4.xi.1926 in both the German and the English editions.

NOLOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 385.

Type-species: *Nolosia marmorata* Hampson, 1900, *ibidem* 2: 385, fig.298, by original designation. MADAGASCAR: Betsileo.

Nolosia was established in the 'Arctiadae'; it was transferred to the Lymantriidae by Collenette, 1959, Annls Soc. ent. Fr. 128: 108.

‡NONODONTA; misspelling. See Notodonta Ochsenheimer, 1810.

NORRACA Moore, 1881, Proc. zool. Soc. Lond. 1881: 340.

Type-species: Norraca longipennis Moore, 1881, ibidem 1881: 340, by monotypy. [MALAYSIA] Malaya: Penang.

Norraca was established in the Noctuidae; it was placed in the Notodontidae by Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 137.

NORRACANA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 153 (key), 205; 1962 October 20, Bonn. zool. Beitr. 13: 235.

Type-species: *Bireta niveipicta* Kiriakoff, 1962 [July 31], *ibidem* 98 (10): 205; 1962 October 20, *ibidem* 13: 235, abb.2, fig.20, pl.2, fig.19 (as *Norracana niveipicta* sp.nov.), by original designation. China: Fukien, [Kuatun]. Additional data cited by Kiriakoff, 1962 October 20, *ibidem* 13: 236.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr. to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper B. niveipicta was described as a new species in the combination Norracana niveipicta and Norracana

was proposed as a new genus. As the dates show, the Belgian paper appeared first. In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

Norracana was first published as a subgenus of Bireta Walker, 1856; it was treated as a genus by Kiriakoff, 1962 October 20, ibidem 13: 235, and again in 1967, in Wytsman, Genera Insect. 217 (B): 54.

NORRACOIDES Strand, 1916, Arch. Naturgesch. 81 (A) 12: 156.

Type-species: *Norracoides discocellularis* Strand, 1916, *ibidem* 81 (A) 12: 156, by original designation. Talwan: Kosempo.

N. discocellularis is a junior subjective synonym of Notodonta basinotata Wileman, 1910, Entomologist 43: 344.

NOTARCTIA Smith, 1938, Revision of the Genus Apantesis Walker. Univ. Illinois: 7.

Type-species: *Chelonia proxima* Guérin-Méneville, [1844] 1829-1838, *Icon. Règne anim. Cuvier* 3 (Insectes): 514, by original designation. [MEXICO]: [Orizaba] Orixaba.

NOTATA Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 4, 47.

Type-species: *Notata parva* Hampson, 1891, *ibidem* 8: 48, pl.139, fig.5, by monotypy. INDIA: Nilgiri plateau.

NOTELA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 263 (key), 324.

Type-species: *Notela jaliscana* Schaus, 1901, *ibidem* 1901: 324, pl.12, fig.3, by original designation. MEXICO: Guadalajara.

NOTHARCTIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 42.

Type-species: *Metarctia flavivena* Hampson, 1901, *Ann. Mag. nat. Hist.* (7) 8: 169, by original designation. [Kenya] Br. E. Africa: Machakos.

Notharctia was established as a subgenus of Metarctia Walker, 1855.

NOTOCERURA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 208.

Type-species: Cerura spiritalis Distant, 1899, Ann. Mag. nat. Hist. (7) 3: 464, by original designation. [SOUTH AFRICA]: Transvaal, Lydenburg district.

NOTODON Meigen, 1830. See Notodonta Ochsenheimer, 1810.

NOTODONTA Ochsenheimer, 1810, Schmett. Eur. 3: 45.

Type-species: *Phalaena dromedarius* Linnaeus, 1767, *Syst. Nat.* (Edn 12) 1 (2): 827, by subsequent designation by Curtis, 1839, *Br. Ent.* 16: 739. Type-Locality: not stated, [Europe].

Invalid type-species designations: *Phalaena dromedarius* Linnaeus, 1767, was designated by Boisduval, 1836, *Hist. nat. Insectes* (Spec. gén. Lépid.) 1: 129.

Phalaena ziczac Linnaeus, 1758, was also designated by Boisduval, 1836, *ibidem* 1: 134, 142.

In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different type-species for each generic name. In his 'Exposé de notre méthode', pages 155-690, no type-species designation was made for any of the genera that he

himself used. Under the *Code*, Article 69(a)(iii), the type-species designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other authors.

NOTODON Meigen, 1830, Syst. Beschreibung eur. Schmett. 2: 171.

An unjustified emendation of *Notodonta* Ochsenheimer, 1810.

‡NONODONTA; Druitt, 1900, Entomologist 33: 202.

An incorrect subsequent spelling of *Notodonta* Ochsenheimer, 1810.

‡NOTODONTULA; Nordström, 1930, Mitt. münch. ent. Ges. 20: 93.

An incorrect subsequent spelling of *Notodonta* Ochsenheimer, 1810.

NOTODONTELLA Roepke, 1943, Natuurh. Maandbl. 32: 78.

Type-species: *Notodontella nieuwenhuisi* Roepke, 1943, *ibidem* 32: 79, fig.7, by monotypy. Java: eastern, Mt Ardjuno, Diunggo.

Notodontella was established in the Notodontidae; it was transferred provisionally to the Thyatiridae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2. Examination of the type specimen of Notodontella nieuwenhuisi has shown it to be closely related to Fentonia ferrifusa Dudgeon, 1898, J. Bombay nat. Hist. Soc. 11 (4): 634, currently included in the genus Fentonia Butler, 1881, in the Notodontidae. Notodontella was restored to the Notodontidae by Fletcher, 1979, in Nye, Generic Names Moths World 3.

‡NOTODONTULA; misspelling. See Notodonta Ochsenheimer, 1810.

NOTOHYBA Holland, 1893, Psyche, Camb. 6: 434, 451.

Type-species: *Notohyba nubifuga* Holland, 1893, *ibidem* 6: 451, 568, pl.18, fig.3, by original designation. [GABON]: [Ogooué] Ogové R. valley.

NOTOLOPHUS Germar, 1812. See Orgyia Ochsenheimer, 1810.

NOTOPHYSON Boisduval, 1870, Considérations Lépid. Guatemala: 87.

Type-species: *Phalaena tiresias* Cramer, 1776, *Uitlandsche Kapellen (Papillons exot.)* 1: 133, 154, pl.85, fig.B, by subsequent designation by Bryk, 1931, *in* Strand, *Lepid. Cat.* 45: 37. SURINAM.

P. tiresias is dated from the wrapper of the part and not from the title-page of the volume, 1779.

ANTHOMYZA Swainson, 1833, Zool. Illust. (2) 3: legend to pl.124.

Type-species: *Phalaena tiresias* Cramer, 1776, by subsequent designation by Duponchel, 1841, *in* d'Orbigny, *Dict. univl Hist.*

A junior homonym of *Anthomyza* Fallén, 1810, *Specimen ent. novam Diptera* . . .: 20, – Insecta, Diptera. The objective replacement name is *Notophyson* Boisduval, 1870.

NOTOPLUSIA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 261 (key), 300.

Type-species: *Phalaena clara* Stoll, 1780, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 43, pl.311, fig.B, by original designation. SURINAM.

P. clara is dated from the wrapper of part 26, not from the title-page of volume 4, which is dated 1782.

‡NOTOPRIOTA Swinhoe, 1903. See Oecura Holland, 1893.

NOTOTRACHUS Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 245.

Type-species: *Nototrachus pierreti* Rambur, 1866, *ibidem* (2): 245, by monotypy. ALGERIA.

NOTOXANTHA Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 484.

Type-species: *Notoxantha sesamiodes* Hampson, 1910, *ibidem* (8) 5: 485, by original designation. [RHODESIA]: Salisbury.

NOVOSIA Toulgoët, 1958, Mém. Inst. scient. Madagascar (E) 9: 192.

Type-species: *Philenora herbuloti* Toulgoët, 1954, *ibidem* (E) 5: 200, text-fig.18, pl.9, figs 28, 29, by original designation. MADAGASCAR: central, massif de l'Ankaratra, Manjakatompo, Ambahona forest.

NUDARIA Haworth, [1809] 1803, Lepid. Br.: 156. LITH Type-species: Phalaena mundana Linnaeus, 1761, Fauna Suecica (Edn 2): 349 (but cited as ‡mundus, an incorrect subsequent spelling by Haworth), by subsequent designation by Curtis, 1832, Br. Ent. 9: 400 (but cited as ‡munda, an incorrect subsequent spelling). Sweden.

NELOPA Billberg, 1820, Enumeratio Insect. Mus. G.J. Billberg: 85.

Type-species: *Phalaena mundana* Linnaeus, 1761, by monotypy (but cited as ‡*muudana*, an incorrect subsequent spelling by Billberg).

A junior objective synonym of Nudaria Haworth, [1809].

AMALEA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 100, 101 (but attributed to Zetterstedt, an incorrect authorship); [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 20, 21.

Type-species: *Phalaena mundana* Linnaeus, 1761, by monotypy (but included by Herrich-Schäffer as *munda* [Fabricius, 1702] an univetified amendation)

1793], an unjustified emendation).

Amalea was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

TRICHOTA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 196.

Type-species: Phalaena mundana Linnaeus, 1761, by monotypy.

A junior objective synonym of Nudaria Haworth, [1809].

DERRHIS Wallengren, 1885, Skand. Heterocer-Fjärilar 2: 259.

Type-species: *Phalaena mundana* Linnaeus, 1761, by monotypy.

A junior homonym of *Derrhis* Agassiz, 1847, *Nomencl. zool*. (Index univl.): 120, – Vermes. The objective replacement name is *Nudaria* Haworth, [1809].

NUDARIDIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvii, 86, 420.

Type-species: *Nudaria muscula* Staudinger, 1887, *in* Romanoff, *Mêm. Lépid.* 3: 185, pl.10, fig.9, by original designation. [U.S.S.R.]: Vladivostok and Sidemi.

NUDINA Staudinger, 1887, in Romanoff, Mém. Lépid. 3: 186, 187.

Type-species: Nudaria nubilosa Staudinger, 1887, ibidem 3: 186, pl.10, fig.10, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 468, but cited as Miltochrista artaxidia Butler, 1881, a nominal species not originally included in Nudina. Hampson, however, placed artaxidia as the senior synonym of nubilosa, a nominal species originally included in Nudina. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. [U.S.S.R.]: Vladivostok.

N. nubilosa is a junior subjective synonym of Miltochrista artaxidia Butler, 1881, Trans. ent. Soc. Lond. 1881: 8.

NUDOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 88, 337.

Type-species: *Nudosia fuscifusa* Hampson, 1900, *ibidem* 2: 337, fig.238, by original designation. PERU.

NUDUR Dyar, 1914, Proc. U.S. natn. Mus. 47: 368. LITH Type-species: Nudur fractivittarum Dyar, 1914, ibidem 47: 368, by original designation. MEXICO: Misantla.

NUMENES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 586 (key), 662.

Type-species: *Numenes siletti* Walker, 1855, *ibidem* 3: 663, by monotypy. [Bangladesh]: [E. Bengal, Sylhet] Silhet, [INDONESIA]: Java.

Numenes was established in the Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 455.

NUMENOIDES Butler, 1879, Ann. Mag. nat. Hist. (5) 4: 238.

Type-species: *Numenoides grandis* Butler, 1879, *ibidem* (5) 4: 238, by original designation. MADAGASCAR: Antananarivo.

NUNUA Karsch, 1895, Ent. Nachr. Berlin 21: 365.

Type-species: Nunua gallans Karsch, 1895, ibidem 21: 365, pl.4, fig.8, by original designation. Togo: Bismarckburg.

‡NYCHTHEMERA; misspelling. See Nyctemera Hübner, [1820].

‡NYCTEMEMERA; misspelling. See Nyctemera Hübner, [1820].

NYCTEMERA Hübner, [1820] 1816, Verz. bekannter Schmett.: 178.

Type-species: *Phalaena lacticinia* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 47, 149, pl.128, fig.E, by subsequent designation by Hampson, 1894, *Fauna Br. India* (Moths) 2: 46. [INDIA]: Coromandel coast.

P. lacticinia is dated from the wrapper of the part and not from the title-page of the volume, 1779.

Nyctemera was placed incorrectly in the Lymantriidae by Bryk, 1934, in Strand, Lepid. Cat. 62: 199. This error was corrected by Collenette, 1935, Stylops 4: 242.

LEPTOSOMA Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 197.

Type-species: *Leptosoma annulatum* Boisduval, 1832, *ibidem* 1: 197, pl.5, fig.9, by subsequent designation by Meyrick, 1912, *Trans. N.Z. Inst.* 44: 93. New Guinea.

A junior homonym of *Leptosoma* Desmarest, 1823, *Dict. Sci. nat.* 28: 421, – Crustacea. Gistl, 1848, *Naturg. Thierreichs*:

ix, established the name Amathes Gistl as an objective replacement name for Leptosoma Boisduval (but cited by Gistl as ‡Leptosomus Quoy, an incorrect subsequent spelling and authorship), but Amathes Gistl, 1848, is itself preoccupied by Amathes Hübner, [1821] 1816, Verz. bekannter Schmett.: 222, – Lepid., Noctuidae. There is no other objective replacement name for Leptosoma Boisduval, 1832, but its type-species is currently considered to be congeneric with the type-species of Nyctemera Hübner, [1820]. The latter is therefore available for use as a subjective replacement name.

Invalid type-species designation: Leptosoma insulare Boisduval, 1833, a nominal species not originally included in Leptosoma and not linked in synonymy with one of the originally included nominal species when designated by Desmarest, (1857), in Chenu, Encycl. hist. Nat. (Papillons nocturnes): 34.

AMATHES Gistl, 1848, Naturg. Thierreichs: ix.

Established as an objective replacement name for *Leptosoma* Boisduval, 1832 (but cited as ‡*Leptosomus* Quoy, an incorrect subsequent spelling and incorrect authorship).

A junior homonym of *Amathes* Hübner, [1821] 1816, *Verz. bekannter Schmett.*: 222, – Lepid., Noctuidae.

‡LEPTOSOMUS; Gistl, 1848, Naturg. Thierreichs: ix.

Gistl established *Amathes* as an objective replacement name for ‡*Leptosomus* Quoy, an incorrect subsequent spelling and incorrect authorship of *Leptosoma* Boisduval, 1832.

‡NYCTHEMERA; Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 98.

An incorrect subsequent spelling of *Nyctemera* Hübner, [1820].

‡Nycthemera was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

‡NICHTHEMERA; Mabille, 1879, Bull. Soc. zool. Fr. 3: 87. An incorrect subsequent spelling of Nyctemera Hübner, [1820].

‡NYCHTHEMERA; Mabille, 1879, Annls Soc. ent. Fr. (5) 9: 304.

An incorrect subsequent spelling of *Nyctemera* Hübner, [1820].

ZONOSOMA Butler, 1881, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 5: ix, 44.

Type-species: Nyctemera interlecta Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 400, by original designation. [Bangladesh]: [Sylhet] Silhet.

A junior homonym of Zonosoma Lederer, 1853, Verh. zool.-bot. Ver. Wien 3: 194, – Lepid., Geometridae. Tristania Kirby, 1892, was established as the objective replacement name but is itself preoccupied by Tristania Böttger, 1878. There is no objective replacement name but the type-species of Zonosoma Butler, 1881, is currently considered to be congeneric with the type-species of Nyctemera Hübner, [1820]. The latter is therefore available for use as a subjective replacement name.

TRISTANIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 423.

Tristania was established as the objective replacement name for Zonosoma Butler, but is itself preoccupied by Tristania Böttger, 1878, Offenbacher Ver.-Naturk 17-18: 19, – Mollusca.

‡NYCTEMEMERA; Pagenstecher, 1901, Jb. nassau. Ver. Naturk. 54: 111.

An incorrect subsequent spelling of *Nyctemera* Hübner, [1820].

‡NYCTEMERIA; Matsumura, 1931, 6000 Illust. Insects Japan.-Empire: 973.

An incorrect subsequent spelling of *Nyctemera* Hübner, [1820].

‡NYCTEMERIA; misspelling. See Nyctemera Hübner, [1820].

NYCTEROPA Turner, 1941, Mem. Qd Mus. 12: 40. THAU Type-species: Nycteropa subovalis Turner, 1941, ibidem 12: 41, by monotypy. AUSTRALIA: Queensland, Injune.

Nycteropa was established in the Notodontidae; it was transferred to the Thaumetopoeidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

NYCTEROTIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.20. NOTO Type-species: Nycterotis poecila Felder, 1874, ibidem 2 (Abt.2): pl.97, fig.20, by monotypy. VENEZUELA: cited by

Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 7.

‡NYCTHEMERA; misspelling. See Nyctemera Hübner, [1820].

NYCTOCHROA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.27. LITH Type-species: Nyctochroa basiplaga Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.27, by monotypy. MEXICO: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 7.

NYCTOSIA Schaus, 1899, Jl N.Y. ent. Soc. 7: 217. LITH Type-species: Nyctosia coccinea Schaus, 1899, ibidem 7: 217, by monotypy. MEXICO: Mexico City.

The name *Nyctosia* probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson as the name for a new genus.

NYCTOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 90, 253.

Type-species: Apistosia tenebrosa Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1881, by original designation. Type-locality: not stated, [Central America].

A junior homonym of *Nyctosia* Schaus, 1899, – Lepid., Lithosiidae. There is no objective replacement name but the type-species of *Nyctosia* Hampson, 1900, is currently considered to be congeneric with the type-species of *Nyctosia* Schaus, 1899. The latter is therefore available for use as a subjective replacement name.

NYCTOSIA Hampson, 1900. See Nyctosia Schaus, 1899.

NYEARCTIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 66.

Type-species: Automolis leucoptera Hampson, 1920, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 2: 144, by original designation. GUYANA: 'bought at Georgetown'.

NYGMIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 193.

Type-species: *Phalaena icilia* Stoll, 1790, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* (Aanhangsel): 158, 183, pl.35, fig.5, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 448. [INDIA]: Coromandel coast, Tranquebar.

NYMPIIYXIS Grote, 1895. See Leucoma Hübner, 1822.

NYRIDELA Lucas, 1857, in Sagra, Hist. phys. politique nat. Ile Cuba [2] (Anim. articulés): 663.

Type-species: *Isanthrene chalciope* Hübner, [1831] 1825, *Zuträge Samm!. exot. Schmett.* 3: 20, pl.[81], figs 469, 470, by monotypy. CUBA: [Havana] Havannah.

See also Mochloptera Butler, 1876.

NYSSIOPSIS Kiriakoff, 1959, Ark. Zool. (2) 12: 325. NOTO
Type-species: Nyssiopsis mystacina Kiriakoff, 1959, ibidem
(2) 12: 325, text-fig.25, pl.1, fig.9, by original designation.
[Burma]: Kambaiti.

NYSTALEA Guenée, 1852, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 6: 122.

Type-species: Nystalea conchyfera Guenée, 1852, ibidem 6: 122, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 618. [WEST INDIES]: [Virgin Islands], St Thomas Island.

N. conchyfera is a junior subjective synonym of Phalaena ebalea Stoll, 1780, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 41, pl.310, fig.C.

Nystalea was established in the Noctuidae; it was placed in the Notodontidae by Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 693.

OCHROCALLIOPE Kiriakoff, 1958, Mém. Inst. scient. Madagascar (E) 9: 218.

Type-species: Ochrocalliope viettei Kiriakoff, 1958, ibidem (E) 9: 218, text-figs 2, 3, pl.4, figs 1, 3, by original designation. MADAGASCAR: Mananjary.

OCHRODOTA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 5, 89.

Type-species: Zatrephes pronapides Druce, 1894, Ann. Mag. nat. Hist. (6) 13: 173, by original designation. Panama: Bugaba.

OCHROGASTER Herrich-Schäffer, [1855 June] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.66, fig.377; [1856], ibidem 1: 12, 67; 1855 December, Syst. Bearbeitung Schmett. Eur. 6: 92.

Type-species: *Ochrogaster lunifer* Herrich-Schäffer, [1855 June], *ibidem* 1 (1): wrapper, pl.66, fig.377, by monotypy. AUSTRALIA.

Ochrogaster was established in the Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 468; it was transferred to the Eupterotidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498; and was placed in the Cnethocampinae, now Thaumetopoeidae by Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 371.

See also Teara Walker, 1855 [August].

‡OCHROGASTER Neave, 1940, Nomencl. zool. 3: 374.

In addition to *Ochrogaster* Herrich-Schäffer, [1855], Neave also cited *Ochrogaster* Felder, 1874, *Reise öst. Fregatte Novara* (Zool.) **2** (Abt.2): pl.94, fig.5, pl.95, fig.9; 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 7. Felder was not however establishing a new genus, but using *Ochrogaster* Herrich-Schäffer, [1855].

‡OCHROGASTER Neave, 1940. See Ochrogaster Herrich-Schäffer, [1855].

OCHROSOMERA Kiriakoff, 1969, Faune Madagascar 29:

Type-species: *Desmeocraera marojejia* Kiriakoff, 1963, *Revue fr. Ent.* 30: 290, fig.3, by original designation. MADA-GASCAR: eastern, Sambava district, Marojejy, Beondroka.

OCHROSTIGMA Hübner, [1819] 1816, Verz. bekannter Schmett.: 146. Type-species: Bombyx austera [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 310, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 602, but cited as velitaris Hufnagel, 1766, a nominal species not originally included in Ochrostigma. However, Kirby placed velitaris as a senior subjective synonym of austera [Denis & Schiffermüller], 1775, a nominal species originally included in Ochrostigma. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. [AUSTRIA]: Vienna district.

B. austera is a junior subjective synonym of Phalaena velitaris Hufnagel, 1766, Berlin. Mag. 3: 394.

See also Drynobia Duponchel, [1845].

OCHROTA Hopffer, 1862, in Peters, Naturw. Reise Mossambique (Zool.) 5: 427.

Type-species: Crocota unicolor Hopffer, 1858, Mber. K. preuss. Akad. Wiss. 1857: 422, by monotypy. MOZAMBIQUE. See also Bettonia Butler, 1898.

OCNERIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 158.

Type-species: *Bombyx rubea* [Denis & Schiffermüller], 1775, *Ankündung syst. Werkes Schmett. Wienergegend*: 51, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 466. [Austria]: Vienna district.

‡OENERIA; Ramann, [1875], Schmett. Dtl. angrenzenden Länder: 177.

An incorrect subsequent spelling of Ocneria Hübner, [1819].

OCNEROGYIA Staudinger, 1892, Dt. ent. Z. Iris 4: 254.

Type-species: Ocnerogyia amanda Staudinger, 1892, ibidem 4: 254, by monotypy. [Turkey]: [Kurdistan], Mardin.

OCNEROPSIS Kozhanchikov, 1950. See Gissarus Nye nom.

OCNOGYNA Lederer, 1853, Verh. zool.-bot. Ver. Wien 2 (Abh.): 78.

Type-species: *Chelonia zoraida* Graslin, [1837] 1836, *Annls Soc. ent. Fr.* 5: 561, pl.17B, fig.3, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 273. Spain: Andalusia, Sierra Nevada.

OCROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 87, 268.

Type-species: *Lithosia reversa* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 105, by original designation. BORNEO: Sarawak. See also *Tampea* Snellen, 1898.

OCYBOLA Turner, 1912, Mem. natn. Mus. Melb. 4: 17, 20.

Type-species: Orgyia semifusca Walker, 1869, Characters undescr. Lepid. Heterocera: 63, 65, by original designation. AUSTRALIA.

ODAGRA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 401.

Type-species: Odagra devestita Walker, 1865, ibidem 32: 402, by monotypy. [INDIA]: [W. Bengal], Darjeeling.

ODNARDA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 153 (key), 174.

Type-species: *Bireta subserena* Kiriakoff, 1962, *ibidem* 98: 174, fig.23, by original designation. BURMA: Fort Hpimaw, near Myitkyina.

Odnarda was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 44.

ODONPTERA Latreille, 1829. See Euchila Billberg, 1820.

ODONTOPERAS Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 165.

Type-species: *Odontoperas archonta* Kiriakoff, 1959, *ibidem* 95: 166, fig.11, by original designation. SIERRA LEONE.

ODONTOSIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 145.

Type-species: *Phalaena carmelita* Esper, [1799], *Die Schmett*. 3 (Suppl.) Abschnitt 3: 65, pl.91, fig.1, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 604. [Germany]: Gunsenhausen.

ODONTOSIANA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 288.

Type-species: *Odontosiana schistacea* Kiriakoff, 1964, *ibidem* 14: 288, abb.5, fig.49, pl.5, fig.52, by original designation. China: Shansi, Mien-Shan.

ODONTOSINA Gaede, 1933, in Seitz, Gross-Schmett. Erde 2 (Suppl.): 182.

Type-species: *Odontosina nigronervata* Gaede, 1933, *ibidem* **2** (Suppl.): 183, pl.15 row f, by monotypy. [China]: [western], Tse-ku.

The date of publication of the part of Seitz containing page 182 was cited on page 177 as 22.iii.1933 in both the German and the English editions.

ODOXIA Dognin, 1923, Hétérocères nouv. Am. Sud 21: 10.

Type-species: *Odoxia concocta* Dognin, 1923, *ibidem* **21**: 11, by original designation. FRENCH GUIANA.

ODOZANA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 109.

Type-species: Odozana floccosa Walker, [1865] 1864, ibidem 31: 110, by monotypy. [Brazil]: [R. Amazon], [Tefé] Ega.

OECURA Holland, 1893, Psyche, Camb. 6: 385, 389. LYMA Type-species: Oecura goodii Holland, 1893, ibidem 6: 390, pl.9, figs 1, 3, 4, by monotypy. WEST AFRICA: tropical (see page 385).

‡NOTOPRIOTA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 466.

‡Notopriota is an unavailable name under the Code, Article 11(d). It was first published by Swinhoe in the synonymy of Oecura ocellifera Holland, 1893, and has not subsequently been treated as an available name.

OEDALEOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 85, 219.

Type-species: *Oedaleosia nigricosta* Hampson, 1900, *ibidem* 2: 219, fig.138, by original designation. SIERRA LEONE, GAMBIA.

OEDEMASIA Packard, 1864, Proc. ent. Soc. Philad. 3: 359.

Type-species: *Phalaena concinna* Smith, 1797, *in* Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 169, pl.85, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 567. [U.S.A.]: Georgia.

OENARCTIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 51.

Type-species: *Metarctia erlangeri* Rothschild, 1910, *Novit. zool.* 17: 442, by original designation. [ETHIOPIA] Abyssinia: Arussi Galla Country, Djedda to Sibbe.

Oenarctia was established as a subgenus of Metarctia Walker, 1855.

‡OENERIA; misspelling. See Ocneria Hübner, [1819]. LYMA

‡OENISTIS; misspelling. See Oeonistis Hübner, [1819].

OENOSANDA Walker, 1856 [May 10]. See Oenosandra Walker, 1856 [April 21].

OENOSANDRA Walker, 1856 [April 21], in Newman, Trans. ent. Soc. Lond. (2) 3 (8): 286.

Type-species: Oenosandra boisduvalii Newman, 1856 [April 21], ibidem (2) 3 (8): 286, by monotypy. AUSTRALIA: Victoria, Mt Alexander Range, 80 miles from Melbourne.

Newman, 1856, ibidem (2) 3 (8): 283 stated that the generic names he had adopted were those of Walker and attributed them to him. Three weeks after the publication of Newman's paper, Walker, 1856 [May 10], List Specimens lepid. Insects Colln Br. Mus. 7: 1713 redescribed Oenosandra boisduvalii, emending the generic name to Oenosanda and replacing the specific name with duponchelii, which he erroneously attributed to Newman. Wheeler, 1911, Trans. ent. Soc. Lond. 1911: 755 stated that Trans. ent. Soc. Lond. (2) 3 (8), the part in which Newman's paper appeared, was received from the printers on 21 April 1856. Nye, 1975, Generic Names Moths World 1: 6 stated that Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7 was available for issue on 10 May 1856.

Oenosandra was established in the Bombycidae; placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 468; it was placed in the Notodontidae by Turner, 1903, Proc. Linn. Soc. N.S.W. 28: 57; and was subsequently placed in the Cnethocampinae, now Thaumetopoeidae, by Turner, 1922, ibidem 47: 373-4.

See also Lomatosticha Möschler, 1872.

OENOSANDA Walker, 1856 [May 10], List Specimens lepid. Insects Colln Br. Mus. 7: 1713.

An unjustified emendation of Oenosandra Walker, 1856 [April 21].

PTERYGOSOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.98, fig.7.

Type-species: Pterygosoma squamipunctum Felder, 1874, ibidem 2 (Abt.2): pl.98, fig.7, by monotypy. AUSTRALIA: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 8.

A junior homonym of Pterygosoma Peters, 1849 July 24, Vossische Ztg (Königl. privilegirte Berl. Ztg) 24.vii.1849 (Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin), - Arachnida. There is no objective replacement name, but P. squamipunctum is a junior subjective synonym of Oenosandra boisduvalii Newman, 1856, the type-species of Oenosandra Walker, 1856; the latter is therefore available for use as a subjective replacement name. The accounts of the meetings of the Gesellschaft Naturforschender Freunde zu Berlin were not published by the Society between 1839 and 1859, but were recorded in the Vossische Zeitung. The Society subsequently published a summary of all these meetings in one volume, 1912, Sber. Ges. naturf. Freunde Berl. 1839-1859; the description of Pterygosoma and its type-species was reprinted on page 87.

Pterygosoma Felder, 1874, when it was established was not placed in a family; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 591; and was placed in the Cnethocampinae, now Thaumetopoeidae, by

Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 374.

‡OENOSIA; misspelling. See Oeonosia Hampson, 1914.

‡OEOLOSIA; misspelling. See Aeolosia Hampson, 1900.

OEONISTIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 165.

Type-species: Phalaena entella Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 27, pl.208, fig.D, by subsequent designation by Moore, 1878, Proc. zool. Soc. Lond. 1878: 14. [INDIA]: Coromandel coast.

P. entella is dated from the wrapper of the part and not from the title-page of the volume, 1782.

‡OENISTIS; Kirby, 1892, Synonymic Cat. Lepid. Heterocera

An incorrect subsequent spelling of *Oeonistis* Hübner, [1819], also used by Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 115.

PHILAGRIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera

Type-species: Phalaena entella Cramer, 1779, by original designation.

A junior objective synonym of Oeonistis Hübner, [1819].

OEONOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xx, 513.

Type-species: Oeonistis aurifera Rothschild, 1912, Novit. zool. 19: 225, by original designation. New Guinea: [Irian Jaya], Snow Mts, near Oetakwa River.

‡OENOSIA; Neave, 1940, Nomencl. zool. 3: 395. An incorrect subsequent spelling of *Oeonosia* Hampson, 1914.

OGLANA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 109.

Type-species: Pydna discoidalis Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 621, pl.82 row b, by original designation. [VIETNAM]: Cochin.

OGOA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1763.

Type-species: Ogoa simplex Walker, 1856, ibidem 7: 1764, by monotypy. [South Africa]: [Durban] Port Natal.

Ogoa was established in the Drepanulidae, now Drepanidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 437.

OGOVODONTA Gaede, 1928, in Seitz, Gross-Schmett. Erde **14**: 438.

Type-species: Ogovodonta transversa Gaede, 1928, ibidem 14: 438, pl.72 row c, by monotypy. [GABON]: [Ogooué R.]

The date of publication of the part of Seitz containing page 438 was cited on page 433 as 30.v.1928 in the German edition, and as 26.xi.1928 in the English edition.

OGULINA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98:

Type-species: Pydna plusioides Bryk, 1950, Ark. Zool. 42 (A) 19: 17, pl.4, fig.7, by original designation. BURMA: northern, Kambaiti.

Ogulina was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 44.

OLAPA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 823.

Type-species: Olapa temperata Walker, 1855, ibidem 4: 823, by monotypy. South Africa.

OLCECLOSTERA Butler, 1878, Trans. ent. Soc. Lond. 1878: APATELODIDAE

Type-species: Olceclostera irrorata Butler, 1878, ibidem 1878: 70, by original designation. [BRAZIL]: Amazon region, Boa Vista, Rio Jutahi.

Olceclostera was established in the Notodontidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 852; and was placed in the Zanolinae, now Apatelodidae by Draudt, 1929, in Seitz, Gross-Schmett. Erde 6: 703.

OLENE Hübner, 1823, Zuträge Samml. exot. Schmett. 2: 19.

LYMA

Type-species: Olene mendosa Hübner, 1823, ibidem 2: 19, pl.[51], figs 293, 294, by subsequent designation by Moore, [1883] 1882-3, Lepid. Ceylon 2: 96. JAVA.

See also Turriga Walker, 1869.

‡OLENOPTERA Herrich-Schäffer, 1855. See Olenoptera Herrich-Schäffer, [1856].

OLENOPTERA Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): 11, 84, wrapper, pl.91, fig.523.

Type-species: Olenoptera flavilimbata Herrich-Schäffer, [1856] 1850-1858, ibidem 1 (1): 11, 84, wrapper, pl.91, fig.523,

by monotypy. BRAZIL.

O. flavilimbata is a junior subjective synonym of Rolepa delineata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 770.

See also Rolepa Walker, 1855, and Athrula Walker, 1862.

‡OLENOPTERA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 91.

A nomen nudum containing one species ‡flavilimbata Her-

rich-Schäffer, also a nomen nudum.

Olenoptera was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which appeared first in Syst. Bearbeitung Schmett. Eur. 6: 82-133. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) appeared in 1855.

The same Synopsis was also included in Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 2-52; the title-page of this work is dated 1850-1858. Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, stated that pages 1-52 were issued in 1856. Further information is given under Dates of important

Publications in Generic Names Moths World 3.

OLEPA Watson nom. n. for Alope Walker, 1855. ARCT
Type-species: Alone acellifera Walker, 1855. List Specimens

Type-species: Alope ocellifera Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 620, by subsequent designation (for Alope Walker, 1855) by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 247, but cited as Bombyx ricini Fabricius, 1775, a nominal species not originally included in Alope. Kirby, however, placed ricini as the senior synonym of ocellifera, a nominal species originally included in Alope. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included name as the type-species. India: N., and Malabar Coast, Canara, [Bangladesh]: [Sylhet] Silhet, [Sri Lanka] Ceylon.

A. ocellifera is a junior subjective synonym of Bombyx ricini

Fabricius, 1775, Syst. Ent.: 583.

ALOPE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 619.

Type-species: Alope ocellifera Walker, 1855, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 247.

A junior homonym of *Alope* White, 1847 April, *List Crust. Colln Br. Mus.* 1: 75, – Crustacea. The objective replacement name is *Olepa* Watson nom. n.

OLIGAESCHRA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 271.

Type-species: Oligaeschra subvelutina Kiriakoff, 1964, ibidem 14: 271, abb.3, fig.28, pl.3, fig.28, by original designation. China: north Yunnan, Li-Kiang.

OLIGAMATITES Kusnezov, 1928, C. r. Acad. Sci. URSS 1928: 431. FOSSIL CTENUCHIDAE Type-species: Oligamatites martynovi Kusnezov, 1928, ibidem 1928: 431, fig., by monotypy. [U.S.S.R.]: [Kazakhstan], R. Irtysh, Mt Ashutas. Upper Oligocene.

OLIGERIA Turner, 1921, Proc. Linn. Soc. N.S.W. 45: 476 (key), 480.

Type-species: Orgyia hemicalla Lower, 1905, Trans. Proc. R. Soc. S. Aust. 29: 176, by monotypy. Australia: Victoria, Melbourne.

OLIGOCENTRIA Herrich-Schäffer, 1855 December, Syst. Bearbeitung Schmett. Eur. 6: 91; [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 11. NOTO

Type-species: *Notodonta violascens* Herrich-Schäffer, [1855 June], *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.67, fig.385, by monotypy. [BRAZIL] Brasil?: Probably correct, for the series in the BMNH includes specimens from C. America. Venezuela and Brazil.

Oligocentria was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-133. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given in Generic Names Moths World 3 under Dates of some important Publications.

‡*OLIGOCHLAENA*; misspelling. See *Oligoclona* Felder, 1874.

OLIGOCLONA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.94, fig.10, pl.95, fig.8.

Type-species: Oligoclona chordigera Felder, 1874, ibidem 2 (Abt.2): pl.94, fig.10, by PRESENT DESIGNATION. [INDIA]: Darjeeling, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 7.

O. chordigera is a junior subjective synonym of Liparis chrysolopha Kollar, [1844] 1848, in Hügel, Kaschmir und das

Reich der Siek 4: 470.

Oligoclona when established was not placed in a family; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 439, where the name was misspelt ‡Oligochlaena. The type-species was placed in the Notodontidae by Grünberg, 1912, in Seitz, Gross-Schmett. Erde 2: 317; and placed in the Thaumetopoeidae by Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 15.

‡OLIGOCHLAENA; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 439.

An incorrect subsequent spelling of Oligoclona Felder, 1874.

ANSONIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 57.

Type-species: Liparis chrysolopha Kollar, [1844] 1848, in Hügel, Kaschmir und das Reich der Siek 4: 470, by original designation. HIMALAYAS.

A junior homonym of Ansonia Stoliczka, 1870, J. Asiat. Soc. Beng. 39 (2): 152; 1870, Proc. Asiat. Soc. Beng. 1870: 104, – Reptilia. There is no objective replacement name, but L. chrysolopha is a senior subjective synonym of Oligoclona chordigera Felder, 1874, the type-species of Oligoclona Felder, 1874; the latter is therefore available for use as a subjective replacement name.

Ansonia was established in the Notodontidae; it was transferred to the Thaumetopoeidae by Kiriakoff, 1968, *ibidem* 217 (C): 2.

OLINA Walker, 1869. See Aphra Watson nom. n. CTEN

OLISTHERIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 194.

Type-species: *Phalaena zatima* Stoll, 1782, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 182, pl.381, fig.F, by subsequent designation by Forbes, 1955, *Ent. Tijdschr.* 98: 132. Type-locality: stated to be Surinam, but this was evidently an error. *P. zatima* is a form of *Phalaena lutea* Hufnagel, 1766, *Berl. Mag.* 2: 412, occurring in Europe.

Invalid designation of type-species: *Phalaena zatima* Stoll, 1782, was designated by Aurivillius, 1901, *Bih. K. svenska VetenskAkad. Handl.* 27 (4) (7): 15, but the designation was doubtful, being accompanied by an interrogation mark, and is therefore invalid. Aurivillius at the same time placed *Olistheria* in the synonymy of *Phiala* Wallengren, 1860, in the Striphnopterygidae, now Eupterotidae. Forbes, 1955, *ibidem* 98: 132, transferred *Olistheria* to the Arctiidae.

Olistheria Hübner, [1820], is a senior subjective synonym of Spilosoma Curtis, 1825, but has been unused as a valid generic name since it was established in 1820. Under the Code, Articles 23(a-b) and 79(b), an unused senior synonym should not be used to upset a long-established name in current use. This case should therefore be submitted to the Commission for a ruling.

See also Spilarctia Butler, 1875.

OMESTIA Bethune-Baker, 1908, Novit. zool. 15: 177. NOTO Type-species: Omestia bella Bethune-Baker, 1908, ibidem 15: 177, by original designation. New Guinea: [Papua New Guinea], Kebea Range.

Omestia was established in the Eupterotidae; it was placed in the Notodontidae by Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 609.

OMICHLIS Hampson, 1895, Trans. ent. Soc. Lond. 1895: NOTO

Type-species: *Omichlis rufotincta* Hampson, 1895, *ibidem* **1895**: 279, by original designation. [Burma]: E. Pegu.

OMIOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 84, 269.

Type-species: Tospitis fuscipennana Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 431, by original designation. Borneo: Sarawak.

OMOCERINA Kiriakoff, 1970, Lambillionea 70: 26. NOTO Type-species: Omocerina viettei Kiriakoff, 1970, ibidem 70: 28, fig., by original designation. MADAGASCAR: eastern, 42 km. N. of Sambava, Ankala forest.

OMOCHROA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 253.

Type-species: *Omochroa spurca* Rambur, 1866, *ibidem* (2): 254, by monotypy. Spain: Andalusia.

The name ‡*spurca* was used first by Rambur, 1858, *ibidem*: pl.4, fig.3, but was non-binominal and therefore nomenclaturally unavailable until 1866.

HOMOCHROA Spuler, 1906, in Hofmann & Spuler, Schmett. Eur. 2: 132.

An unjustified emendation of Omochroa Rambur, 1866.

A junior homonym of *Homochroa* Hübner, [1825] 1816, Verz. bekannter Schmett.: 358, – Lepid., Pyralidae. The objective replacement name is *Omochroa* Rambur, 1866.

OMOIALA Grote, 1863, Proc. ent. Soc. Philad. 2: 334. PERI Type-species: Omoiala vermiculata Grote, 1863, ibidem 2: 334, pl.6, fig.1, by monotypy. [U.S.A.]: [Colorado], Pike's Peak.

Omoiala was established as the objective replacement name for Lamprosoma Grote, used in the legend to pl.6. Grote stated that he discovered that Lamprosoma was preoccupied too late to correct the plate.

LAMPROSOMA Grote, 1863, Proc. ent. Soc. Philad. 2: pl.6, fig.1.

Type-species: *Omoiala vermiculata* Grote, 1863, by monotypy.

A junior homonym of *Lamprosoma* Kirby, 1819, *Trans. Linn. Soc. Lond.* **12** (2): 445, – Insecta, Coleoptera. The objective replacement name is *Omoiala* Grote, 1863.

CALLALUCIA Grote, 1865, Proc. ent. Soc. Philad. 4: 315. Type-species: Omoiala vermiculata Grote, 1863, by monotypy.

A junior objective synonym of Omoiala Grote, 1863.

ONOPHALERA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 418. NOTO

Type-species: Onophalera duplex Gaede, 1928, ibidem 14: 418, pl.70 row c, by monotypy. [Malawi] Nyassa Land: [Zomba Plateau]. This additional type-locality data is taken from the label on the type-specimen in the BMNH.

The date of publication of the part of Seitz containing page 418 is cited on page 417 as 29.iii.1928 in both the German and the English editions.

ONYCHIPODIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 92, 250.

Type-species: Ghoria nigricostata Butler, 1894, Proc. zool. Soc. Lond. 1894: 583, pl.37, fig.5, by original designation. [Kenya]: Kikuyu escarpment, Kedong.

ONYMAPATA Hulstaert, 1924, Ann. Mag. nat. Hist. (9) 13: 131.

Type-species: Stenocyttara vittata Hulstaert, 1923, ibidem (9) 11: 186, fig., by original designation (for Stenocyttara Hulstaert, 1923). PHILIPPINES: E. Mindanao, Surigao province, Bacuag.

Onymapata was established as the objective replacement name for Stenocyttara Hulstaert, 1923.

STENOCYTTARA Hulstaert, 1923, Ann. Mag. nat. Hist. (9) 11: 186.

Type-species: Stenocyttara vittata Hulstaert, 1923, by original designation.

A junior homonym of Stenocyttara Turner, 1918, Trans. ent. Soc. Lond. 1918: 164, – Lepid., Cossidae. The objective replacement name is Onymapata Hulstaert, 1924.

ONYTHES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 749.

Type-species: *Onythes pallidicosta* Walker, 1855, *ibidem* 3: 749, by monotypy. VENEZUELA.

OPHARUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 728.

Type-species: *Opharus procroides* Walker, 1855, *ibidem* 3: 729, by monotypy. Brazil: Rio [de] Janeiro.

OPHITIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt. 2): pl.97, fig.10.

Type-species: *Ophitis magnaria* Felder, 1874, *ibidem* **2** (Abt.2): pl.97, fig.10, by monotypy. French Guiana: cited by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 7.

OPOBOA Tessmann, 1921, Mitt. zool. Mus. Berl. 10: 215.

Type-species: *Opoboa sexguttata* Tessmann, 1921, *ibidem* 10: 215, by original designation. NIGERIA: S., Opobo.

Opoboa was established in the Pterothysanidae; it was transferred to the Lymantriidae by Collenette, 1932, Novit. zool. 37: 173.

OPSAROA Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 437.

Type-species: *Opsaroa fulvinota* Hampson, 1905, *ibidem* (7) 15: 437, by original designation. South Africa: Cape Province, Triangle.

ORANA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 225.

LYM

Type-species: *Dasychira grammodes* Hering, 1926, *in* Seitz, *Gross-Schmett. Erde* 14: 177, pl.27, row f, by original designation. MADAGASCAR.

ORAURA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 152 (key), 204.

Type-species: *Bireta aurora* Kiriakoff, 1962 [July 31], *ibidem* 98 (10): 204; 1962 October 20, *Bonn. zool. Beitr.* 13: 234, abb.2, fig.18, pl.2, fig.17 (as *Bireta aurora* sp.nov.), by original designation. China: [Chekiang, Mokanshan]. Additional data cited by Kiriakoff, 1962 October 20, *ibidem* 13: 234.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper B. aurora was described as a new species.

As the dates show, the Belgian paper appeared first.

In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

Oraura was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1967, in

Wytsman, Genera Insect. 217 (B): 53.

ORCYNIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 274.

Type-species: Euchromia calcarata Walker, 1854, ibidem 1:

274, by monotypy. Brazil: Santarem.

Orcynia was established as a subgenus of Euchromia Hübner, [1819], in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, as a valid genus by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 154.

‡ORDEMNIAS; misspelling. See Grammia Rambur, 1866.

ORDISHIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 25.

Type-species: *Sphinx rutilus* Stoll, 1782, *in* Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 183, 252, pl.382, fig.B, by original designation. Surinam.

‡OREGYIA; misspelling. See Orgyia Ochsenheimer, 1810.

OREGYIA Meigen, 1830. See Orgyia Ochsenheimer, 1810.

OREINOBIA Wallengren, 1865, K. svenska VetenskAkad. Handl. (N.F.) 5 (4): 34.

Type-species: *Psilura scurrilis* Wallengren, 1860, *Wien. ent. Monatschr.* 4: 163, by monotypy. [SOUTH AFRICA]: Boiesmansrand.

P. scurrilis is a junior subjective synonym of Polymona rufifemur Walker, 1855, the type-species of Polymona Walker, 1855.

See also Polymona Walker, 1855.

OREOCERURA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 216.

Type-species: Cerura dissodectes Kiriakoff, 1958, Ruwenzori

Expdn 1952 1 (3): 47, figs 7, 13, by original designation. UGANDA: Ruwenzori, Nyinabitaba.

OREOCERYX Kiriakoff, 1953, Lambillionea 53: 65.
Type-species: Oreoceryx aenea Kiriakoff, 1953, ibidem 53: 66, pl.1, fig.3, by original designation. [ZAIRE]: Kivu, Loashi.

OREODONTA Kiriakoff, 1967, Tijdschr. Ent. 110: 54. NOTO Type-species: Notodonta gigantea Elwes, 1890, Proc. zool. Soc. Lond. 1890: 399, by original designation. INDIA: Darjeeling.

OREOPOLA Turner, 1940, Proc. R. Soc. Qd 51: 59. LITH Type-species: Oreopola athola Turner, 1940, ibidem 51 RP 59, by monotypy. AUSTRALIA: Tasmania, Mt Wellington.

‡ORGYA; misspelling. See Orgyia Ochsenheimer, 1810.

ORGYA Zetterstedt, [1839]. See Orgyia Ochsenheimer, 1810.

ORGYIA Ochsenheimer, 1810, Schmett. Eur. 3: 208. LYMA Type-species: Phalaena antiqua Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 503, by subsequent designation by Curtis, 1831, Br. Ent. 8: 378. Type-locality: not stated, [Europe].

NOTOLOPHUS Germar, 1812, Diss. sistens Bombycum Species: 35.

Type-species: *Phalaena antiqua* Linnaeus, 1758, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 493.

Notolophus is a junior objective synonym of Orgyia Ochsenheimer, 1810.

‡ORGYJA; Hübner, [1819] 1816, Verz. bekannter Schmett.:

An incorrect subsequent spelling of *Orgyia* Ochsenheimer, 1810.

‡OREGYIA; Billberg, 1820, Enumeratio Insecţ. Mus. G.J. Billberg: 84.

An incorrect subsequent spelling of *Orgyia* Ochsenheimer,

‡ORGYA; Lefebvre, 1826, Mém. Soc. linn. Paris 5: 487.

An incorrect subsequent spelling of *Orgyia* Ochsenheimer, 1810.

‡*Orgya* Lefebvre was cited by Neave, 1940, *Nomencl. zool.* 3: 457, as an available generic name.

OREGYIA Meigen, 1830, Syst. Beschreibung eur. Schmett. 3: 14

An unjustified emendation of *Orgyia* Ochsenheimer, 1810. *Oregyia* is dated from the wrapper of the part and not from the title-page of the volume, 1832.

ORGYA Zetterstedt, [1839] 1840, Insecta Lapponica: 926.
An unjustified emendation of Orgyia Ochsenheimer, 1810.
Orgya Zetterstedt is not a junior homonym of ‡Orgya
Lefebvre, 1826, Mém. Soc. linn. Paris 5: 487, an incorrect subsequent spelling.

MICROPTEROGYNA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 281.

Type-species: *Phalaena antiqua* Linnaeus, 1758, by original designation.

Micropterogyna is dated from the wrapper of livraison 2, and not from the title-page of the work, 1858.

Micropterogyna is a junior objective synonym of Orgyia Ochsenheimer, 1810.

APTEROGYNIS Guenée, [1875] 1867, Statistique scient., Dép. Eure-et-Loire Lépid.: 78.

Type-species: Phalaena antiqua Linnaeus, 1758, by subse-

quent designation by Bryk, 1934, in Strand, Lepid. Cat. 62: 367

Apterogynis is a junior objective synonym of Orgyia Ochsenheimer, 1810.

‡ORGYJA; misspelling. See Orgyia Ochsenheimer, 1810.

ORICIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 476.

Type-species: Oricia truncata Walker, 1854, ibidem 2: 476, by monotypy. Honduras, Guatemala.

ORITHREPTA Collenette, 1939, Ruwenzori Exped. 1934-5 3:

Type-species: Orithrepta edwardsi Collenette, 1939, ibidem 3: 102, fig.1, by original designation. UGANDA: Kigezi, Mt Muhavura.

ORMETICA Clemens, 1861, Proc. Acad. nat. Sci. Philad. 1860: 544.

Type-species: Ormetica sphingiformis Clemens, 1861, ibidem 1860: 545, by monotypy. MEXICO: near Jalapa.

O. sphingiformis is a junior subjective synonym of Chelonia taeniata Guérin-Méneville, [1844], Icon. Règne anim. G. Cuvier (Insectes): 514.

ORNITHOPSYCHE Wallengren, 1863, Wien. ent. Monatschr. 7: 143.

Type-species: Ornithopsyche hypoxantha Wallengren, 1863, ibidem 7: 143, by monotypy. [South Africa]: E. Caffraria orientali.

O. hypoxantha is a junior subjective synonym of Aroa discalis Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 792.

See also Aroa Walker, 1855.

ORODEMNIAS Wallengren, 1885. See Grammia Rambur,

ORONCUS Seitz, 1910, Gross-Schmett. Erde 2: 82. ARCT Type-species: Phragmatobia urania Püngeler, 1904, Societas ent. 19: 121, by subsequent designation by Hampson, 1920, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 2: 342. [U.S.S.R.]: asiatic near Aksu; and Korla.

The date of publication of the part of Seitz containing page 82 was cited on page 81 as 20.iv.1910 in both the German and English editions.

ORPHANOS Hübner, [1825] 1816, Verz. bekannter Schmett.: 306.

Type-species: *Phalaena tripunctaria* Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 523, by PRESENT DESIGNATION. [MALAYSIA OF INDONESIA] Asia.

ORPHANUS Agassiz, 1847, Nomencl. zool. (Index univl.): 264.

An unjustified emendation of Orphanos Hübner, [1825]. Also used by Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 94, 96.

The 'Index universalis' was published as Fascicle 12 of the Nomenclator zoologicus; Orphanos is dated from the wrapper of Fascicle 12, not from its title-page dated 1846.

ORPHANUS Agassiz, 1847. See Orphanos Hübner, [1825].

ORTHOCLOSTERA Butler, 1878, Trans. ent. Soc. Lond. 1878: NOCTUIDAE

Type-species: Orthoclostera peculiaris Butler, 1878, ibidem 1878: 70, by original designation. [BRAZIL]: Amazon region, Rio Negro.

Orthoclostera was established in the Notodontidae; it was placed in the Noctuidae by Hampson, 1912, Cat. Lepid. Phalaenae Br. Mus. 11: 109.

ORTHOLOMIA Felder, 1861, Sher. Akad. Wiss. Wien 43

Type-species: Ortholomia moluccana Felder, 1861, ibidem (Abt.1): 40, by monotypy. [Indonesia]: [Moluccas], Ambon Island.

See also Osica Walker, 1865.

ORTHORHINA Meigen, 1830. See Euchila Billberg, 1820.

ORTHORHINIA Agassiz, 1847. See Euchila Billberg, 1820.

‡ORTHORINA; misspelling. See Euchila Billberg, 1820. NOTO

ORTHORINIA Boisduval, 1828. See Euchila Billberg, 1820.

ORTHOSOMA Felder, 1874. See Rhuda Walker, 1857. NOTO

ORVASCA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 502.

Type-species: Orvasca subnotata Walker, 1865, ibidem 32: 502, by monotypy. [INDIA]: Hindostan.

Orvasca was established in the Limacodidae; it was included in the Liparidae, now Lymantriidae, by Moore, [1883] 1882-3, Lepid. Ceylon 2: 86.

OSICA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 33: 766.

Type-species: Osica glauca Walker, 1865, ibidem 33: 767, by monotypy. [Australia]: [Queensland], Moreton Bay.

O. glauca is a subspecies of Ortholomia moluccana Felder, 1861, Sber. Akad. Wiss. Wien 43 (Abt.1): 40.

Osica was established in the Noctuidae; it was placed in the Notodontidae by Turner, 1903, Proc. Linn. Soc. N.S.W. 28: 71.

See also *Ortholomia* Felder, 1861.

OSMACOLA Watson nom. n. for Calosoma Geyer, 1832.

Type-species: Calosoma flammula Geyer, 1832, in Hübner, Zuträge Samml. exot. Schmett. 4: 14, pl.[110], figs 637, 638, by PRESENT DESIGNATION. JAVA.

CALOSOMA Geyer, 1832, in Hübner, Zuträge Samml. exot. Schmett. 4: 14, 39.

Type-species: Calosoma flammula Geyer, 1832, by PRES-ENT DESIGNATION for Osmacola Watson nom. n.

A junior homonym of Calosoma Weber, 1801, Observationes ent.: 20, – Insecta, Coleoptera, and Calosoma Fabricius, 1801, Syst. Eleutheratorum 1: 211, - Insecta, Coleoptera. objective replacement name is Osmacola Watson nom.n.

OSMOCNEME Draudt, 1917, in Seitz, Gross-Schmett. Erde

Type-species: Osmocneme bracata Draudt, 1917, ibidem 6: 209, pl.29 row d, by original designation. Colembia: eastern, Villavicencio and Rio Negro.

The date of publication of the part of Seitz containing page 209 was cited on that page as 17.ix.1917 in both the German and English editions.

See Otroeda Walker, 1854. ‡OTHROEDA; misspelling.

LYMA

OTPADA Kiriakoff, 1965, Bull. Inst. r. Sci. nat. Belg. 41 (8): 21.

Type-species: Otpada recussa Kiriakoff, 1965, ibidem 41 (8): 23, fig.7, by original designation. ZAIRE: Maniema, Kalima.

OTROEDA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 402.

Type-species: *Phalaena hesperia* Cramer, 1779, *Uitlandsche Kapellen (Papillons exot.)* 3: 99, pl.251, figs A, B, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 428. [West Africa]: Cote de Guinée, près de la fortresse de Delmina.

Otroeda was established in the Lithosiidae, now a subfamily of the Arctiidae.

‡OTHROEDA; Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 138.

An incorrect subsequent spelling of *Otroeda* Walker, 1854. *Otroeda* was incorrectly attributed to Hübner and was misspelled in both the German and the English editions of this work.

OTTACHANA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 153 (key), 179.

Type-species: Bireta sideridis Kiriakoff, 1962 [July 31], ibidem 98 (10): 179; 1962 October 20, Bonn. zool. Beitr. 13: 230, abb.2, figs 12, 13, pl.2, fig.11 (as Bireta sideridis sp.nov.), by original designation. [CHINA]: Tibet, Batang, [Yangtse valley]. Additional data cited by Kiriakoff, 1962 October 20, ibidem 13: 230.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr. to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper B. sideridis was described as a new species. As the dates show, the Belgian paper appeared first.

In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

Ottachana was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 46.

OVENNA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 93.

Type-species: *Eilema subgriseola* Strand, 1912, *Arch. Naturgesch.* 78 (A) 9: 98, by original designation. [CAMEROUN or EQUATORIAL GUINEA] Süd-Kamerun or Spanisch Guinea: Alen, road from Mabungo to Abea.

OVERLAETIA Kiriakoff, 1954. See Tricholoba Hampson, 1910.

OVIPENNIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 88, 410.

Type-species: *Nudaria dudgeoni* Elwes, 1890, *Proc. zool. Soc. Lond.* **1890**: 388, pl.32, fig.10, by original designation. INDIA: near Darjeeling.

OWAMBARCTIA Kiriakoff, 1957, Revue Zool. Bot. afr. 55: 274.

Type-species: Owambarctia owamboensis Kiriakoff, 1957, ibidem 55: 274, fig.6, by original desination. [Namibia] Afrique du sud-ouest: [Ovamboland] région de l'Owambo, Tsumeb.

OXACME Hampson, 1894, Fauna Br. India (Moths) 2: vii, 40, 94.

Type-species: Oxacme dissimilis Hampson, 1894, ibidem 2: 94, fig.55, by original designation. [INDIA]: Sikkim.

OXOIA Kiriakoff, 1967, Tijdschr. Ent. 110: 53. NOTO Type-species: Exaereta smaragdiplena Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 134, by original designation. [BORNEO]: Sarawak.

OXYMETOPA Turner, 1943, Mem. Qd Mus. 12: 105. THAU Type-species: Oxymetopa phaeogramma Turner, 1943, ibidem 12: 105, by monotypy. Australia: Queensland, Talwood.

Oxymetopa was established in the Notodontidae; it was transferred to the Thaumetopoeidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

OZODANIA Dyar, 1899, Psyche, Camb. 8: 359.

Type-species: Ozodania schwarziorum Dyar, 1899, ibidem 8: 359, by original designation. U.S.A.: Arizona, Oracle, Mexico: Vera Cruz State, Rancho Hannover.

PACHASURA Dognin, 1923, Hétérocères nouv. Am. Sud 24:
4. LITH
Type-species: Pachasura stellata Dognin, 1923, ibidem 24:

5, by original designation. Colombia: Pacho.

‡PACHICISPIA; misspelling. See Pachycispia Butler, 1882.

PACHYCERMA Van Duzee, 1897. See Cerma Hübner, 1818.
NOCTUIDAE

PACHYCEROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 85, 270.

Type-species: *Pachycerosia bipuncta* Hampson, 1900, *ibidem* 2: 270, fig.189, by original designation. [INDIA]: [Bay of Bengal], Andamans.

PACHYCERYX Kiriakoff, 1957, Revue Zool. Bot. afr. 55: 283.

Type-species: *Pachyceryx albomaculata* Kiriakoff, 1957, *ibidem* 55: 284, fig.14, by original designation, [ZAIRE]: Tshuapa, Bokuma.

PACHYCISPIA Butler, 1882, Cistula ent. 3: 11.

Type-species: Pachycispia picta Butler, 1882, ibidem 3: 12, by monotypy. MADAGASCAR: Ankafana.

Pachycispia was established in the Liparidae, now Lymantriidae; it was transferred tentatively to the Notodontidae by Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 188; the transfer was confirmed by Collenette, 1955, Trans. R. ent. Soc. Lond. 107: 188.

‡PACHICISPIA; Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 188.

An incorrect subsequent spelling of *Pachycispia* Butler, 1882, in both the German and the English editions.

PACHYDOTA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 4, 92.

Type-species: *Phaegoptera iodea* Herrich-Schäffer, 1855, *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.75, fig.432 (but cited as ‡*Phegoptera*, an incorrect subsequent spelling by Herrich-Schäffer), by original designation (but cited as ‡*Phegoptera*). MEXICO.

P. iodea is dated from the wrapper of the Heft, and not from the title-page of the volume, 1850-1858.

PACHYLAELIA Butler, 1874, Trans. ent. Soc. Lond. 1878: 63.

Type-species: *Pachylaelia translucida* Butler, 1878, *ibidem* **1878**: 63, by monotypy. [BRAZIL]: [Amazonas], Prainha.

Pachylaelia was established in the Liparidae, now Lymantriidae; it is here transferred to the Arctiidae, Arctiinae.

PACHYLISCHIA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 240.

Type-species: *Trichosoma boeticum* Rambur, [1837] 1836, *Annls Soc. ent. Fr.* 5: 587, pl.17A, figs 1-4, by subsequent designation by Kirby, 1870, *Zool. Rec.* 6: 388. Spain: southern.

PACHYPHILONA Butler, 1875, Trans. ent. Soc. Lond. 1875: 325.

Type-species: *Hypsa correcta* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 217, by original designation. [SRI LANKA] Ceylon.

See also Migoplastis Felder, 1868.

PADENIA Moore, [1882] 1882-3, Lepid. Ceylon 2: 58. LITH Type-species; Cyllene transversa Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 544, by monotypy. [SRI LANKA] Ceylon.

PADENODES Rothschild, 1912, Novit. zool. 19: 234. LITH Type-species: Padenodes unifascia Rothschild, 1912, ibidem 19: 235, by subsequent designation by Collins, 1962, Ann. Mag. nat. Hist. (13) 5: 123. New Guinea: [Papua New Guinea], Mambare R., Biagi.

The name *Padenodes* probably originated from Hampson but was used and made nomenclaturally available by Rothschild prior to its proposal by Hampson as the name for a new genus.

PADENODES Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxii, 567.

Type-species: *Padenodes cuprizona* Hampson, 1914, *ibidem* 1: 567, fig.158, by original designation. New Guinea: [Irian Jaya], Snow Mts, upper Setekwa R.

A junior homonym of *Padenodes* Rothschild, 1912, *Novit. zool.* 19: 234, — Arctiidae, Lithosiinae. There is no objective replacement name but the type-species of *Padenodes* Hampson, 1914, is currently considered to be congeneric with the type-species of *Padenodes* Rothschild, 1912. The latter is therefore available for use as a subjective replacement name.

PADENODES Hampson, 1914. See Padenodes Rothschild, 1912.

PAGARA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1678.

Type-species: *Pagara simplex* Walker, 1856, *ibidem* 7: 1679, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 362. Type-locality: not stated, [U.S.A.].

See also Comacla Walker, [1865], and Vanessodes Grote & Robinson, 1870.

‡PAGELLA; misspelling. See Pegella Walker, 1866. LYMA

PAGRASANA Schaus, 1939, Ann. Carnegie Mus. 27: 329.

Type-species: *Pagrasana bermejona* Schaus, 1939, *ibidem* 27: 329, pl.31, fig.14, by original designation. ARGENTINA: Salta, Rio Bermejo.

‡PAIDA; misspelling. See Paidia Hübner, [1819]. LITH

PAIDIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 159.

Type-species: *Phalaena murina* Hübner, 1790, *Beitr. Gesch. Schmett.* 2 (3): 66; (4): 125, pl.(3)2, fig.K, by subsequent designation by Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* 2: 466. GERMANY: Frankfurt.

‡PAIDA; Mabille, 1878, Petites Nouv. Ent. 2: 274.

An incorrect subsequent spelling of Paidia Hübner, [1819].

PAIDINA Staudinger, 1887, in Romanoff, Mém. Lépid. 3: 184.

LITH

Type-species: *Lithosia ramosula* Staudinger, 1887, *ibidem* 3: 184, pl.10, fig.13, by monotypy. [U.S.S.R.]: Raddefka; Vladivostok; Suifun; Ussuri; and Chingan.

Paidina was established as a subgenus of Lithosia Fabricius,

1798.

PAIS Hübner, 1821, Index exot. Lepid.: [4]. AGARISTIDAE Type-species: Phalaena decora Linnaeus, 1764, Mus. Ludovicae Ulricae Reginae: 382, by monotypy. [SOUTH AFRICA]: [Cape of Good Hope] Cap. b. Spei.

Pais has been used as the original generic name for species

now placed in the Arctiidae.

PALAEOCOLEUS Robinson, 1975, Macrolepid. Fiji and Rotuma: 235.

Type-species: *Bocana sypnoides* Butler, 1886, *Trans. ent. Soc. Lond.* 1886: 416, by original designation. Type(s) 9, Fiji: Viti Is.

Palaeocoleus in the Noctuidae, Hypeninae, is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PALAEOMOLIS Hampson, 1909, Ann. Mag. nat. Hist. (8) 4: 362.

Type-species: Palaeomolis purpurascens Hampson, 1909, ibidem (8) 4: 363, by original designation. Peru: Agualani.

PALAEOPSIS Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 15, 86.

Type-species: *Palaeopsis diaphanella* Hampson, 1893, *ibidem* 9:15,87, pl.158, fig.8, by monotypy. [SRI LANKA] Ceylon: Nawala-pittia.

PALAEOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 92, 227.

Type-species: *Lithosia bicosta* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 2: 506, by original designation. [AUSTRALIA]: Sydney.

PALAEOSICCIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 92, 505.

Type-species: *Palaeosiccia punctata* Hampson, 1900, *ibidem* 2: 505, fig. 361, by original designation. SIERRA LEONE.

PALAEOSTAUROPUS Okagaki & Nakamura, 1953, Butterfl. Moths 4: 17.

Type-species: Stauropus obliterata Wileman & South, 1917, Entomologist 50: 29, by original designation. [JAPAN]: Oiwake.

PALAEOTYPE Hampson, 1918, Novit. zool. 25: 109. LITH Type-species: Trischalis submarginata Rothschild, 1915, Rep. Br. orn. Un. Expedition N. Guinea 2 (15): 50, by original designation. New Guinea: [Irian Jaya], [Oetakwa] Utakwa River.

PALAEOZANA Dyar, 1914, Proc. U.S. natn. Mus. 47: 167.

Type-species: *Palaeozana mida* Dyar, 1914, *ibidem* 47: 167, by original designation. Panama: La Chorrera.

PALAEXERA Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 691, 699.

Type-species: *Palaexera phyllodes* Meyrick, 1886, *ibidem* (2) 1: 699, by monotypy. Australia: New South Wales, Sydney.

‡PALASARA Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 421.

An incorrect subsequent spelling of *Palaexera* Meyrick, 1886. ‡*Palasara* was cited by Neave, 1940, *Nomencl. zool.* 3: 535, as 'pro *Palirisa* Moore 1884' in the Eupterotidae. However,

from the content of Pagenstecher's work it is almost certain that he was referring to Palaexera Meyrick, 1886.

‡PALASARA; misspelling. See Palaexera Meyrick, 1886. LITH

PALASEA Wallengren, 1863, Wien. ent. Monatschr. 7: 142.

Type-species: Palasea albimacula Wallengren, 1863, ibidem 7: 143, by monotypy. [South Africa]: E. Caffraria orientali.

PALESSA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 153 (key), 190.

Type-species: Pydna alboflavida Bryk, 1950, Ark. Zool. 42 (A) 19: 18, pl.4, fig.5, by original designation. BURMA: north, Kambaiti.

Palessa was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 100.

PALILEMA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 77.

Type-species: Eilema uelleburgensis Strand, 1912, Arch. Naturgesch. 78 (A) 9: 98, by original designation. [CAMEROUN] or EQUATORIAL GUINEA] Süd-Kamerun or Spanisch Guinea: Uelleburg, road to Uelle.

Palilema was established as a subgenus of Archilema Birket-

Smith, 1965.

PALLENE Walker, 1854. See Prinasura Hampson, 1903.

PALPARCTIA Spuler, 1906. See Acymba Rambur, 1866.

PALPIDIA Dyar, 1898, Jl N.Y. ent. Soc. 6: 33. NOCTUIDAE Type-species: Palpidia pallidior Dyar, 1898, ibidem 6: 34, by monotypy. Holotype 9, U.S.A.: Florida, Cocoanut Grove, (USNM, Washington).

Palpidia was established in the Lithosiidae, now a subfamily of the Arctiidae; in a personal communication Prof. J.G. Franclemont has stated that Palpidia is a senior subjective synonym of Echinocampa Franclemont, 1949.

Palpidia now in the Noctuidae, Ophiderinae was not included

in Nye, 1975, Generic Names Moths World 1.

PALUSTRA Bar, 1873, Annls Soc. ent. Fr. (5) 3: 300. ARCT Type-species: Palustra laboulbeni Bar, 1873, ibidem (5) 3: 301, pl.8(2), figs 1-8, by monotypy. French Guiana.

‡PAMCALOMA; misspelling. See Pamcoloma Schaus, 1905.

PAMCOLOMA Schaus, 1905, Proc. U.S. natn. Mus. 29: 280.

Type-species: Pamcoloma marita Schaus, 1905, ibidem 29: 281, by original designation. FRENCH GUIANA: Maroni River, St Laurent.

‡PAMCALOMA; Schaus, 1928, Proc. U.S. natn. Mus. 73 (19): 55.

An incorrect subsequent spelling of Pamcoloma Schaus, 1905.

PAMPA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 238. ZYGAENIDAE

Type-species: Euchromia mystica Walker, 1854, ibidem 1: 239, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 112. HONDURAS.

Pampa has been used as the original generic name for species

now placed in the Ctenuchidae.

Pampa was established in the 'Zygaenides' as a subgenus of Euchromia Hübner, [1819], now in the Ctenuchidae.

PANACHRANTA Turner, 1922, Proc. R. Soc. Vict. 35: 30.

Type-species: Panachranta lirioleuca Turner, 1922, ibidem 35: 30, by monotypy. AUSTRALIA: N. Queensland; Cairns and [Kuranda] Karanda.

PANASSA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 607.

Type-species: Panassa cingalesa Walker, 1865, ibidem 32: 607, by monotypy. [SRI LANKA] Ceylon.

P. cingalesa is a junior subjective synonym of Aemene taprobanis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 542.

See also Aemene Walker, 1854, and Autoceras Felder, 1874.

PANAXIA Tams, 1939. See Callimorpha Latreille, 1809.

PANGORA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 42.

Type-species: Pangora distorta Moore, 1879, ibidem (1): 42, pl.2, fig.14, by monotypy. [INDIA]: [Mussooree] Masuri, Simla, and N.W. Himalayas.

PANIA Dalman, 1823, Analecta Ent.: 92. NOTO Type-species: Bombyx phantoma Dalman, 1823, ibidem: 92, by monotypy. Sweden: northern, Lapland.

Pania was established as a subgenus of Bombyx Linnaeus,

‡PANORGYIA; misspelling. See Parorgyia Packard, 1864.

PANTANA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 819.

Type-species: Pantana dispar Walker, 1855, ibidem 4: 820, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 464, but cited as 'P. Visum, Hübn.' a nominal species not originally included in Pantana. Kirby, however, placed visum as the senior synonym of dispar, a nominal species originally included in Pantana. Under the Code, Article 69(a)(iv), this constitutes a type-species designation. EAST INDIES.

PANTANOPSIS Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 387.

Type-species: Pantanopsis diehli Kiriakoff, 1974, ibidem 17: 388, text-fig.10, pl.2, fig.3, by original designation. SUMATRA: northern, Deli, Dolok Merangir.

PANTELECLITA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 418.

Type-species: Panteleclita viridipicta Kiriakoff, 1974, ibidem 17: 419, text-fig.30, pl.5, fig.4, by original designation. SUMATRA: northern, Dairi.

PANTHERINUS Nakamura, 1973, Ent. Rev. Japan 25: 58.

Type-species: Pseudofentonia bipunctata Okano, 1960, Rep. Gakugei Fac. Iwate Univ. 16: 16, text-figs 15, 16, pl.7, fig.7, by original designation. [TAIWAN]: central, Jen-ai-hsiang, Nantow-hsien.

PAPUAROA Collenette, 1955, Nova Guinea (N.S.) 6: 55.

Type-species: Papuaroa miccyla Collenette, 1955, ibidem 6: 55, pl.1, fig.17, by original designation. New Guinea: [Irian Jaya], Snow Mts, Top Camp.

PARABASIS Bethune-Baker, 1904, Novit. zool. 11: 428. NOTO

Type-species: *Parabasis pratti* Bethune-Baker, 1904, *ibidem* 11: 429, pl.4, fig.26, by original designation. New Guinea: [Papua New Guinea], Dinawa.

Parabasis was established in the Noctuidae; it was placed in the Notodontidae by Clench, 1956, Lepid. News 10: 15.

PARABITECTA Hering, 1926, in Draeseke, Dt. ent. Z. Iris 40: 50.

Type-species: *Parabitecta flava* Hering, 1926, *in* Draeseke, *ibidem* 40: 50, by monotypy. CHINA: Szechwan.

PARABLAVIA Schaus, 1899, Jl N.Y. ent. Soc. 7: 215. LITH Type-species: Parablavia xanthura Schaus, 1899, ibidem 7: 215, by monotypy. [BRAZIL]: Parana, Castro.

The name *Parablavia* probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson as the name for a new genus.

PARABLAVIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 81, 125.

Type-species: Lithosia sadima Schaus, 1896, Jl N.Y. ent. Soc. 4: 151, by original designation. [BRAZIL]: Parana, Castro.

A junior homonym of *Parablavia* Schaus, 1899. There is no objective replacement name but the type-species of *Parablavia* Hampson, 1900, is currently considered to be congeneric with the type-species of *Parablavia* Schaus, 1899. The latter is therefore available for use as a subjective replacement name.

PARABLAVIA Hampson, 1900. See Parablavia Schaus, 1899.

PARACERYX Bethune-Baker, 1904, Novit. zool. 11: 383.

Type-species: *Paraceryx aroa* Bethune-Baker, 1904, *ibidem* 11: 383, by original designation. New Guinea: [Papua New Guinea], upper Aroa River.

PARACHADISRA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 636.

Type-species: *Damata varians* Bethune-Baker, 1908, *Novit. zool.* 15: 176, by monotypy. New Guinea: [Papua New Guinea], Kebea Range.

The date of publication of the part of Seitz containing page 636 was cited on page 633 as 21.ii.1930 in the German edition.; and as 17.v.1930 in the English edition.

PARACHELONIA Aurivillius, 1900, Ent. Tidskr. 20: 234, 239.

Type-species: Chelonia rubriceps Mabille, 1879, Bull. Soc. zool. Fr. 3: 88, by original designation. MADAGASCAR.

PARACINCIA Field, 1950, Proc. U.S. natn. Mus. 100: 321.

Type-species: *Paracincia butleri* Field, 1950, *ibidem* 100: 322, pl.9, fig.14, by original designation. JAMAICA: St Andrew Parish, Newcastle.

PARACLEAPA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 57.

Type-species: Cleapa psecas Druce, 1901, Ann. Mag. nat. Hist. (7) 7: 78, by original designation. [Tanzania]: [Tanganyika], Dar-es-Salaam.

PARACLES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 717.

Type-species: *Paracles contraria* Walker, 1855, *ibidem* 3: 717, by monotypy. [BRAZIL]: Para.

PARADESTOLMIA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 227.

Type-species: *Notodonta nigrolinea* Lucas, 1895, *Trans. nat. Hist. Soc. Qd* 1: 107, by original designation. AUSTRALIA: [Queensland], Brisbane.

PARADIASTEMA Aurivillius, 1901, Ent. Tidskr. 22: 120.

Type-species: *Paradiastema nigrocincta* Aurivillius, 1901, *ibidem* **22**: 121, fig.22, by monotypy. [ZAIRE] Congo: Kinshasa.

PARADINIA Draudt, 1915, in Seitz, Gross-Schmett. Erde 6: 110.

Type-species: *Paradinia chrysogastrides* Draudt, 1915, *ibidem* 6: 110, pl.18 row c, by original designation. Colombia.

The date of publication of the part of Seitz containing page 110 was cited on page 105 as 16.iv.1915 in both the German and English editions.

PARADIOPTIS Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 521.

Type-species: *Dioptis otanes* Druce, 1893, *Proc. zool. Soc. Lond.* **1893**: 294, by monotypy (of *Eudioptis* Prout, 1918). ECUADOR: Rio Napo.

Paradioptis was established as the objective replacement

name for Eudioptis Prout, 1918.

The date of publication of the part of Seitz containing page 521 was cited on that page as 24.x.1925 in both the German and the English editions.

EUDIOPTIS Prout, 1918, Novit. zool. 25: 397 (key), 416. Type-species: Dioptis otanes Druce, 1893, by monotypy.

A junior homonym of *Eudioptis* Hübner, 1823, *Zuträge* Samml. exot. Schmett. 2: 24, – Lepid., Pyralidae. The objective replacement name is Paradioptis Hering, 1925.

PARADOHERTYA Bethune-Baker, 1904, Novit. zool. 11: 421.

Type-species: *Paradohertya trifascia* Bethune-Baker, 1904, *ibidem* 11: 421, pl.4, fig.36, by original designation. New GUINEA: [Papua New Guinea], Aroa River.

PARADOXOSIA Hampson, 1918, Novit. zool. 25: 105. LITH Type-species: Paradoxosia rufipex Hampson, 1918, ibidem 25: 106, by original designation. UGANDA: S.E. Buddu, Tero Forest.

PARADRALLIA Bethune-Baker, 1908, Ann. Mag. nat. Hist. (8) 2: 255.

Type-species: *Paradrallia rhodesi* Bethune-Baker, 1908, *ibidem* (8) 2: 256, by original designation. [ZAMBIA] N.E. Rhodesia: Fort Jameson.

Paradrallia was established in the Eupterotidae; it was placed in the Notodontidae by Hulstaert, 1924, Rev. Zool. Bot. afr. 12: 398; and subsequently placed in the Thaumetopoeidae by Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 396.

PARADRYMONIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 122. NOTO

Type-species: *Notodonta vittata* Staudinger, 1892, *Dt. ent.* Z. *Iris* 4: 262, by original designation. [TURKEY]: Hadjin [approx. 36E 38N].

PARAETHRIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xvi, 28, 353.

Type-species: Gnophaela triseriata Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.49, fig.261, by original designation. BRAZIL.

GNOPHAELA Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.49, fig.261.

Type-species: *Gnophaela triseriata* Herrich-Schäffer, [1855], by monotypy.

A junior homonym of Gnophaela Walker, 1854. The objective replacement name is Paraethria Hampson, 1898.

Gnophaela Herrich-Schäffer probably originated from Boisduval who attributed it to himself in Boisduval, 1870, Considérations Lépid. Guatemala: 87.

PARAFENTONIA Roepke, 1944, Natuurh. Maandbl. 33: 19.

Type-species: Parafentonia sundana Roepke, 1944, ibidem 33: 20, fig. 20, by monotypy. [JAVA]: Mt Gedeh, Goalpara.

PARAFENTONIA Kiriakoff, 1964. See Hemifentonia Kiriakoff, 1967.

PARAGLUPHISIA Djakonov, 1927, Ezheg. zool. Muz. 27: 219.

Type-species: Paragluphisia oxiana Djakonov, 1927, ibidem 27: 220, pl.15, figs 1, 3, 7, 9, pl.16, fig.12, by original designation. U.S.S.R.: [Uzbekistan], S. Bukhara, Termez.

PARAGYLLA Dognin, 1899, Annls Soc. ent. Belg. 43: 132.

Type-species: Paragylla endophaea Dognin, 1899, ibidem 43:

132, by monotypy. [ECUADOR]: Loja.

Paragylla probably originated from Hampson but was used and made nomenclaturally available by Dognin prior to its proposal by Hampson as the name for a new genus.

PARAGYLLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 91, 213.

Type-species: Areva amoureli Dognin, 1890, Naturaliste 12: 128, by original designation. [ECUADOR]: near Loja.

A junior homonym of Paragylla Dognin, 1899. There is no objective replacement name but the type-species of Paragylla Hampson, 1900, is currently considered to be congeneric with the type-species of Paragylla Dognin, 1899. The latter is therefore available for use as a subjective replacement name.

PARAGYLLA Hampson, 1900. See Paragylla Dognin, 1899.

PARAKANCHIA Bethune-Baker, 1908, Novit. zool. 15: 189.

Type-species: Parakanchia grisea Bethune-Baker, 1908, ibidem 15: 189, by original designation. New Guinea: [Papua New Guinea], [Papua], Babooni.

PARALACYDES Aurivillius, 1899, Ent. Tidskr. 20: 234, 239.

Type-species: Phalaena vocula Stoll, 1790, in Cramer, Uitlandsche Kapellen (Papillons exot.) (Aanhangsel): 143, 184, pl.31, fig.5, by subsequent designation by Hampson, 1900, Ann. S. Afr. Mus. 2: 54 (for Maenas Hübner, [1819]). [SOUTH AFRICA]: Cape of Good Hope.

Paralacydes was established as the objective replacement

name for Maenas Hübner, [1819].

P. vocula is dated from the wrapper of the part and not from the title-page of the volume, 1791.

MAENAS Hübner, [1819] 1816, Verz. bekannter Schmett.: 167.

Type-species: Phalaena vocula Stoll, 1790, by subsequent designation by Hampson, 1900, Ann. S. Afr. Mus. 2: 54.

A junior homonym of Maenas Walbaum, 1792, in Artedi, P. Artedi renovati ... Ichthyologiae (Edn 2) 3: 585, - Pisces. The objective replacement name is Paralacydes Aurivillius,

EUTAENIA Wallengren, 1875, Ofvers. K. Vetensk Akad. Förh. Stockh. 32 (1): 102.

Type-species: Eutaenia scapulosa Wallengren, 1875, ibidem 32 (1): 102, by monotypy. South Africa: Transvaal.

A junior homonym of Eutaenia Thomson, 1857, Archs ent.

Paris 1: 184, - Insecta, Coleoptera. There is no objective replacement name but the type-species of Eutaenia Wallengren, 1875, is currently considered to be congeneric with the typespecies of Paralacydes Aurivillius, 1899. The latter is therefore available for use as a subjective replacement name.

E. scapulosa is a junior subjective synonym of Lacydes

arborifera Butler, 1875, Cistula ent. 2: 26.

CALIGULA Aurivillius, 1879, Ofvers. K. VetenskAkad. Förh. Stockh. 36: 56.

Type-species: Caligula wallengreni Aurivillius, 1879, ibidem **36**: 56, by monotypy.

A junior homonym of Caligula Moore, 1862, Trans. ent. Soc. Lond. (3) 1: 321, - Lepid., Saturniidae. There is no objective replacement name but the type-species of Caligula Aurivillius, 1879, is currently considered to be congeneric with the typespecies of Paralacydes Aurivillius, 1899. The latter is therefore available for use as a subjective replacement name.

C. wallengreni is a junior subjective synonym of Lacydes

arborifera Butler, 1875, Cistula ent. 2: 26.

PARALAETHIA Hampson, 1907, Ann. Mag. nat. Hist. (7) 19: 224.

Type-species: Ceryx subformicina Bethune-Baker, 1904, Novit. zool. 11: 382, by original designation. New Guinea: [Papua New Guinea], upper Aroa River.

PARALERODES Kiriakoff, 1969, Faune Madagascar 29: 23.

Type-species: Lerodes albosparsatus Kenrick, [1914] 1913, Trans. ent. Soc. Lond. 1913 (4): 591, pl.32, fig.17, by original designation. MADAGASCAR.

L. albosparsatus was established in the Lasiocampidae; it was transferred to the Notodontidae by Kiriakoff, 1969, ibidem

PARALITHOSIA Daniel, 1954, Bonn. zool. Beitr. 5: 131.

Type-species: Paralithosia hoenei Daniel, 1954, ibidem 5: 131, text-fig.93, pl.3, fig.75 (but cited as \$\pm\$hönei, an incorrect original spelling), by original designation. CHINA: N. Yunnan, Li-kiang.

PARAMAENAS Grünberg, 1911, Stettin. ent. Ztg 72: 381.

Type-species: Paramaenas strigosus Grünberg, 1911, ibidem 72: 381, by monotypy. [Namibia] South West Africa: [Windhoek] Windhuk.

PARAMARBLA Collenette, 1937, Ann. Mag. nat. Hist. (10) 20: 605.

Type-species: Paramarbla teroensis Collenette, 1937, ibidem (10) **20**: 606, pl.13, fig.7, by original designation. UGANDA: S.E. Buddu, Tero Forest.

PARAMECES Turner, 1936, Proc. R. Soc. Qd 47: 29. NOTO Type-species: Parameces glauca Turner, 1936, ibidem 47: 29, by monotypy. Australia: Queensland, Kuranda.

‡PARAMELESE Zikan, 1968, in Zikan & Zikan, Pesquisa Agropec. bras. 3: 92.

An unavailable name under the Code, Article 13(a) and (b), having neither description, indication, nor type-species. The single included nominal species, ‡Paramelese itatiayensis Zikan, 1968, ibidem 3: 92 is also a nomen nudum.

PARAMELISA Aurivillius, 1905, Ark. Zool. 3 (1): 13. THYRET

Type-species: Paramelisa lophura Aurivillius, 1905, ibidem 3 (1): 13, pl.1, fig.2, by monotypy. [ZAIRE] Kongo: Mukimbungu [5°6'S 14°4'E].

Paramelisa was established in the Arctiidae; it was placed

in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 7.

PARAMEVANIA Draudt, 1915, in Seitz, Gross-Schmett. Erde 6: 161.

Type-species: *Paramevania inconspicua* Draudt, 1915, *ibidem* 6: 161, pl.23, row k, by monotypy. Type-locality: not stated, [AMERICA]: [tropical].

The date of publication of the part of Seitz containing page 161 was cited on that page as 3.v.1915 in both German and English editions.

PARAMSACTA Hulstaert, 1923, Ann. Mag. nat. Hist. (9) 11: 187.

Type-species: *Paramsacta pura* Hulstaert, 1923, *ibidem* (9) 11: 187, by original designation. [New Guinea]: [Irian Jaya], Okaba.

PARAMULONA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 84, 386.

Type-species: *Mieza albulata* Herrich-Schäffer, 1866, *CorrespBl. zool.-min. Ver. Regensburg* **20**: 120, by original designation. CUBA.

PARAMYA Druce, 1898 May. See Metamya Travassos, 1946.

CTEN

PARAMYA Hampson, 1898 [December]. See *Metamya* Travassos, 1946.

PARAMYOPSYCHE Debauche, 1942, Explor. Parc natn. Albert 41: 7. CTEN

Type-species: *Paramyopsyche wittei* Debauche, 1942, *ibidem* 41: 7, text-figs a, b, pl.1, figs 1-3, by original designation. [ZAIRE] Congo Belge: Mikeno volcano, Rweru.

PARANERICE Kiriakoff, 1964, Bonn. zool. Beitr. 14: 280.

Type-species: *Paranerice hoenei* Kiriakoff, 1964, *ibidem* 14: 280, abb.6, fig.60, pl.5, fig.41, by original designation. China: Shansi, Mien-shan.

PARANERITA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xv, 3, 439.

Type-species: *Evius polyxenus* Druce, 1883, *Proc. zool. Soc. Lond.* **1883**: 383, pl.40, fig.6, by original designation. ECUADOR: Sarayacu.

PARAONA Moore, 1878, Proc. zool. Soc. Lond. 1878: 8.

Type-species: *Crambomorpha splendens* Butler, 1877, *Trans. ent. Soc. Lond.* **1877**: 357, by monotypy. [INDIA]: Bombay.

PARAONAGYLLA Berio, 1939, Missione Biol. Paese dei Borana 3: 12.

Type-species: *Paraonagylla zavattarii* Berio, 1939, *ibidem* 3: 12, text-figs 1-4, by original designation. [ETHIOPIA]: Neghelli.

PARAPALOSIA Dyar, 1905, Proc. U.S. natn. Mus. 29: 174.

Type-species: *Parapalosia cinderella* Schaus, 1905, *ibidem* **29**: 197, by original designation (in anticipation of Schaus's description of *cinderella*). FRENCH GUIANA: St Jean.

Parapalosia was available but was without an available included species between its date of publication 1905 October 3rd and that of its type-species 1905 October 11th by Schaus.

PARAPELOSIA Bethune-Baker, 1908, Novit. zool. 15: 192.

Type-species: *Parapelosia grisescens* Bethune-Baker, 1908, *ibidem* 15: 192, by original designation. New Guinea: [Papua New Guinea], Owgarra.

PARAPHLEBS Aurivillius, 1921, Ann. S. Afr. Mus. 18: 240.

Type-species: *Paraphlebs singularis* Aurivillius, 1921, *ibidem* 18: 241, by monotypy. South Africa: Transvaal, Barberton and Potgietersrust.

‡PARAPHLEPS; Gaede, 1934, in Strand, Lepid. Cat. 59: 202. An incorrect subsequent spelling of Paraphlebs Aurivillius, 1921.

‡PARAPHLEPS; misspelling. See Paraphlebs Aurivillius, 1921.

PARAPHRYGIA Hampson, 1920, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 2: xx, 493.

Type-species: *Phryganopteryx rectangulata* Kenrick, 1914, *Trans. ent. Soc. Lond.* **1913**: 600, fig., by original designation. MADAGASCAR.

PARAPIRGA Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 552.

Type-species: *Parapirga neurabrunnea* Bethune-Baker, 1911, *ibidem* (8) 7: 552, by original designation. [ANGOLA]: [Vila Salazar] N'Dalla Tando.

PARAPISA Kiriakoff, 1952, Revue fr. Ent. 19: 173. THYRET Type-species: Apisa bourgognei Kiriakoff, 1952, ibidem 19: 173, by original designation. IVORY COAST: Bingerville.

Parapisa was established as a subgenus of Apisa Walker, 1855.

PARAPLASTIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 507.

Type-species: *Migoplastis hampsoni* Swinhoe, 1889, *Proc. zool. Soc. Lond.* **1889**: 402, pl.43, fig.2, by original designation. [INDIA]: [Madras], Nilgiri Hills.

PARAPLESIS Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 95, 97.

Type-species: *Paraplesis collaris* Herrich-Schäffer, 1855, *ibidem* 6: 97, by monotypy. Brazil.

A nomen dubium, but associated by its author with the nominal genus *Gnophria* Stephens, 1829.

Paraplesis was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

PARAPLOCIA Aurivillius, 1909. See *Calpoparia* Watson nom. n.

PARAPREPIA Dyar, 1913, Insecutor Inscit. menstr. 1: 75.

Type-species: *Talara ruficollis* Schaus, 1896, *Jl N.Y. ent.* Soc. 4: 149, by original designation. BRAZIL: São Paulo. See also *Parillice* Dyar, 1913.

PARAPROCTIS Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 543.

Type-species: *Paraproctis osiris* Bethune-Baker, 1911, *ibidem* (8) 7: 543, by original designation. [NIGERIA]: Lokoja District.

PARARCTIA Sotavalta, 1965, Suom. hyönt. Aikak. 31: 173.

Type-species: *Bombyx lapponica* Thunberg, 1791, *Diss. ent. sistens Insecta Suecica* 2: 40, pl., fig.7, by original designation. Sweden: [Lappland] Lapponia.

PARARETHONA Janse, 1920, Ann. Transv. Mus. 7: 173.

Type-species: *Desmeocraera hierax* Distant, 1897, *Ann. Mag. nat. Hist.* (6) **20**: 204, by original designation. South Africa: Transvaal, Pretoria and Lydenburg district.

PARARHENEA Kiriakoff, 1960, Revue Zool. Bot. afr. 61: 55.

Type-species: *Pararhenea viridescens* Kiriakoff, 1960, *ibidem* 61: 56, fig.8, by original designation. [ZAIRE]: Uele, Paulis.

PARARHENEADES Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 314.

Type-species: *Pararheneades plumosa* Kiriakoff, 1965, *ibidem* 101: 314, fig.4, by monotypy. UGANDA: Ankole, Kalinzu forest.

PARAROTHIA Kiriakoff, 1974, in Kiriakoff & Viette, Faune Madagascar 39: 6 (key), 72.

AGARISTIDAE

Type-species: *Rothia vieui* Viette, 1966, *Lambillionea* 65: 88, fig.2, by original designation. Holotype & MADAGASCAR: E., environs de Perinet, foret d'Analamazaotra, (MNHN, Paris).

Pararothia in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PARASCAPTIA Bethune-Baker, 1908, Novit. zool. 15: 195.

LITH

Type-species: *Parascaptia biplagata* Bethune-Baker, 1908, *ibidem* 15: 195, by original designation. New Guinea: [Papua New Guinea], Babooni.

PARASCEPSIS Dognin, 1923, Hétérocères nouv. Am. Sud 23: 2. CTEN

Type-species: *Parascepsis solox* Dognin, 1923, *ibidem* 23: 3, by original designation. [Brazil]: [Amazonas], São Paulo de Olivença.

PARASCOLIA Watson nom. n. for Pseudoscolia Hampson, 1914.

Type-species: *Pseudoscolia tenebrifera* Hampson, 1914, *Cat. Lepid. Phalaenae Br. Mus.* (Suppl.) 1: 450, fig., by original designation (for *Pseudoscolia* Hampson, 1914). New Guinea: [Papua New Guinea], Angabunga River.

PSEUDOSCOLIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xix, 450.

Type-species: *Pseudoscolia tenebrifera* Hampson, 1914, by original designation.

A junior homonym of *Pseudoscolia* Radoszkovsky, 1876, *Horae Soc. ent. ross.* 12: 103, – Insecta, Hymenoptera. The objective replacement name is *Parascolia* Watson nom. n.

PARASCOTIA Hübner, [1825] 1816, Verz. bekannter Schmett.: 314.

Type-species: Geometra carbonaria [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 108, by subsequent designation by Warren, 1913, in Seitz, Gross-Schmett. Erde 3: 399.

Further information in Nye, 1975, Generic Names Moths World 1: 370.

KARA Matsumura, 1925, J. Coll. Agric. Hokkaido imp. Univ. 15: 160.

Type-species: *Kara sachalinensis* Matsumura, 1925, *ibidem* 15: 160, pl.8, fig.13, by original designation. Syntypes 16, 19, [U.S.S.R.]: [S. Sakhalin] South-Saghalien, Shimizu, (EIHU, Sapporo).

A junior homonym of Kara Strebel, 1910, Abh. Geb. Naturw.

Hamburg 19 (3): 16, — Mollusca. There is no objective replacement name, but *K. sachalinensis* is a junior subjective synonym of *Geometra carbonaria* [Denis & Schiffermüller], 1775, the type-species of *Parascotia* Hübner, [1825]. The latter is therefore available for use as a subjective replacement name.

Kara was established in the Geometridae; it was transferred to the Noctuidae, Ophiderinae by Fletcher, 1979, in Nye, Generic Names Moths World 3.

Kara in the Noctuidae, Ophiderinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PARASCRANCIA Kiriakoff, 1969, Bull. Inst. fond. Afr. noire
(A) 30 (4): 1446.

NOTO

Type-species: *Parascrancia chadisroides* Kiriakoff, 1969, *ibidem* (A) **30** (4): 1447, fig.8, by original designation. IVORY COAST: Adiopodoumé.

The date of publication of part 4, January 1969, is taken from the foot of page 1762 of that part, not from its title-page which is dated October 1968.

PARASEMIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 181.

Type-species: *Phalaena plantaginis* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 501, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 250. Type-local-ITY: not stated, [EUROPE].

See also Eupsychoma Grote, 1865.

NEMEOPHILA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 55, 72.

Type-species: *Phalaena plantaginis* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 501, by monotypy.

A junior objective synonym of *Parasemia* Hübner, [1820]. Nemeophila was proposed again by Stephens, 1829, Syst. Cat. Br. Insects (2): 55.

CHIONOPHILA Guenée, 1865, Annls Soc. ent. Fr. (4) 4: 686. Type-species: Phalaena plantaginis Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 501, by PRESENT DESIGNATION.

A junior objective synonym of Parasemia Hübner, [1820].

PARASHADA Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 433.

Type-species: *Parashada truncata* Hampson, 1905, *ibidem* (7) **15**: 434, by monotypy. [INDONESIA]: [Moluccas], Sula Is, [Mangole] Mongoli.

PARASICCIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 90, 407.

Type-species: *Aemene maculifascia* Moore, 1878, *Proc. zool. Soc. Lond.* **1878**: 33, fig., by original designation. [INDIA]: Darjeeling.

PARASINGA Kiriakoff, 1967, Tijdschr. Ent. 110: 51. NOTO Type-species: Somera lichenina Butler, 1880, Ann. Mag. nat. Hist. (5) 6: 67, by original designation. BORNEO.

PARASOLOE Kiriakoff, 1954, Bull. Inst. r. Sci. nat. Belg. 30 (29): 8.

Type-species: *Parasoloe tetrasticta* Kiriakoff, 1954, *ibidem* **30** (29): 8, fig.11, by original designation. [ZAIRE] Congo Belge: Kafakumba.

Parasoloe was proposed as ‡Parasoloe. Under the Code, Article 27, the diaeresis must be deleted.

PARASTATIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xix, 28, 468.

Type-species: *Marissa parnassia* Möschler, 1878, *Verh. zool.-bot. Ges. Wien* **27** (Abh.): 636, pl.8, fig.5, by original designation. SURINAM: Paramaribo.

PARASTAURA Gaede, 1928, in Seitz, Gross-Schmett, Erde 14: 429.

Type-species: Parastaura divisa Gaede, 1928, ibidem 14: 429,

pl.71 row c, by monotypy. [ZIMBABWE-]RHODESIA.

The date of publication of the part of Seitz containing page 429 was cited on page 425 as 30.iii.1928 in both the German and the English editions.

PARATALARA Dyar, 1905, Proc. U.S. natn. Mus. 29: 175.

Type-species: Paratalara inversa Schaus, 1905, ibidem 29: 203, by original designation (in anticipation of Schaus's description of inversa). FRENCH GUIANA: St Jean.

Paratalara was available but was without an available included species between its date of publication 1905 October 3rd and that of its type-species 1905 October 11th by Schaus.

PARATHEMERASTIS Bethune-Baker, 1916, Ann. Mag. nat. Hist. (8) 17: 383.

Type-species: Parathemerastis melanistis Bethune-Baker, 1916, ibidem (8) 17: 383, by subsequent designation by Collins, 1962, ibidem (13) 5: 5. NEW GUINEA: [Papua New Guinea], Mt Kebea.

P. melanistis was established as a variety of Osica turneri Bethune-Baker, 1904, in the combination Parathemerastis turneri var. melanistis Bethune-Baker. Var. melanistis was subsequently raised to specific rank by Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 638.

PARATHISANOTIA Kiriakoff, 1976. See Cyphocampa Harris, 1869. AGARISTIDAE

PARATHYRIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 158.

Type-species: Phalaena cedonulli Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 108, pl.346, figs A, B, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 852. SURINAM.

P. cedonulli is dated from the wrapper of the part and not

from the title-page of the volume, 1782.

THYRARCTIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 3, 9.

Type-species: Phalaena cedonulli Stoll, 1781, by original designation.

A junior objective synonym of Parathyris Hübner, [1819].

‡PARATROTENOTUS; misspelling. See Paratrotonotus Aurivillius, 1925.

PARATROTONOTUS Aurivillius, 1925, Ergebn. zweiten dt. zent.-Afr. Exped. 1910-1911 (Zool.) 1 (18): 1267.

Type-species: Ctenogyna ogovensis Holland, 1893, Ent. News 4: 342, pl.15, fig.12, by monotypy. [GABON]: [Ogooué R. valley], Kangwé.

When established *Paratrotonotus* contained a second nominal species, doubtfully included; under the Code, Articles 67(h) and 68(c), a doubtfully included species is not eligible for selection as type-species.

‡PARATROTENOTUS; Hering, 1930, Revue Zool. Bot. afr. 17: 405.

An incorrect subsequent spelling of *Paratrotonotus* Aurivillius, 1925.

PARATYPE Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.104, fig.11.

Type-species: Paratype laeticolor Felder, 1874, ibidem 2 (Abt.2): pl.104, fig.11, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 8.

PARAUCHMIS Holland, 1897, in Smith, Through unknown African Countries: 414.

Type-species: Parauchmis smithii Holland, 1897, ibidem: 414, fig.2, by monotypy. [ETHIOPIA]: Dombalok [38°27'E 5°0'N].

P. smithii is a junior subjective synonym of Eutimia marpissa Wallengren, 1858, Ofvers. K. VetenskAkad. Förh. Stockh. 15:

Parauchmis was established in the Noctuidae; it was placed in the Notodontidae by Nye, 1975, Generic Names Moths World 1: 371.

See also Eutimia Wallengren, 1858.

PARAXENA Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 544.

Type-species: Paraxena esquamata Bethune-Baker, 1911. ibidem (8) 7: 545, by original designation. [ANGOLA]: [Vila Salazar] N'Dalla Tando.

PARAZANA Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 555.

Type-species: Parazana radiata Bethune-Baker, 1911, ibidem (8) 7: 555, by original designation. [NIGERIA]: Lokoja district. P. radiata is a junior subjective synonym of Anticyra rufovittata Aurivillius, 1901, Ent. Tidskr. 22: 122.

PARELICTIS Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 691, 709.

Type-species: Parelictis saleuta Meyrick, 1886, ibidem (2) 1: 709, by monotypy. AUSTRALIA: New South Wales, Sydney; Victoria.

PAREMONIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xx, 86, 557.

Type-species: Hemonia luteicincta Holland, 1893, Psyche, Camb. 6: 412, by original designation. [CAMEROUN]: [Ogooué] Ogové River valley.

PAREUCHAETES Grote, 1865, Proc. ent. Soc. Philad. 5: 245.

Type-species: Pareuchaetes cadaverosa Grote, 1865, ibidem 5: 245, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 237. CUBA.

P. cadaverosa is a junior subjective synonym of Halysidota insulata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus 3: 734 (but proposed as \$\frac{1}{2}Halesidota\$, an incorrect subsequent spelling).

PAREUGOA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 88, 405.

Type-species: Eugoa multistrigata Hampson, 1898, J. Bombay nat. Hist. Soc. 11: 439, by original designation. [INDIA]: [Assam], Khasia Hills.

PAREUGOA Daniel, 1951. See Apogurea Watson nom. n.

PAREVIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 3, 38.

Type-species: Evius sisenna Druce, 1899, Ann. Mag. nat. Hist. (7) 3: 466, by original designation. BRAZIL: [Espirito Santo] Espiritu Santo.

PAREXILISIA Toulgoët, 1958, Mém. Inst. scient. Madagascar (E) 9: 192.

Type-species: Philenora bipunctoides Toulgoët, 1953, ibidem (E) 4: 358, text-fig.11, pl.1, fig.11, by original designation. MADAGASCAR: north east, near Maroantsetra, Ambodivoangy.

PARILLICE Dyar, 1913, Insecutor Inscit. menstr. 1: 75. LITH

Type-species: Parillice deceptans Dyar, 1913, ibidem 1: 75, by original designation. BRAZIL: Parana, Castro.

P. deceptans is a junior subjective synonym of Talara rust-collis Schaus, 1896, Jl N.Y. ent. Soc. 4: 149.

See also Paraprepia Dyar, 1913.

PAROCNERIA Dyar, 1897, Can. Ent. 29: 13, 14 (key), 15.

Type-species: Phalaena detrita Esper, 1785, Die Schmett. 3:

229, pl.44, fig.6, by original designation. EUROPE.

P. detrita is dated from the wrapper of the part in which it was established, and not from the title-page of the volume, 1782.

‡DANIELA Hartig, 1963, in Hartig & Prola, Mitt. münch. ent. Ges. 53: 65.

Not nomenclaturally available. ‡Daniela was proposed for *Phalaena detrita* Esper, 1785, and three other species, but no type-species was designated. Under the *Code*, Article 13(b), a genus-group name published after 1930 must, in order to be available, be accompanied by the fixation of a type-species. In addition to this, the name is preoccupied by *Daniela* Koch, 1891, *Mitt. zool. Stn Neapel* 9: 669, —Coelenterata.

PARORGYIA Packard, 1864, Proc. ent. Soc. Philad. 3: 332.

Type-species: *Phalaena achatina* Smith, 1797, in Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia*: 153, pl.77, (a junior primary homonym, see below), by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 471. [U.S.A.]: Georgia.

The type-species is a junior primary homonym of *Phalaena* (Noctua) achatina Sulzer, 1776, Abgekürzte Gesch. Insecten 1: 160; 2: pl.22, fig.4, – Noctuidae, Catocalinae. There is no objective replacement name but *Phalaena achatina* Smith is a subjective synonym of Dasychira meridionalis (Barnes & McDunnough, 1913) ssp. memorata Ferguson, 1978, Moths Am. N. of Mexico 22 (2): 39, text-fig.8d, pl.3, figs 30-37, pl.4, figs 3, 8, 16, 17; the latter is therefore available for use as a subjective replacement name.

‡PANORGYIA; Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 227.

An incorrect subsequent spelling of Parorgyia Packard, 1864.

PAROSICA Kiriakoff, 1967, Tijdschr. Ent. 110: 37. NOTO Type-species: Parosica nomo Kiriakoff, 1967, ibidem 110: 38, fig.1, by original designation. New Guinea: [Irian Jaya], Mt Nomo, south of Mt Bougainville.

PARVICINCIA Field, 1950, Proc. U.S. natn. Mus. 100: 323.

Type-species: *Parvicincia belli* Field, 1950, *ibidem* **100**: 324, pl.9, figs 16, 17, by original designation. JAMAICA: Trelawny Parish, near Jackson Town, Baron Hill.

PASSINEURA Butler, 1876, J. Linn. Soc. (Zool.) 12: 412.

Type-species: Antichloris phemonoides Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 639, by original designation (but cited as Pampa fusiformis Walker, see below). SURINAM.

The original designation by Butler of *Pampa fusiformis* Walker, 1856, as type-species of *Passineura*, was an error as *P. fusiformis* Walker sensu Butler, 1876, was a misidentification for *Antichloris phemonoides* Möschler, 1878, as shown by Möschler, 1878, *ibidem* 27: 640 and Field, 1975, *Smithson. Contr. Zool.* 198: 8. This is a case of a misidentified type-species which should be placed before the Commission under the *Code*, Article 70(a).

PASTEOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xi, 84, 217.

Type-species: *Pasteosia irrorata* Hampson, 1900, *ibidem* 2: 218, fig.135, by original designation. [SOUTH AFRICA]: Cape Province, Annshaw.

PATRELIURA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xviii, 30, 413.

Type-species: *Sphinx capys* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 88, pl.56, figs E, F, by original designation. Surinam.

S. capys is dated from the wrapper of the part and not from the title-page of the volume, 1779.

PAULIANOSIA Toulgoët, 1958, Mém. Inst. scient. Madagascar (E) 9: 198.

Type-species: *Paulianosia clathrata* Toulgoët, 1958, *ibidem* 9: 198, text-fig.21, pl.2, fig.14, by original designation. MADAGASCAR: E., Ronomafana.

PAULISANA Kiriakoff, 1960, Revue Zool. Bot. afr. 61: 54.

Type-species: *Paulisana rufina* Kiriakoff, 1960, *ibidem* 61: 54, fig.6, by original designation. [ZAIRE]: Uele, Paulis.

PAULUMA Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 260 (key), 292.

Type-species: *Pauluma nubila* Schaus, 1901, *ibidem* **1901**: 292, pl.11, fig.9, by original designation. [BRAZIL]: Parana, Castro.

PAUROPHLEPS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xix, 91, 531.

Type-species: *Paurophleps minuta* Hampson, 1900, *ibidem* 2: 531, fig.385, by original designation. [SOUTH AFRICA]: Cape [Province] Colony, Transkei.

PECTINIDIA Holloway, 1977, Series Ent. 13: 91.

Type-species: *Pectinidia silvicola* Holloway, 1977, *ibidem* 13: 93, text-figs 37, 38, pl.22, fig.2, by original designation. Holotype &, Norfolk Island (South Pacific Ocean): Mt Bates, (BMNH).

Pectinidia in the Noctuidae, Ophiderinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PECTINOPHORA Janse, 1920. See Eujansea Fletcher nom.

NOTO

PEGELLA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1922.

Type-species: *Pegella curvifera* Walker, 1866, *ibidem* 35: 1922, by monotypy. PHILIPPINES.

‡PAGELLA; Kozhanchikov, 1950, Fauna SSSR (N.S.) 42 (Lepid. 12): 351.

An incorrect subsequent spelling of Pegella Walker, 1866.

PELLEWIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 135.

Type-species: *Fentonia cantiana* Schaus, 1928, *Proc. U.S. natn. Mus.* **73** (19): 79, by original designation. CHINA: Szechwan, Suifu.

PELLUCENS Bethune-Baker, 1910, Ann. Mag. nat. Hist. (8) 6: 447.

Type-species: *Pellucens lactea* Bethune-Baker, 1910, *ibidem* (8) 6: 447, by original designation. New Guinea: [Irian Jaya] Dutch N.G., N. coast.

PELOBROCHIS Lucas, 1892, Proc. Linn. Soc. N.S.W. (2) 7: 250 (but attributed to Meyrick, an incorrect authorship).

Type-species: Scaeodora rava Lucas, 1890, Proc. Linn. Soc. N.S.W. (2) 4: 1079, by monotypy. AUSTRALIA: Brisbane.

S. rava is a junior subjective synonym of Thumatha fuscescens Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1901.

See also Thumatha Walker, 1866.

PELOCHYTA Hübner, [1819] 1816, Verz. bekannter Schmett.: 171, 172.

Type-species: Sphinx arontes Stoll, 1782, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 184, 248, pl.382, fig.E, by subsequent designation by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 103. Surinam.

The type-species designations by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 194, 195, cannot be accepted under the Code, Article 69(a)(iii), as he designated more than one valid species and it is not clear which he considered to be the type-species.

See also Sontia Walker, 1856.

PELOROSES Collenette, 1955, Naturaliste malgache 7: 177.

Type-species: *Numenes praestans* Saalmüller, 1884, *Lepid. Madagascar* (Abt.1): 187, by original designation. MADAGASCAR: N.W., Nossi-Bé Island.

PELOSIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 165.

Type-species: *Phalaena muscerda* Hufnagel, 1766, *Berlin. Mag.* 3: 400, by subsequent designation by Moore, 1878, *Proc. zool. Soc. Lond.* 1878: 27. GERMANY: Berlin region.

SAMERA Wallengren, 1863, Wien. ent. Monatschr. 7: 146, 147.

Type-species: *Phalaena muscerda* Hufnagel, 1766, by original designation.

A junior objective synonym of *Pelosia* Hübner, [1819].

‡PETOSIA; Swinhoe, 1893, Ann. Mag. nat. Hist. (6) 12: 216. An incorrect subsequent spelling of Pelosia Hübner, [1819]. Cited by Neave, 1940, Nomencl. zool. 3: 679, as 'pro Petasia Stephens, 1829', – Lepid., Noctuidae.

PELTOSTAURA Kiriakoff, 1967, Biologia gabon. 3: 337.

Type-species: *Peltostaura pelias* Kiriakoff, 1967, *ibidem* 3: 337, fig.24, by original designation. GABON: Makokou.

‡*PEMPHEGOSTOLA*; misspelling. See *Pemphigostola* Strand, 1909. AGARISTIDAE

PEMPHIGOSTOLA Strand, 1909, Dt. ent. Z. 1909: 663.

Type-species: Pemphigostola synemonistis Strand, 1909, ibidem 1909: 665, by monotypy. Holotype & MADAGASCAR, (MNHU, Berlin).

Pemphigostola was established in the Castniidae, Pemphigostolinae; it was transferred to the Apoprogenidae, now Apoprogonidae, by Hampson 1918, Novit. zool. 25: 377; and transferred to the Agaristidae by Jordan, 1921, Novit. zool. 28: 73.

Pemphigostola in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡PEMPHEGOSTOLA; Hampson, 1918, Novit. zool. 25: 377. An incorrect subsequent spelling of Pemphigostola Strand, 1909.

‡PENARA; misspelling. See Penora Walker, 1855. LYMA

PENDRIA Swinhoe, 1906, Ann. Mag. nat. Hist. (7) 17: 540.

Type-species: Redoa rinaria Moore, [1860] 1858-9, in Hors-

field & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 336, by original designation. JAVA.

PENORA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 821.

Type-species: *Penora remota* Walker, 1855, *ibidem* 4: 821, by subsequent designation by Schaus, 1927, *in* Seitz, *Gross-Schmett. Erde* 6: 560. SOUTH AMERICA: West coast.

‡PENARA; Bryk, 1931, in Strand, Lepid. Cat. 45: 9.
An incorrect subsequent spelling of Penora Walker, 1855.

PENTELIA Kiriakoff, 1976. See Sergiusia Nye nom. n. AGARISTIDAE

PENTHOPHERA Germar, 1812, Diss. sistens Bombycum Species: 33.

Type-species: *Phalaena morio* Linnaeus, 1767, *Syst. Nat.* (Edn 12) 1 (2): 828, by subsequent designation by Curtis, 1828, *Br. Ent.* 5: 213. [Austria]: [Vienna] Wiennae.

‡HIPOGYMNA Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) (Opinion 97): 19. Also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 (Opinion 278): 140. Only included species: Phalaena morio Linnaeus, 1767.

LIPARIS Ochsenheimer, 1810, Schmett. Eur. 3: 186.

Type-species: *Phalaena morio* Linnaeus, 1767, by subsequent designation by Hampson, 1918, *Novit. zool.* **25**: 384.

Incorrect type-species designations: (1) Phalaena dispar Linnaeus, 1758, cited by Boisduval, 1836, Hist. nat. Insectes (Spec. Gén. Lépid.) 1: 134. Boisduval in his lengthy 154 page Introduction, reviewed earlier classifications and designated up to three different type-species for each generic name. In his 'Exposé de notre méthode' from page 155-690, no type-species designations were made for the genera he himself used. Under the Code, Article 69(a)(iii), the type-designation of an author is eligible for consideration if he states that it is the type '... and if it clear that he himself accepts it as the type-species'. Boisduval's type-designations although clearly stated do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not in the past been accepted by Hemming or other authors. Acceptance of Boisduval's designation in this case would make no difference to the generic concept.

(2) Phalaena monacha Linnaeus, 1758, cited by Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 424, after Hampson's valid designation.

A junior homonym of *Liparis* Scopoli, 1777, *Introductio Hist. nat.*: 453, – Pisces. The objective replacement name is *Penthophera* Germar, 1812.

LIPARUS Harris, 1841, Rep. Insects Mass. injurious to Vegn 4: 260.

An unjustified emendation of *Liparis* Ochsenheimer, 1810. A junior homonym of *Liparus* Olivier, 1807, *Entomologie*, ou Hist. nat. Insectes 5: 284, — Coleoptera. The objective replacement name is *Penthophera* Germar, 1812.

ERASTA Gistl, 1848, Naturg. Thierreichs: ix.

Erasta was established as an objective replacement name for Liparis Ochsenheimer, 1810.

‡PENTHOPHORA; Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 789.

An incorrect subsequent spelling of *Penthophera* Germar, 1812.

‡PENTOPHORA; Heylaerts, 1884, Annls Soc. ent. Belg. 28 (Bull.); xli.

An incorrect subsequent spelling of *Penthophera* Germar,

‡PENTOPHERA; Neave, 1940, Nomencl. zool. 3: 654.

An incorrect subsequent spelling of *Penthophera* Germar, 1812.

‡Pentophera was a misspelling by Neave in citing ‡Pentophora Heylaerts, 1884.

‡PENTHOPHORA; misspelling. See Penthophera Germar, 1812.

PENTOBESA Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 258 (key), 269.

Type-species: *Edema xylinoides* Walker, 1866, *List Specimens lepid. Insects Colln Br. Mus.* **35**: 1931, by original designation. [COLOMBIA]: Santa Martha.

‡PENTOPHERA; misspelling. See Penthophera Germar, 1812.

‡PENTOPHORA; misspelling. See Penthophera Germar, 1812.

PERATODONTA Aurivillius, 1904, Ark. Zool. 2 (4): 6.

Type-species: *Peratodonta brunnea* Aurivillius, 1904, *ibidem* 2 (4): 6, text-figs 14, 15, by monotypy. CAMEROUN.

PERCOTE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 492.

Type-species: *Percote signatura* Walker, 1854, *ibidem* 2: 493, by monotypy. WEST INDIES.

PERICALLIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 182. ARCT

Type-species: *Phalaena matronula* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 509, by monotypy. GERMANY.

PLERETES Lederer, 1853, Verh. zool.-bot. Ver. Wien 2 (Abh.): 77.

Type-species: *Phalaena matronula* Linnaeus, 1758, by monotypy.

A junior objective synonym of *Pericallia* Hübner, [1820].

‡PLEREDES; Hofmann, 1887, Schmett. Eur.: 41.

An incorrect subsequent spelling of Pleretes Lederer, 1853.

PERICOPIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 175.

Type-species: *Phalaena zerbina* Stoll, 1790, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* (Aanhangsel): 173, 184, pl.40, fig.3, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 184. SURINAM.

P. zerbina is dated from the wrapper of the part and not

from the title-page of the volume, 1791.

P. zerbina is a junior subjective synonym of Phalaena eurocilia Cramer, 1777, Uitlandsche Kapellen (Papillons exot.) 2: 126, 148, pl.178, fig.C. P. eurocilia is dated from the wrapper and not from the title-page of the volume, 1779.

PERIDEA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 32.

Type-species: Bombyx serrata Thunberg, 1792, Diss. ent. sistens Insecta Suecica (3): 47, by monotypy. Sweden.

B. serrata is a junior subjective synonym of Phalaena anceps

Goeze, 1781, Ent. Beytr. 3 (3): 207.

Sherborn, 1902, *Index Anim.* (1758-1800): 893 cited an earlier reference to *B. serrata*, namely Thunberg, 1788, *D. D. Mus. nat. Upsal.* (6): 70; in this publication *B. serrata* was a nomen nudum.

Peridea was again proposed by Stephens, 1829 [June], Nom. Br. Insects: 39; and 1829 [July], Syst. Cat. Br. Insects (2): 45.

‡PERIDIA;, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 5, 60.

An incorrect subsequent spelling of Peridea Stephens, 1828.

MESODONTA Matsumura, 1920, Zool. Mag. Tokyo 32: 145.

Type-species: Notodonta monetaria Oberthür, 1879, Diagnoses Espèces nouv. Lépid. ile Askold: 12, by original designation. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

A junior homonym of *Mesodonta* Baly, 1865, *Entomologist's mon. Mag.* 2: 99, – Insecta, Coleoptera. There is no objective replacement name, but *N. monetaria* is congeneric with *Bombyx serrata* Thunberg, 1792, the type-species of *Peridea* Stephens, 1828; the latter is thus available for use as a subjective replacement name.

Mesodonta was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 790.

‡PERIDIA; misspelling. See Peridea Stephens, 1828.

PERIDOPSIS Nakamura, 1976. See Pulia Kiriakoff, 1974.

PERIDROME Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 444.

Type-species: *Hypsa orbicularis* Walker, 1854, *ibidem* 2: 445, by monotypy. INDIA:N, [BANGLADESH]: [Sylhet] Silhet, 'EAST INDIES'.

Peridrome was established as a subgenus of Hypsa Hübner, [1819].

See also Aganopis Herrich-Schäffer, [1856].

PERIERGOS Kiriakoff, 1959, Ark. Zool. (2) 12: 321. NOTO Type-species: Periergos obsoleta Kiriakoff, 1959, ibidem (2)
12: 322, text-fig.17, pl.1, fig.6, by original designation. [Burma]: Kambaiti.

PERINA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 966.

Type-species: *Perina basalis* Walker, 1855, *ibidem* 4: 966, by monotypy. [Nepal] Nepaul, India: North, Hong Kong.

Perina was established in the Psychidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 467.

P. basalis is a junior subjective synonym of Bombyx nuda Fabricius, 1787, Mantissa Insect. 2: 119.

PERINETIA Collenette, 1936, Novit. zool. 40: 165. LYMA Type-species: Perinetia leucocloea Collenette, 1936, ibidem 40: 166, pl.12, fig.8, by original designation. MADAGASCAR: E., 149km E. of Tananarivo, Périnet.

PERIPHALERA Kiriakoff, 1959, Ark. Zool. (2) 12: 314.

Type-species: *Phalera albicauda* Bryk, 1949, *ibidem* 42 (A) 19: 8, pl.1, fig.2, by original designation. Burma: Kambaiti.

PERNAMBIS Schaus, 1920, Proc. U.S. natn. Mus. 57: 124.

PYROMORPHIDAE

Type-species: *Pernambis intervenata* Schaus, 1920, *ibidem* 57: 124, by original designation. BRAZIL: Pernambuco.

Pernambis was established in the 'Lymantriadae'; it was placed in the Cossidae by Dyar, 1937, in Seitz, Gross-Schmett. Erde 6: 1276; and is here transferred to the Cossoidea, Pyromorphidae.

PEROARA Schaus, 1905, Proc. U.S. natn. Mus. 29: 245.

Type-species: *Peroara sylvestris* Schaus, 1905, *ibidem* 29: 245, by monotypy. French Guiana: Maroni River, St Jean.

PEROMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1022.

Type-species: *Peroma anomala* Walker, 1855, *ibideni* 5: 1022, by monotypy. [Brazil]: Rio de Janeiro.

PERONETIS Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 242.

Type-species: *Peronetis xenodora* Meyrick, 1886, *ibidem* (2) 1: 242, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 300. New Guinea: [Irian Jaya], Fly River.

PETALIA Walker, 1854. See Asota Hübner, [1819]. AGAN

PETALOPLEURA Meyrick, 1889, Trans. ent. Soc. Lond. 1889: 458.

Type-species: *Petalopleura phaeocephala* Meyrick, 1889, *ibidem* **1889**: 459, by monotypy. New Guinea.

P. phaeocephala is a junior subjective synonym of Lithosia impervia Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 230.

‡PETOSIA; misspelling. See Pelosia Hübner, [1819]. LITH

PETRINIA Walker, 1869, in Chapman, Proc. nat. Hist. Soc. Glasgow 1: 347.

NOCTUIDAE

Type-species: Petrinia lignosa Walker, 1869, ibidem 1: 348,

by monotypy. [ZAIRE] Congo.

Petrinia was established in the Noctuidae; it was included doubtfully in the Notodontidae by Gaede, 1934, in Strand, Lepid. Cat. 59: 319, and restored to the Noctuidae by Gaede, 1935, in Seitz, Gross-Schmett. Erde 15: 187.

PEZAPTERA Butler, 1876, J. Linn. Soc. (Zool.) 12: 404.

Type-species: *Eunomia sordida* Walker, 1856, *List Specimens lepid. Insects Colln Br. Mus.* 7: 1617, by original designation. [Brazil]: Santarem.

PEZOPTERA Zerny, 1912, in Wagner, Lepid. Cat. 7: 101. An unjustified emendation of Pezaptera Butler, 1876.

PEZOPTERA Zerny, 1912. See Pezaptera Butler, 1876.

PHACUSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 150. ZYGAENIDAE

Type-species: Glaucopis tenebrosa Walker, 1854, ibidem 1:

150, by monotypy. INDIA: N.

Phacusa Walker, 1854, ibidem 1: 150, was established as a subgenus of Glaucopis Fabricius, 1807.

PHACUSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 175. See Leucotmemis Butler, 1876. CTEN

The same name, *Phacusa*, was established by Walker, 1854, *ibidem* 1 pages 150 and 175, for two different subgenera of *Glaucopis* Fabricius, 1807.

PHACUSOSIA Hampson, 1911, Ann. Mag. nat. Hist. (8) 8: 404.

Type-species: *Phacusosia xanthosoma* Hampson, 1911, *ibidem* (8) 8: 405, by original designation. New Guinea: [Papua New Guinea], Babooni.

PHAEGOPTERA Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pls 14, 15, figs 58-68.

Type-species: *Phaegoptera histrionica* Herrich-Schäffer, [1853] 1850-1858, *ibidem* 1 (1): wrapper, pl.15, fig.68 (but attributed to Boisduval, an incorrect authorship), by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 212. Brazil.

Phaegoptera Herrich-Schäffer, [1853], and the similar name Phoegoptera Boisduval, 1869, are discussed by Watson, 1973, Smithson. Contr. Zool. 128: 2.

‡PHEGOPTERA; Herrich-Schäffer, [1855] 1850-1858,

Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pls 52-53, figs 283-294; wrapper, pl.75, fig.432.

An incorrect subsequent spelling of *Phaegoptera* Herrich-Schäffer, [1853], repeated by Herrich-Schäffer, [1856], *ibidem* 1 (1): wrapper, pl.92, fig.528; and [1856], *ibidem* 1: 16.

PHAEGORISTA Boisduval, [1836], Hist. nat. Insectes (Spec. gén. Lépid.) 1 (Atlas): explanation to pl.14.

AGAN

Type-species: *Phaegorista agaristoides* Boisduval, [1836], *ibidem* 1 (Atlas): explanation to pl.14, by monotypy. SENEGAL.

The spelling *Phaegorista* was adopted by Neave, 1940, *Nomencl. zool.* 3: 686, as first reviser, from a multiple original spelling.

‡PHOEGORISTA Boisduval, [1836], Hist. nat. Insectes (Spec. gén. Lépid.) 1 (Atlas): legend to pl.14.

An incorrect (of a multiple) original spelling of *Phaegorista* Boisduval, [1836].

PHAENARETE Druce, 1896, Ann. Mag. nat. Hist. (6) 18: 31.

Type-species: *Phaenarete diana* Druce, 1896, *ibidem* (6) **18**: 31, by original designation. PANAMA: Chiriqui.

PHAEO Zerny, 1912. See Phaio Neumoegen, 1894. CTEN

PHAEOCHLAENA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 18, 32.

Type-species: *Phaeochlaena tendinosa* Hübner, 1818, *ibidem* 1: 18, pl.[16], figs 89, 90, by subsequent designation by Butler, 1878, *Trans. ent. Soc. Lond.* 1878: 62. BRAZIL.

P. tendinosa is a junior subjective synonym of Zygaena gyon Fabricius, 1787, Mantissa Insect. 2: 105.

PHLAEOCHLOENA; Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 156.

An incorrect subsequent spelling of *Phaeochlaena* Hübner, 1818.

‡PHLAEOCHLAENA; Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1876.

An incorrect subsequent spelling of *Phaeochlaena* Hübner, 1818.

‡*PHAEOCHLOENA*; Druce, 1885, *Proc. zool. Soc. Lond.* **1885**: 536, legend pl.33, fig.16.

An incorrect subsequent spelling of *Phaeochlaena* Hübner, 1818.

‡PHAEOCLAENA; Dognin, 1900, Annls Soc. ent. Belg. 44: 438.

An incorrect subsequent spelling of *Phaeochlaena* Hübner, 1818.

‡PHAEOCLENA; Neave, 1940, Nomencl. zool. 3: 688.

An incorrect subsequent spelling of *Phaeochlaena* Hübner, 1818. Neave attributed this misspelling to Druce, 1885, *Proc. zool. Soc. Lond.* **1885**: 535, but on this page *Phaeochlaena* was correctly spelt. On page 536, however, Druce misspelt the name as ‡*Phaeochloena*.

‡PHAEOCHLOENA; misspelling. See Phaeochlaena Hübner, 1818.

‡PHAEOCLAENA; misspelling. See Phaeochlaena Hübner, 1818.

‡PHAEOCLENA; misspelling. See Phaeochlaena Hübner, 1818.

PHAEOMOLIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 3, 20.

Type-species: Neritos obscurata Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: xii, 51, pl.17, fig.6, by original designation. [BRAZIL]: [Amazonas], Prainha.

PHAEOPHLEBOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 82, 109.

Type-species: Setina furcifera Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 520, by original designation. [AUSTRALIA]: [Tasmania] Van Dieman's Land.

PHAEOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 81, 128.

Type-species: *Lithosia lutea* Druce, 1885, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 1: 131, by original designation. GUATEMALA: San Geronimo.

PHAEOSIA Spuler, 1903. See Pheosia Hübner, [1819].

‡PHAEOSINA; misspelling. See Pheosina Gaede, 1916.

PHAEOSPHECIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xii, 26, 153.

Type-species: Myrmecopis opaca Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1620, by original designation. [Brazil]: Para, Amazon valley.

PHAIO Neumoegen, 1894, Can. Ent. 26: 334. CTEN Type-species: Phaio longipennis Neumoegen, 1894, ibidem 26: 335, by monotypy. CUBA.

PHAEO Zerny, 1912, in Wagner, Lepid. Cat. 7: 96.
An unjustified emendation of Phaio Neumoegen, 1894.

‡*PHALAE*; misspelling. See *Phaloe* Guérin-Méneville, [1838].

PHALAENA Linnaeus, 1758. See Naenia Stephens, 1827.
NOCTUIDAE

‡PHALAESIA; misspelling. See Phaloesia Walker, 1854.

PHALANNA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 218.

Type-species: *Sphinx polymena* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 494, by subsequent designation by Hampson, 1898, *Cat. Lepid. Phalaenae Br. Mus.* 1: 293. ASIA: [tropical southern].

Phalanna was established as a subgenus of Euchromia Hübner, [1819].

PHALCIDON Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 301.

Type-species: *Josia integra* Walker, 1854, *ibidem* 2: 301, by subsequent designation by Prout, 1918, *Novit. zool.* 25: 420. HONDURAS.

J. integra is a subspecies of Bombyx megaera Fabricius, 1787, Mantissa Insect. 2: 109.

‡PHALCIDONA; Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 135.

An incorrect subsequent spelling of Phalcidon Walker, 1854.

‡PHALCIDONA; misspelling. See Phalcidon Walker, 1854.

PHALERA Hübner, [1819] 1816, Verz. bekannter Schmett.:

Type-species: *Phalaena bucephala* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 508, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 576. Type-locality: not stated, [Europe].

ACROSEMA Meigen, 1830, Syst. Beschreibung eur. Schmett. 3: 24.

Type-species: *Phalaena bucephala* Linnaeus, 1758, by subsequent designation by Grote, 1897, *Illte Wschr. Ent.* **2**: 389.

A junior objective synonym of *Phalera* Hübner, [1819].

HAMMATOPHORA Westwood, 1843, in Humphreys & Westwood, Br. Moths 1: 63.

Type-species: *Phalaena bucephala* Linnaeus, 1758, by monotypy.

A junior objective synonym of Phalera Hübner, [1819].

PHALERINA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 259.

Type-species: *Phalerina terminalis* Kiriakoff, 1964, *ibidem* 14: 260, abb.2, fig.15, pl.1, fig.14, by original designation. China: Fukien, Kuatun.

PHALERODONTA Staudinger, 1892, in Romanoff, Mém. Lépid. 6: 367.

Type-species: *Notodonta bombycina* Oberthür, 1880, *Etudes d'Entomologie* 5: 63, pl.6, fig.3, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

PHALEROMIMUS Bryk, 1950, Ark. Zool. 42 (A) 19: 9.

Type-species: *Phaleromimus albocalceolata* Bryk, 1950, *ibidem* **42** (A) 19: 9, pl.1, fig.3, by original designation. BURMA: Kambaiti.

PHALOE Guérin-Méneville, [1838] 1830, in Duperrey, Voyage autour du Monde sur la Corvette la Coquille (Zool.) 2 (2) 1: 283.

Type-species: *Pericopis cruenta* Hübner, 1823, *Zuträge Samml. exot. Schmett.* **2**: 24, pl.[58], figs 329, 330, by monotypy. [Brazil]: Bahia.

‡PHALAE; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1653.

An incorrect subsequent spelling of *Phaloe* Guérin-Méneville, [1838].

PHALOESIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 359.

Type-species: *Phaloesia saucia* Walker, 1854, *ibidem* 2: 359, by monotypy. NICARAGUA.

‡PHALAESIA; Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1653.

An incorrect subsequent spelling of *Phaloesia* Walker, 1854.

PHANEROPSEUSTIS Turner, 1899, Trans. Proc. R. Soc. S. Aust. 23: 12 (but attributed to Meyrick (Meyr.(MSS.)), an incorrect authorship).

Type-species: *Nudaria albida* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 273, by monotypy. [Australia]: [Queensland], Moreton Bay.

PHANOPTIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.104, fig.10. DIOP

Type-species: *Phanoptis cyanomelas* Felder, 1874, *ibidem* 2 (Abt.2): pl.104, fig.10, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 8.

Prout, 1918, *Novit. zool.* **25**: 414, proposed *Phanoptis* Felder as a new genus because he believed Felder's publication of the generic name to be invalid, as it was not accompanied by a description; according to the present *Code*, Article 13, this requirement applies only to names published after 1930.

PHAOS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 627.

Type-species: *Phaos interfixa* Walker, 1855, *ibidem* 3: 627, by monotypy. [Australia]: [Tasmania] Van Dieman's Land.

PHARA Walker, 1854. See Aphra Watson nom. n. CTEN

PHASTIA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 258.

Type-species: *Phastia basalis* Walker, 1862, *ibidem* (3) 1: 258, by monotypy. [Brazil]: Rio de Janeiro.

‡PHASTRA; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 622.

An incorrect subsequent spelling of Phastia Walker, 1862.

‡PHASTRA; misspelling. See Phastia Walker, 1862. NOTO

PHAUDA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 256.

Type-species: *Euchromia flammans* Walker, 1854, *ibidem* 1: 257, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 111. INDIA: N.

Phauda was established as a subgenus of Euchromia Hübner, [1819].

PHAULOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 85, 355.

Type-species: *Eudoliche sordida* Butler, 1878, *Trans. ent. Soc. Lond.* **1878**: 55, by original designation. [BRAZIL]: [Amazon region], Jutahi River.

PHAVARAEA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 315.

Type-species: Centronia rejecta Geyer, 1832, in Hübner, Zuträge Samml. exot. Schmett. 4: 18, [pl.115], figs 663, 664 (but included by Walker as 'Centronia Erynnis Geyer' a misidentification; see below), by monotypy. Type-locality: cited as Java, but this was probably an error as the species illustrated is not known to occur in the Old World, but is distributed in tropical South America.

Walker cited the name erynnis (established as Zygaena erynnis Fabricius, 1777, Genera Insect.: 277, a taxon now placed in the Ctenuchidae) but gave the reference to C. rejecta. Both taxa are of a similar, very distinctive, external appearance. The specimens described by Walker were however of rejecta.

Phavaraea is a nominal genus having a misidentified typespecies. Under the Code, Article 70(a), such cases should be referred to the Commission.

Phavaraea was established as a subgenus of Josia Hübner, [1819], at that time placed in the Lithosiidae; it was placed in the Cyllopodidae, now Geometridae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 401; and was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 426.

SAGARIS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 313.

Type-species: *Josia stygne* Walker, 1854, *ibidem* **2**: 313, by monotypy. [BRAZIL]: Parà.

A junior homonym of Sagaris Panzer, 1806, Kritische Revision Insektenfaune Deutschlands 2: 91, — Insecta, Hymenoptera. There is no objective replacement name but the typespecies of Sagaris Walker, 1854, is conspecific with the typespecies of Phavaraea Walker, 1854. The latter is therefore available for use as a subjective replacement name.

Sagaris was established as a subgenus of Josia Hübner, [1819], at that time placed in the Lithosiidae.

J. stygne is a junior subjective synonym of Centronia rejecta Geyer, 1832.

MILODORA Boisduval, 1870, Considérations Lépid. Guatemala: 85.

Type-species: Centronia rejecta Geyer, 1832, by subsequent designation by Prout, 1918, Novit. zool. 25: 426.

A junior objective synonym of Phavaraea Walker, 1854.

PHECADA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1752.

Type-species: *Phecada confinis* Walker, 1856, *ibidem* 7: 1753, by monotypy. [Brazil]: Rio Janeiro.

P. confinis is a junior subjective synonym of Tepilia biluna Walker, 1855, the type-species of Tepilia Walker, 1855.

Phecada when established contained a second nominal species, but this was doubtfully included and, under the Code, Article 68(c), is not eligible for fixation as type-species.

Phecada was established in the Notodontidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 851; and its type-species was placed in the Lymantriidae by Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 557.

See also Tepilia Walker, 1855.

PHEDOSIA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 691.

Type-species: *Phedosia turbida* Möschler, 1878, *ibidem* 27 (Abh.): 691, pl.10, fig.49, by monotypy. Surinam.

‡PHEGOPTERA; misspelling. See Phaegoptera Herrich-Schäffer, [1853].

PHEIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 146.

Type-species: Glaucopis albisigna Walker, 1854, ibidem 1: 146, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 139. HONDURAS.

Pheia was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 203.

PHIA Zerny, 1912, in Wagner, Lepid. Cat. 7: 61.

An unjustified emendation of Pheia Walker, 1854.

PHELLOE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 320.

Type-species» Dioptis glaucaspis Walker, 1854, ibidem 2: 320, by monotypy. [BRAZIL]: Para.

‡PHELODES Boisduval, 1870. See Dioptis Hübner, 1818.

PHENACOMORPHA Turner, 1940, Proc. R. Soc. Qd 51: 65.

Type-species: *Thallarcha rhabdophora* Turner, 1899, *Trans. Proc. R. Soc. S. Aust.* 23: 19, by monotypy. AUSTRALIA: 'New South Wales (?)'.

‡PHEOSA; misspelling. See Pheosia Hübner, [1819].

PHEOSIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 145.

Type-species: *Phalaena dictaea* Linnaeus, 1767, *Syst. Nat.* (Edn 12) 1 (2): 826, by subsequent designation by Hampson, [1893] 1892, *Fauna Br. India* (Moths) 1: 160. NORTH AFRICA: Barbaria

Incorrect type-species: [Phalaena] tremula Clerck, 1759, a name not originally included in Pheosia, and not linked with one of the two originally included names when designated by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 607.

Hübner established *Pheosia* for *Phalaena dictaea* Linnaeus, 1767, and *Phalaena dictaeoides* Esper, 1789; Kirby placed *P. dictaea* Linnaeus, 1767, as a distinct species without synonyms (but published the name as ‡dictata, an incorrect subsequent spelling) and placed *P. dictaeoides* Esper, 1789, as a junior subjective synonym of *Bombyx gnoma* Fabricius, 1777.

LEIOCAMPA Stephens, 1828, Illust. Br. Ent. (Haustellata)

Type-species: *Phalaena dictaea* Linnaeus, 1767, by subsequent designation by Westwood, 1840, *Introd. mod. Classif. Insects* 2 (Synopsis Genera Br. Insects): 90.

A junior objective synonym of Pheosia Hübner, [1819].

Invalid type-species designation: Phalaena dictaea Linnaeus, 1767, was designated by Boisduval, 1836, Hist. nat. Insectes (Spec. gén. Lépid.) 1: 142. In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different type-species for each generic name. In his 'Exposé de notre méthode', pages 155-690, no type-species designation was made for any of the genera that he himself used. Under the Code, Article 69(a)(iii), the type-species designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other authors. The acceptance of Boisduval's type-species designation in the case of Leiocampa would make no difference to the generic concept.

Leiocampa was again proposed by Stephens, 1829 [June], Nom. Br. Insects: 39; and 1829 [July], Syst. Cat. Br. Insects (2): 42.

‡LEIOCAMPHA; Agassiz, 1846, Nomencl. zool. (Nom. syst. Generum Anim.) Lepid.: 39; 1847, ibidem (Index univl.): 203, 212.

An incorrect subsequent spelling of *Leiocampa* Stephens, 1828. The Lepidoptera part of *Nomencl. zool.* (Nom. syst. Generum Anim.) was included in fascicles 9 and 10 issued in one wrapper dated 1846. The 'Index universalis' formed fascicle 12 of the work; the wrapper is dated 1847, the title-page is dated 1846.

LIOCAMPA Agassiz, 1847, Nomencl. zool. (Index univl.): 203, 212.

An unjustified emendation of *Leiocampa* Stephens, 1828, which was listed by Agassiz as ‡*Leiocampha*, an incorrect subsequent spelling.

The 'Index universalis' formed fascicle 12 of the *Nomencl.* zool.; Liocampa is dated from the wrapper of the fascicle not from its title-page which is dated 1846.

‡PHEOSA; Moore, [1883] 1882-3, Lepid. Ceylon 2: 113. An incorrect subsequent spelling of Pheosia Hübner, [1819].

PHAEOSIA Spuler, 1903, in Hofmann, Schmett. Eur. 1: 95. An unjustified emendation of Pheosia Hübner, [1819], and a junior homonym of Phaeosia Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 128, — Lepid., Arctiidae.

‡PHOESIA; Scheffner, 1927, Societas ent. 42: 5.

An incorrect subsequent spelling of *Pheosia Hübner*, [1819].

‡LEIOCAMPHA; Neave, 1939, Nomencl. zool. 2: 886.

An incorrect subsequent spelling of *Leiocampa* Stephens, 1828. Neave attributed this misspelling to Stephens, 1829, *Syst. Cat. Br. Insects* (2): 42, but at this reference *Leiocampa* was spelt correctly. The error was corrected by Edwards & Hopwood, 1966, in Neave, ibidem 6: 145.

PHEOSILLA Kiriakoff, 1964, Bonn. zool. Beitr. 14: 274.

Type-species: *Pheosilla umbra* Kiriakoff, 1964, *ibidem* 14: 276, abb.4, fig.32, pl.3, fig.34, by original designation. China: Chekiang, West Tien-mu-Shan.

PHEOSINA Gaede, 1916, Int. ent. Z. 9: 125. NOTO Type-species: Pheosina atra Gaede, 1916, ibidem 9: 125, by original designation. [CENTRAL AFRICAN EMPIRE]: Neu Kamerun, Busamtere.

‡PHAEOSINA; Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 433.

An incorrect subsequent spelling of *Pheosina* Gaede, 1916.

PHEOSIOPSIS Bryk, 1950, Ark. Zool. 42 (A) 19: 10. NOTO

Type-species: *Pheosiopsis niveipicta* Bryk, 1950, *ibidem* 42 (A) 19: 11, pl.2, fig.3, by original designation. BURMA: Kambaiti.

PHEOSITIS Strand, 1912, Arch. Naturgesch. 78 (A) 10: 213.

Type-species: *Pheositis excellens* Strand, 1912, *ibidem* 78 (A) 10: 214, by original designation. [TANZANIA] [Tanganyika] D.O. Afrika: Songea.

PHERASPIS Turner, 1903, Proc. Linn. Soc. N.S.W. 28: 46 (key), 61.

Type-species: *Pheraspis polioxutha* Turner, 1903, *ibidem* 28: 62, by original designation. AUSTRALIA: Queensland, Townsville.

‡PHRERASPIS; Turner, 1941, Mem. Qd Mus. 12: 40. An incorrect subsequent spelling of Pheraspis Turner, 1903.

PHERESSACES Turner, 1903, Proc. Linn. Soc. N.S.W. 28: 45 (key), 56.

Type-species: *Notodonta cycnoptera* Lower, 1894, *Trans. Proc. R. Soc. S. Aust.* 18: 78, by original designation. AUSTRALIA: Queensland, Duaringa.

PHIA Zerny, 1912. See Pheia Walker, 1854. CTEN

PHIALA Wallengren, 1860, Wien. ent. Monatschr. 4: 165.
EUPTEROTIDAE

Type-species: *Phiala xanthosoma* Wallengren, 1860, *ibidem* 4: 165, by original designation (in accordance with the *Code*, Article 68(a)(i)). [SOUTH AFRICA] Caffraria.

Phiala when established was not placed in a family; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 446; and was placed in the Striphnopterygidae, now Eupterotidae, by Aurivillius, 1901, Bihang K. svenska Vetensk Akad. Handl. 27 (Afd.4) 7: 15.

PHIDITIA Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 338.

Type-species: *Phalaena diores* Cramer, 1775, *Uitlandsche Kapellen (Papillons exot.)* 1: 119, pl.75, fig.F, by monotypy. SURINAM.

P. diores is dated from the wrapper of the part, and not from the title-page of the volume, 1779.

PHILAGRIA Kirby, 1892. See Oeonistis Hübner, [1819].

PHILARETA Moore, 1881, Proc. zool. Soc. Lond. 1881: 361.
NOCTUIDAE

Type-species: Heliothis taurica Herrich-Schäffer, 1846, Syst. Bearbeitung Schmett. Eur. 2: Noctuides pl.52, figs 263, 264; 1851, ibidem 2: 368, by monotypy. Type(s) Type-locality: cited by Herrich-Schäffer, 1851, ibidem 2: 369, as 'Nach Parreyss aus Corsika', but this was probably an error as this species is not known to occur in Corsica or the western Mediterranean, but is present as its name implies in Turkey: Taurus Mts.

H. taurica is dated from the original wrappers of the parts containing it and not from the title-page, 1845.

H. taurica is a junior subjective synonym of Heliothis treitschkii Frivaldszky, 1835, Magy. Tud. Tars. Evk. 2 (2): 273, pl.7, fig.8, (but cited as ‡Heliotis, an incorrect subsequent spelling).

Philareta in the Noctuidae, Noctuinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PHILEA Zetterstedt, [1839]. See Setina Schrank, 1802.

PHILENORA Rosenstock, 1885, Ann. Mag. nat. Hist. (5) 16: 382.

Type-species: Acontia undulosa Walker, [1858] 1857, List Specimens lepid. Insects Colln Br. Mus. 12: 797, by monotypy. [Australia]: Tasmania.

‡*PHILLOECIA*; misspelling. See *Phylloecia* Guérin-Méneville, [1844].

‡PHILODONTIS; misspelling. See Euchila Billberg, 1820.

‡PHILOMACRA; misspelling. See Ptilomacra Walker, 1855.

PHILONA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 456.

Type-species: *Hypsa inops* Walker, 1854, *ibidem* 2: 457, by monotypy. [BANGLADESH]: [Sylhet] Silhet.

Philona was established as a subgenus of Hypsa Hübner, [1819].

PHILOROS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 283.

Type-species: Ctenucha venosa Walker, 1854, ibidem 2: 284, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 176. VENEZUELA, MEXICO, and an unstated locality.

Philoros was established as a subgenus of Ctenucha Kirby, 1837.

PHILORUS Zerny, 1912, in Wagner, Lepid. Cat. 7: 152.

An unjustified emendation of Philoros Walker, 1854.

A junior homonym of *Philorus* Kellog, 1903, *Proc. Calif. Acad. Sci.* (3) 3: 199, – Insecta, Diptera. The objective replacement name is *Philoros* Walker, 1854.

PHILORUS Zerny, 1912. See Philoros Walker, 1854. CTEN

PHINTIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 306.

Type-species: *Josia podarce* Walker, 1854, *ibidem* 2: 306, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 403. [Brazil]: Para.

PHISSAMA Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 362.

Type-species: Amphissa vacillans Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 685, by monotypy (of Amphissa Walker, 1855). Hong Kong.

Phissama was established as the objective replacement name for Amphissa Walker, 1855.

A. vacillans is a junior subjective synonym of Spilosoma transiens Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 675.

AMPHISSA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 684.

Type-species: Aniphissa vacillans Walker, 1855, by monotypy.

A junior homonym of *Amphissa* Adams & Adams, 1853, *Genera recent Mollusca* 1: 111, – Mollusca. The objective replacement name is *Phissama* Moore, 1858.

‡PHLAEOCIILAENA; misspelling. See Phaeochlaena Hübner, 1818.

‡PHLAEOCHLOENA; misspelling. See Phaeochlaena Hübner, 1818. DIOP

PHLOGOMERA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxviii, 797.

Type-species: Asura bicolor Rothschild, 1913, Novit. zool.

20: 210, by original designation. New GUINEA: [Irian Jaya], Snow Mts, near Octakwa River and upper Octakwa River.

PHLYCTAENOGASTRA Gaede, 1915, Dt. ent. Z. Iris 29: 145.

Type-species: *Phlyctaenogastra rangei* Gaede, 1915, *ibidem* **29**: 145, by monotypy. [Namibia] South West Africa: Churutabis.

‡PHYCTAENOGASTRA; Neave, 1940, Nomencl. zool. 3: 720. An incorrect subsequent spelling of Phlyctaenogastra Gaede, 1915.

PHOEGOPTERA Boisduval, 1869, Annls Soc. ent. Belg. 12: 180.

Type-species: *Phoegoptera salicis* Boisduval, 1869, *ibidem* 12: 81, by subsequent designation by Watson, 1973, *Smithson. Contr. Zool.* 128: 2. [U.S.A.]: California.

P. salicis is a junior subjective synonym of Lophocampa maculata Harris, 1841, Rep. Insects Mass. injurious to Vegn: 258.

Phoegoptera Boisduval, 1869, and the similar name Phaegoptera Herrich-Schäffer, [1853], are discussed by Watson, 1973, ibidem 128: 2.

‡PHOEGORISTA; misspelling. See Phaegorista Boisduval, [1836].

PHOENICOPROCTA Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 404.

Type-species: *Phoenicoprocta metachrysea* Druce, 1898, *ibidem* (7) 1: 404, by monotypy. [French Guiana]: Cayenne.

The name *Phoenicoprocta* probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson, 1898 December, as the name for a new genus.

P. metachrysea is a junior subjective synonym of Eunomia vacillans Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1617.

HYELA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 172.

Type-species: Glaucopis sanguinea Walker, 1854, ibidem 1: 172, by monotypy. Type-locality: not stated, but the holotype has been subsequently labelled 'Honduras', [AMERICA]: [tropical].

Hyela was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 196.

A junior homonym of *Hyela* Stephens, 1850, *List Specimens Br. Anim. Colln Br. Mus.* 5: 149, – Lepid., Noctuidae. There is no objective replacement name but the type-species of *Hyela* Walker, 1854, is currently considered to be congeneric with the type-species of *Phoenicoprocta* Druce, 1898. The latter is therefore available for use as a subjective replacement name.

PHOENICOPROCTA Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xiii, 23, 196.

Type-species: Glaucopis sanguinea Walker, 1854, by monotypy (of Hyela Walker, 1854).

Phoenicoprocta was established as the objective replacement name for Hyela Walker, 1854.

A junior homonym of *Phoenicoprocta* Druce, 1898 May. There is no objective replacement name but the type-species of *Phoenicoprocta* Hampson is currently considered to be congeneric with the type-species of *Phoenicoprocta* Druce. The latter is therefore available for use as a subjective replacement

PHOENICOPROCTA Hampson, 1898 [December]. See Phoenicoprocta Druce, 1898 May.

‡PHOENICOSTACTA; misspelling. See Phoeniostacta Hampson, 1898.

PHOENIOSTACTA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xiv, 259.

Type-species: *Phoeniostacta haematobasis* Hampson, 1898, *ibidem* 1: 260, by original designation. BRAZIL: [Teresopolis] Therosopolis.

The spelling *Phoeniostacta* is here chosen from a multiple original spelling of both *Phoeniostacta* and ‡*Phoenicostacta*.

‡PHOENICOSTACTA; Hampson, 1898, Cat. Lepid. Pha-laenae Br. Mus. 1: 26 (key).

An incorrect (of a multiple) original spelling of *Phoeniostacta* Hampson, 1898.

‡PHOESIA; misspelling. See Pheosia Hübner, [1819].

PHRAGMATOBIA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 55, 73.

Type-species: *Phalaena fuliginosa* Linnaeus, 1758, *Syst. Nat.* (Edn 10): 509, by monotypy. Type-Locality: not stated, [Europe].

Phragmatobia was proposed again as a new name by Stephens, 1829 [June], Nom. Br. Insects: 40; and 1829 [July], Syst. Cat. Br. Insects (2): 55.

PHREATA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 403.

Type-species: *Phreata glaucoalba* Walker, 1865, *ibidem* 32: 404, by monotypy. Bolivia.

‡PHRERASPIS; misspelling. See Pheraspis Turner, 1903.

PHRYGANEOMORPHA Wallengren, 1858, Ofvers. K. Vetensk Akad. Förh. Stockh. 15: 214.

Type-species: Chelonia madagascariensis Boisduval, 1847, in Delegorgue, Voyage Afr. australe 2: 598, by original designation. MADAGASCAR.

See also Amblythyris Mabille, 1879.

PHRYGANIDIA Packard, 1864, Proc. ent. Soc. Philad. 3: 348.

Type-species: *Phryganidia californica* Packard, 1864, *ibidem* 3: 349, by monotypy. [U.S.A.]: California.

Phryganidia was established in the Psychidae; it was placed in the Zygaenidae by Stretch, 1872, Illust. Zygaenidae Bombycidae N. Am. 1: 91; and placed in the Dioptidae by Grote, 1882, New Check List N. Am. Moths: 18.

PHRYGANOPSIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.30.

Type-species: *Phryganopsis sordida* Felder, 1874, *ibidem* 2 (Abt.2): pl.106, fig.30, by monotypy. [SOUTH AFRICA]: [Cape of Good Hope] Cap. b. sp., cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 8.

‡PHRYGANOPTERA; misspelling. See Phryganopteryx Saalmüller, 1884.

PHRYGANOPTERYX Saalmüller, 1884, Lepid. Madagascar (Abt.1): 155.

Type-species: *Macrobrochis strigilata* Saalmüller, 1878, *Ber. senckenb. naturf. Ges.* **1877-78**: 91, by monotypy. MADAGASCAR.

MAXIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 206.

An unnecessary replacement name for Phryganopteryx Saal-

müller, 1884, which is not preoccupied by *Phryanopterus* Kolenati, 1860, *Wien. ent. Monatschr.* 4: 393, – Insecta, Diptera.

A junior homonym of *Maxia* Heyden, 1891, *in* Saalmüller, *Lepid*. *Madagascar* 2: 488, – Lepid., Noctuidae.

‡PHRYGANOPTERA; Rothschild, 1933, Ann. Mag. nat. Hist. (10) 11: 188.

An incorrect subsequent spelling of *Phryganopteryx* Saalmüller, 1884.

PHRYNE Grote, 1865. See Calybia Kirby, 1892.

PHUSIANA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 141.
NOTO

Type-species: *Phusiana albifrons* Walker, 1862, *ibidem* (Zool.) 6: 141, by monotypy. Borneo: Sarawak.

Study of the type-specimen of *P. albifrons* in the University Museum, Oxford, has proved it to be a senior subjective synonym of *Phalera javana amplificata* Gaede, 1930, in Seitz, *Gross-Schmett. Erde* 10: 613. Syn. n.

PHYA Druce, 1887. See Ianassa Walker, 1855. NOTO

PHYCIDOPSIS Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 16, 91. NOTO

Type-species: *Phycidopsis albovittata* Hampson, 1893, *ibidem* 9: 16, 91, pl.161, fig.13, by monotypy. [SRI LANKA] Ceylon.

Phycidopsis was established in the Noctuidae; it was placed in the Notodontidae by Hampson, 1897, J. Bombay nat. Hist. Soc. 11: 281.

PHYCITIMORPHA Janse, 1920, Ann. Transv. Mus. 7: 208.

Type-species: *Phycitimorpha stigmatica* Janse, 1920, *ibidem* 7: 208, pl.8, figs 1-5, pl.14, fig.8, by original designation. [ZIMBABWE-]RHODESIA: Umvuma.

‡PHYCTAENOGASTRA; misspelling. See Phlyctaenogastra Gaede, 1915.

‡PHYLLAECIA; misspelling. See Phylloecia Guérin-Méneville, [1844].

PHYLLALIODES Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 474.

Type-species: *Phyllaliodes agramma* Hampson, 1910, *ibidem* (8) 5: 474, by original designation. [SOUTH AFRICA]: Natal, Pt Shepstone.

PHYLLOECIA Guérin-Méneville, [1844] 1829-1838, Icon. Règne anim. Cuvier 3 (Insectes): 504.

Type-species: *Phylloecia punctata* Guérin-Méneville, [1844] 1829-1838, *ibidem*: 504, fig., by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 129. [Mexico]: Baiade Campeche.

The spelling *Phylloecia* was adopted from a multiple original spelling by Sherborn, 1929, *Index Anim.* (1801-1850): 4913, 4931, as first reviser.

P. punctata is a junior subjective synonym of Sphinx plumipes Drury, 1773, Illust. nat. Hist. exot. Insects 2: index, 47, pl.27, fig.3.

‡PHILLOECIA; Guérin-Méneville, [1844] 1829-1838, Icon. Règne anim. Cuvier 3 (Insectes): 504.

An incorrect (of a multiple) original spelling of *Phylloecia* Guérin-Méneville, [1844].

‡PHYLLAECIA; Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 248, 252.

An incorrect subsequent spelling of *Phylloecia* Guérin-Méneville, [1844].

PHYLLOPALPIA Draudt, 1932, in Seitz, Gross-Schmett. Erde 6: 939.

Type-species: *Phyllopalpia triangulum* Draudt, 1932, *ibidem* 6: 939, pl.147 row a, by original designation. South America.

The date of publication of the part of Seitz containing page 939 was cited on page 937 as 16.iii.1932 in both the German and the English editions.

PHYSETOCNEME Felder, 1861, Sher. Akad. Wiss. Wien 43 (Abt.1): 37.

Type-species: *Physetocneme ciliosa* Felder, 1861, *ibidem* 43 (Abt.1): 38, by monotypy. [Indonesia]: [Ambon I.] Aboinensium.

‡PHYTOCNEMA; Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 276.

An incorrect subsequent spelling of *Physetocneme* Felder, 1861.

‡PHYTOCNEMA; misspelling. See Physetocneme Felder, 1861.

PIDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 399.

Type-species: *Pida apicalis* Walker, 1865, *ibidem* 32: 400, by monotypy. [INDIA]: [W. Bengal], Darjeeling.

PIESTA Billberg, 1820, Enumeratio Insect. Mus. G.J. Billberg: 91. OECOPHORIDAE

Type-species: Haemilis pastinacella Duponchel, 1838, in Godart & Duponchel, Hist. nat. Lépid. Papillons Fr. 11: 153, by subsequent designation by Walsingham, 1907, Proc. zool. Soc. Lond. 1907: 958, as Phalaena heracliana Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 532, (but cited as ‡heracleana, an incorrect subsequent spelling), a misidentification of H. pastinacella Duponchel. This is a case of a misidentified type-species which should be placed before the Commission under Article 70(a) of the Code. France.

The name *Piesta* has been used in the Arctiidae based on the later type-species designation by Fletcher, 1966, *Entomologist's Gaz.* 17: 16 of *Phalaena deplana* Linnaeus, [1771], *Mantissa Plantarum* 2: 539, – Arctiidae, Lithosiinae.

‡PIGETERA; misspelling. See Cataphractes Felder, 1874.

PIKA Matsumura, 1929. See Tarsolepis Butler, 1872. NOTO

PILODONTA Dognin, 1911, Hétérocères nouv. Am. Sud 2: 42.

Type-species: *Pilodonta subcrinita* Dognin, 1911, *ibidem* 2: 43, by original designation. FRENCH GUIANA.

PINHEYA Kiriakoff, 1973. See Pinheyata Nye nom. n. THYRET

PINHEYATA Nye nom. n. for Pinheya Kiriakoff, 1973.

Type-species: Automolis quinta Kiriakoff, 1973, Mitt. münch. ent. Ges. 63: 64, fig.9, by original designation (for Pinheya Kiriakoff, 1973). [ZIMBABWE-]RHODESIA: Vumba Mts, Umtali.

Prof. Dr Kiriakoff was informed of this homonymy and suggested the replacement name.

PINHEYA Kiriakoff, 1973, Mitt. münch. ent. Ges. 63: 63.

Type-species: Automolis quinta Kiriakoff, 1973, by original designation.

A junior homonym of *Pinheya* Dlabola, 1963, *Publções cult*. *Co. Diam. Angola* 66: 51, – Insecta, Hemiptera. The objective replacement name is *Pinheyata* Nye nom. n.

Pinheya was established as a subgenus of *Automolis* Hübner, [1819].

PINHEYIA Kiriakoff, 1971, Bull. Annls Soc. r. ent. Belg. 107:

Type-species: *Pinheyia lymantrioides* Kiriakoff, 1971, *ibidem* **107**: 318, text-fig.2, by original designation. MOZAMBIQUE: Chirondo, 30 miles east of Inhamingo.

PIONIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 255.

Type-species: *Euchromia lycoides* Walker, 1854, *ibidem* 1: 256, by monotypy. VENEZUELA.

Pionia was established as a subgenus of *Euchromia* Hübner, [1819].

See also Correbia Herrich-Schäffer, [1855].

PIRGA Aurivillius, 1891, Ent. Tidskr. 12: 228.

Type-species: Pirga mirabilis Aurivillius, 1891, ibidem 12:

228, pl.2, fig.2, by monotypy. [GABON] Gabun.

Pirga and P. mirabilis were made nomenclaturally available by a figure on a plate and its legend. The written descriptions were published a year later by Aurivillius, 1892, *ibidem* 13: 192, 193.

‡PIRGALASEA; misspelling. See Somatoxena Aurivillius, 1904.

PIRGULA Tessmann, 1921, Mitt. zool. Mus. Berl. 10: 215.

Type-species: *Pirgula decempunctata* Tessman, 1921, *ibidem* **10**: 216, by original designation. [Tanzania]: Tabora; Ipala, S.W. Kalitu.

P. decempunctata is a junior subjective synonym of Antiphella atrinotata Butler, 1896, Proc. zool. Soc. Lond. 1896: 854, pl.43 fig.5.

Pirgula was established in the Pterothysanidae; P. decempunctata was placed in the Lymantriidae by Collenette, 1931, Ann. Mag. nat. Hist. (10) 7: 511.

PIRGULINA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 125.

Type-species: *Pirgulina jordani* Hering, 1926, *ibidem* 14: 125, pl.19, row b, by monotypy. MADAGASCAR: Diego Suarez.

The date of publication of the part of Seitz containing page 125 was cited on page 121 as 9.x.1926 in both the German and the English editions.

Pirgulina was established in the Pterothysanidae, but was placed in the Lymantriidae by Collenette, 1953, Ann. Mag. nat. Hist. (12) 6: 567.

PITANE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 462. ARCT

Type-species: *Pitane fervens* Walker, 1854, *ibidem* **2**: 462, by monotypy. Type-locality: not known, [South America]: [tropical].

PITANE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 531. See Nepita Moore, 1858.

PITASILA Moore, 1877, Proc. zool. Soc. Lond. 1877: 599.

Type-species: *Pitasila leucospilota* Moore, 1877, *ibidem* **1877**: 599, pl.59, fig.7, by monotypy. [INDIA]: Andaman Is, Port Blair.

PITHEA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1701. ARCT

Type-species: *Pithea ferruginea* Walker, 1856, *ibidem* 7: 1701, by monotypy. [COLOMBIA]: Santa Fé de Bogotà.

PITTHEIDES Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 58.

Type-species: Somera chloauchena Holland, 1893, Psyche,

Camb. 6: 475, pl.17, fig.18, by original designation. [GABON]: [Ogooué] Ogové R. valley.

PLASTYSTAURA Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 338.

Type-species: *Plastystaura murina* Kiriakoff, 1965, *ibidem* 101: 339, fig.6, by original designation. [ZIMBABWE-]RHODE-SIA: Wankie.

PLATARCTIA Packard, 1864, Proc. ent. Soc. Philad. 3: 109.

Type-species: Arctia parthenos Harris, 1850, in Agassiz, Cabot et al, Lake Superior: 390, pl.7, fig.4, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 257. CANADA: northern shore of Lake Superior.

PLATYCERURA Packard, 1864, Proc. ent. Soc. Philad. 3: 373.
NOCTUIDAE

Type-species: Platycerura furcilla Packard, 1864, ibidem 3:

374, by monotypy. [U.S.A.]: Massachusetts.

Platycerura was established in the Notodontidae; it was placed in the Noctuidae by Grote, 1882, New Check List N. Am. Moths: 22.

PLATYCHASMA Butler, 1881, Trans. ent. Soc. Lond. **1881**: 596. NOCTUIDAE

Type-species: Platychasma virgo Butler, 1881, ibidem 1881:

596, by monotypy. JAPAN: Tokyo.

Platychasma was established in the Notodontidae; it was transferred to the Noctuidae by Kiriakoff, 1963, J. Lepid. Soc. 17: 33.

PLATYODONTA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.16. NOTO

Type-species: *Platyodonta calpe* Felder, 1874, *ibidem* 2 (Abt.2): pl.97, fig.16, by monotypy. [BRAZIL]: River Amazon, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 8.

PLATYPREPIA Dyar, 1897, Can. Ent. 29: 212, 215. ARCT Type-species: Chelonia virginalis Boisduval, 1852, Annls Soc. ent. Fr. (2) 10: 321, by monotypy. U.S.A.: California, near San Francisco.

‡PLEREDES; misspelling. See Pericallia Hübner, [1820].

PLERETES Lederer, 1853. See Pericallia Hübner, [1820].

PLEUROMNEMA Gaede, 1914, Int. ent. Z. 8: 127. AGAN Type-species: Pleuromnema reussi Gaede, 1914, ibidem 8: 127, fig. [TANZANIA]: Dar es Salaam.

PLEUROSOMA Orfila, 1935, Revta Soc. ent. argent. 7: 178.

Type-species: *Sphecosoma angustata* Möschler, 1878, *Verh. zool.-bot. Ges. Wien* **27** (Abh.): 634, pl.8, fig.2, by original designation. Surinam.

Not preoccupied by ‡*Pleurosoma*, Guérin-Méneville, 1844, *Magasin Zool. Paris* (2) 6: pl.136, – Insecta, Coleoptera, an incorrect subsequent spelling of *Pleurasoma* Guérin-Méneville, 1844, *Revue zool.* 7: 9.

ASTRIDIA Kiriakoff, 1948, Bull. Annls Soc. r. ent. Belg. 84: 267.

Type-species: Sphecosoma angustata Möschler, 1878, by original designation.

Astridia is a junior objective synonym of Pleurosoma Orfila, 1935.

PLINIOLA Strand, 1912, Arch. Naturgesch. 78 (A) 7: 185.

Type-species: *Tegulata nigristriata* Holland, 1893, *Ent. News* 4: 341, pl.15, fig.10, by original designation. [GABON]: [Ogoové river valley], Kangwe.

PLUMAREOLA van Eecke, 1920, Zool. Meded. Leiden 5: 137.

Type-species: *Plumareola lineola* van Eecke, 1920, *ibidem* 5: 137, fig., by monotypy. JAVA: W., Preanger; Malabargebergte, Tjinjiroean.

PLUSIOGRAMMA Hampson, 1895, Trans. ent. Soc. Lond. 1895: 278.

Type-species: *Plusiogramma aurosigna* Hampson, 1895, *ibidem* **1895**: 278, fig., by original designation. [BURMA]: Tenasserim, Donaut Range.

PODOCRYPTULA Bryk, 1950, Ark. Zool. 42 (A) 19: 44.
NOTO

Type-species: *Podocryptula nana* Bryk, 1950, *ibidem* **42** (A) 19: 44, pl.1, fig.7, by original designation. [BURMA]: Tenasserim, Sukli, 75 km east of Moulmein.

PODOMACHLA Strand, 1909, Annls Soc. ent. Belg. 53: 326, 339.

Type-species: *Nyctemera apicalis* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 2: 395, by PRESENT DESIGNATION. [GHANA]: Ashanti, [SOUTH AFRICA]: Zululand and [Durban] Port Natal, SIERRA LEONE.

Podomachla was established as a subgenus of Deilemera Hübner, [1820].

POECILARCTIA Aurivillius, 1921, Ann. S. Afr. Mus. 18: 235.

Type-species: *Poecilarctia venata* Aurivillius, 1921, *ibidem* 18: 235, by monotypy. [ZIMBABWE-]RHODESIA: Salisbury.

POECILOPHEOSIA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 203.

Type-species: *Pheosia grisea* Swinhoe, 1892, *Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus.* 1: 298, pl.8, fig.7, by original designation (but cited by Kiriakoff as *Pheosia excurvata* Hampson, a misidentification; see below). INDIA.

After examining the relevant type material and genitalia slides, my colleague Mr T. Willett-Whittaker has found that the generic diagnosis of *Poecilopheosia* and the genitalia illustration that accompanied it refer to *Pheosia grisea* Swinhoe, 1892, and not to *Pheosia excurvata* Hampson, [1893] 1892, *Fauna Br. India* (Moths) 1: 161. Thus *Pheosia excurvata* Hampson sensu Kiriakoff, 1968 = *Pheosia grisea* Swinhoe, 1892. He has also found that *Pheosia excurvata* and *Pheosia grisea* are not congeneric; *Pheosia excurvata* Hampson, [1893], should be placed in the genus *Fentonia* Butler, 1881, close to *Harpyia ocypete* Bremer, 1861. Under the *Code*, Article 70(a), the case of a misidentified type-species should be referred to the Commission.

POECILOSOMA Hübner, [1819] 1816, Verz. bekannter Schmett.: 126. Available but without included available species until Hübner, 1823, Zuträge Samml. exot. Schmett. 2: 8.

Type-species: *Poecilosoma chrysis* Hübner, 1823, *ibidem* 2: 8, pl.[37], figs 211, 212, by subsequent monotypy. Surinam. *Poecilosoma* Hübner was nomenclaturally available from [1819] but both the included nominal species, ‡*P. chrysis* and ‡*P. pulchra*, were *nomina nuda*.

POLELASSOTHYS Janse, 1920, Ann. Transv. Mus. 7: 229.

Type-species: *Polelassothys plumitarsus* Janse, 1920, *ibidem* 7: 230, pl.12, figs 2-8, pl.14, fig.17, by original designation. [ZIMBABWE-]RHODESIA: Sawmills.

POLIENUS Distant, 1903, Insecta Transv. 1: 93. NOTO Type-species: Polienus modestus Distant, 1903, ibidem 1: 94, fig.10, by monotypy. [SOUTH AFRICA]: Transvaal, Pretoria.

P. modestus is a junior subjective synonym of Simyra capillata Wallengren, 1875, Ofvers K. Vetensk Akad. Förh. Stockh. 32 (1): 104.

POLIODULE Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 90, 339.

Type-species: *Scoliacma xanthodelta* Lower, 1897, *Proc. Linn. Soc. N.S.W.* **22**: 10, by original designation. AUSTRALIA: New South Wales, Broken Hill.

‡POLIOPASTA; misspelling. See Poliopastea Hampson, 1898.

POLIOPASTEA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xvi, 26, 336.

Type-species: *Poliopastea plumbea* Hampson, 1898, *ibidem* 1: 337, pl.12, fig.26, by original designation. [BRAZIL]: Amazon basin, Paratins.

‡POLIOPASTA; Rothschild, 1911, Novit. zool. 18: 40.

An incorrect subsequent spelling of *Poliopastea* Hampson, 1898, incorrectly attributed to 'Kirby 1911, Novit. zool. 18, 40' by Neave, 1940, *Nomencl. zool.* 3: 845.

POLIOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 82, 106.

Type-species: *Lithosia muricolor* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 105, by original designation. BORNEO: Sarawak.

POLIOSTAUROPUS Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 192. NOTO

Type-species: Stauropus griseus Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 59, pl.141, fig.16 (legend misspelt as ‡grisea), by original designation. INDIA: southern, Nilgiri district.

POLYAESCHRA Kiriakoff, 1967, Tijdschr. Ent. 110: 60.

Type-species: *Hyperaeschra dentata* Hampson, [1893 January 10] 1892, *Fauna Br. India* (Moths) 1: 165, by original designation. [SRI LANKA] Ceylon: Trincomalee.

H. dentata was again published as a new species by Hampson, 1893 [February 25], Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 5, 60, pl.160, fig.11.

See also Allodontoides Matsumura, 1922.

POLYCHOA Turner, 1907, Proc. Linn. Soc. N.S.W. 31: 681.

Type-species: *Polychoa styphlopis* Turner, 1907, *ibidem* 31: 681, by monotypy. Australia: Queensland, Kuranda.

POLYMONA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 768.

Type-species: *Polymona rufifemur* Walker, 1855, *ibidem* 3: 768, by monotypy. South Africa: interior.

Polymona was established in the Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 481.

See also Oreinobia Wallengren, 1865.

‡POLYPAETES; misspelling. See Polypoetes Druce, 1885.

POLYPOETES Druce, 1885, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 159.

Type-species: *Polypoetes deldon* Druce, 1885, *ibidem* 1: 160, pl.14, fig.24, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 414. GUATEMALA: Cerro Zunil, Las Mercedes, Pantaleon.

ARINA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1662.

Type-species: Arina obtusa Walker, 1856, ibidem 7: 1663,

by monotypy. Brazil.

A junior homonym of Arina Robineau-Desvoidy, 1830, Mém. prés. div. Sav. Acad. Sci. Inst. Fr. 2: 696, — Insecta, Diptera. There is no objective replacement name but A. obtusa is currently treated as congeneric with Polypoetes deldon Druce, 1885, the type-species of Polypoetes Druce, 1885; the latter is therefore available for use as a subjective replacement name.

‡POLYPAETES; Dognin, 1894, Lépid. Loja et environs (Equateur) (3): pl.9, fig.13, legend.

An incorrect subsequent spelling of Polypoetes Druce, 1885.

‡POLYPTOETES; Dognin, 1894, Lépid. Loja et environs (Equateur) (3): explan. pl.9.

An incorrect subsequent spelling of *Polypoetes* Druce, 1885.

‡POLYPTOETES; misspelling. See Polypoetes Druce, 1885.

POLYPTYCHIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.104, fig.7.

Type-species: *Polyptychia fasciculosa* Felder, 1874, *ibidem* **2** (Abt.2): pl.104, fig.7, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, *ibidem* **2** (Abt.2) Erklärung to pls 75-107: 8

Prout, 1918, *Novit. zool.* **25**: 428, proposed *Polyptychia* Felder as a new genus because he believed Felder's publication of the generic name to be invalid, as it was not accompanied by a description; according to the present *Code*, Article 13, this requirement applies only to names published after 1930.

POLYSTICTINA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 178.

Type-species: Heterocampa maculata Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 60, by monotypy. INDIA: Darjeeling.

POMPILIODES Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xii, 24, 162.

Type-species: Euchromia aliena Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 241, by original designation. [BRAZIL]: Para.

‡POMPILIOPSIS; misspelling. See Pompilopsis Hampson, 1898.

POMPILOPSIS Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xii, 161.

Type-species: Glaucopis tarsalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 196, by original designation. [Brazil]: Para.

The spelling *Pompilopsis* is here chosen from a multiple original spelling which included *Pompilopsis* (cited twice) and ‡*Pompiliopsis* (cited once).

‡POMPILIOPSIS; Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 24.

An incorrect (of a multiple) original spelling of *Pompilopsis* Hampson, 1898.

POMPOSTOLA Hübner, [1819] 1816, Verz. bekannter Schmett.: 120. ZYGAENIDAE

Type-species: Sphinx hipparchus Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 7, pl.197, fig.C, by subsequent designation by Butler, 1876, J. Linn. Soc. (Zool.) 12: 421. SIERRA LEONE.

Pompostola has been used as the original generic name for species now placed in the Ctenuchidae.

S. hipparchus is dated from the wrapper of the part and not from the title-page of the volume, 1782.

PONCETIA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 152 (key), 177.

Type-species: Niganda albistriga Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 64, by original designation. INDIA: Darjeeling.

Poncetia was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 68.

PONEOPIRGA Berio, 1937, Annali Mus. civ. Stor. nat. Giacomo Doria 59: 373.

Type-species: Poneopirga sakania Berio, 1937, ibidem 59: 374, by original designation. [ZAIRE] Congo Belga: Katanga, Sakania.

PONTALA Walker, 1864, List Specimens lepid. Insects Colln Br. Mus. 30: 954.

Type-species: Pontala rubrana Walker, 1864, ibidem 30: 954,

by monotypy. [BRAZIL]: Para.

Pontala was established in the Crambites, now Pyralidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 326.

PONTALAMIA Bryk, 1953, Ark. Zool. (2) 5: 166. Type-species: Pontalamia grotesca Bryk, 1953, ibidem (2) 5: 166, by original designation. BRAZIL: E. Tabatinga, Solimões, PERU: W. Tabatinga, Iquitos.

PORELA Walker, 1855 [June 9], List Specimens lepid. Insects Colln Br. Mus. 3: 772. LASIOCAMPIDAE

Type-species: Porela vetusta Walker, 1855, ibidem 3: 772, by monotypy. [AUSTRALIA]: New South Wales, Sydney.

Porela was established in the Arctiidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 843.

CALLIA Walker, 1855 [December 8], List Specimens lepid. Insects Colln Br. Mus. 6: 1387 (key), 1482.

Type-species: Callia albifinis Walker, 1855, ibidem 6: 1483, by monotypy. [Australia]: [New South Wales], Sydney.

A junior homonym of Callia Audinet-Serville, 1835, Annls Soc. ent. Fr. 4: 60, - Insecta, Coleoptera. There is no objective replacement name, but C. albifinis is congeneric with Porela vestusta Walker, 1855, the type-species of Porela Walker, 1855 [June 9]; the latter is thus available as a subjective replacement name.

Callia was established in the Bombycidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892 [July 9], Synonymic Cat. Lepid. Heterocera 1: 469; and was placed in the Lasiocampidae by Swinhoe, 1892 [November 4], Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 309.

PORESTA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 259

Type-species: Edema lanassa Druce, 1890, Proc. zool. Soc. Lond. 1890: 509, by original designation. Panama: Chiriqui.

PORPHYROCHRYSA Turner, 1940, Proc. R. Soc. Qd 51: 114.

Type-species: Porphyrochrysa dochmoschema Turner, 1940, ibidem 51: 114, by monotypy. Australia: N. Queensland, Cape York.

PORSICA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1823.

Type-species: Porsica ingens Walker, 1866, ibidem 35: 1823,

by monotypy. [BANGLADESH]: Sylhet.

Porsica was established in the Tineidae; it was transferred to the Notodontidae by Butler, 1886, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 6: vii, 21.

PORTHESAROA Hering, 1926, in Seitz, Gross-Schmett. Erde **14**: 151.

Type-species: Porthesaroa aureopsis Hering, 1926, ibidem 14: 151, pl.28, row e, by original designation. MADAGASCAR: Diego Suarez.

The date of publication of the part of Seitz containing page 151 was cited on page 145 as 4.xi.1926 in both the German and the English editions.

PORTHESIA Stephens, 1828. See Euproctis Hübner, [1819].

PORTHETRIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 160.

Type-species: Phalaena dispar Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 501, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 475. Type-locality: not stated, [EUROPE].

SERICARIA Berthold, 1827, in Latreille, Nat. Familien Thierreichs: 480. Available but without included species until Latreille, 1829, in Cuvier, Règne Animal (Edn 2) 5: 404.

Type-species: Phalaena dispar Linnaeus, 1758, by subsequent designation by Blanchard, [1847], in Cuvier, Règne Animal (Disciples Edn) 14 (Insectes, Orthoptères-Diptères) Atlas: titlepage, pl.152.

Sericaria is a junior objective synonym of Porthetria Hübner, [1819].

PORTHMEIA Bethune-Baker, 1908, Novit. zool. 15: 186. LYMA

Type-species: Porthmeia subnigra Bethune-Baker, 1908, ibidem 15: 186, by original designation. New Guinea: [Papua New Guinea], [Papua], Owgarra.

Incorrect type-species: Porthmeia bicolora Bethune-Baker, 1908, cited by Bryk, 1934, in Strand, Lepid. Cat. 62: 214.

POSTANITA Dyar, 1918, Proc. U.S. natn. Mus. 54: 357.

Type-species: Postanita decurrens Dyar, 1918, ibidem 54: 357, by original designation. MEXICO.

PRABHASA Moore, 1878, Proc. zool. Soc. Lond. 1878: 25.

Type-species: Prabhasa venosa Moore, 1878, ibidem 1878: 26, pl.2, fig.16, by subsequent designation by Hampson, 1894, Fauna Br. India (Moths) 2: 76. INDIA: Darjeeling.

‡PRAEMOLIS; misspelling. See Premolis Hampson, 1901. ARCT

PRAESCHAUSIA Benjamin, 1932, Pan-Pacif. Ent. 8: 57.

Type-species: Kalkoma zapata Schaus, 1920, Proc. U.S. natn. Mus. 57: 140, by original designation. Mexico: Presidio River, Sinaloa.

PREMOLIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: [vii], 2, 26.

Type-species: Halysidota semirufa Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1708 (but cited as ‡Halesidota by Walker, an incorrect subsequent spelling), by original designation (but cited as \$\pm\$Halesidota). [Brazil]:

‡PRAEMOLIS; Rothschild, 1909, Novit. zool. 16: 52. An incorrect subsequent spelling of *Premolis* Hampson, 1901.

PREPARCTIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 5, 219.

Type-species: Chelonia mirifica Oberthür, 1892, Etudes d'Entomologie 16: 8, pl.1, fig.7, by original designation. CHINA: [Szechwan] Ta-tsien-lou.

PREPIELLA Schaus, 1899, Jl N.Y. ent. Soc. 7: 215. LITH Type-species: Prepiella pexicera Schaus, 1899, ibidem 7:

215, by monotypy. VENEZUELA: Aroa.

The name *Prepiella* probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson as the name for a new genus.

PREPIELLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 89, 356.

Type-species: *Odozana hippona* Druce, 1855, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 1: 125, pl.12, fig.15, by original designation. GUATEMALA: Cubilguitz and Pamima.

A junior homonym of *Prepiella* Schaus, 1889. There is no objective replacement name but the type-species of *Prepiella* Hampson, 1900, is currently considered to be congeneric with the type-species of *Prepiella* Schaus, 1899. The latter is therefore available for use as a subjective replacement name.

PREPIELLA Hampson, 1900. See Prepiella Schaus, 1899.

PREPODES Herrich-Schäffer, 1855. See Lycomorpha Harris, 1839.

PRINASURA Hampson, 1903, Ann. Mag. nat. Hist. (7) 11: 349.

Type-species: *Prinasura pyrrhopsamma* Hampson, 1903, *ibidem* (7) **11**: 350, by original designation. [AUSTRALIA]: New South Wales, Sydney.

P. pyrrhopsamma is a junior subjective synonym of Pallene structa Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 543.

PALLENE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 542.

Type-species: *Pallene structa* Walker, 1854, *ibidem* 2: 543, by subsequent designation by Hampson, 1900, *Cat. Lepid. Phalaenae Br. Mus.* 2: 426. [AUSTRALIA]: Sydney.

A junior homonym of *Pallene* Dejean, 1821, *Cat. Coléoptères* 1: 87, – Insecta, Coleoptera. There is no objective replacement name but the type-species of *Pallene* Walker, 1854, is currently considered to be congeneric with the type-species of *Prinasura* Hampson, 1903. The latter is therefore available for use as a subjective replacement name.

PRIONOCENTRUM Janse, 1920, Ann. Transv. Mus. 7: 226.

Type-species: Prionocentrum oneili Janse, 1920, ibidem 7: 226, pl.10, figs 18-22, pl.11, fig.1, pl.14, fig.16, by original designation. [ZIMBABWE-]RHODESIA: Umvuma and Salisbury. P. oneili is a junior subjective synonym of Trotonotus bettoni Butler, 1898, Proc. zool. Soc. Lond. 1898: 431, pl.33, fig.2. See also Trotonotus Butler, 1898.

PRISTOCERAEA Karsch, 1895, Ent. Nachr. Berlin 21: 349.

AGARISTIDAE

Type-species: Agarista eriopis Herrich-Schäffer, [1853] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.7, fig.31; [1856]: 13; [1858]: 71, 78, by original designation.

Further information in Nye, 1975, Generic Names Moths World 1: 403, 404.

ERIOPANA Kiriakoff, 1974, in Kiriakoff & Viette, Faune Madagascar 39: 6 (key), 80.

Type-species: Agarista eriopis Herrich-Schäffer, [1853]

1850-1858, by original designation.

Eriopana is a junior objective synonym of Pristoce

Eriopana is a junior objective synonym of Pristoceraea Karsch, 1895.

Eriopana in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PROANITA Dognin, 1911, Hétérocères nouv. Am. Sud 2: 43.

Type-species: *Proanita squalida* Dognin, 1911, *ibidem* 2: 43, by original designation. FRENCH GUIANA: Maroni [river].

P. squalida is a junior subjective synonym of Anita lassa Schaus, 1905, Proc. U.S. natn. Mus. 29: 279.

PROCALYPTA Butler, 1876, J. Linn. Soc. (Zool.) 12: 411.

Type-species: *Euchromia subcyanea* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 230, by original designation. MEXICO.

PROCANTHIA Hampson, 1900, Ann. S. Afr. Mus. 2: 59.

Type-species: *Procanthia argentea* Hampson, 1900, *ibidem* 2: 59, by original designation. SOUTH AFRICA: [Cape Province] Cape Colony.

‡PROCERIDIA; misspelling. See Procridia Hampson, 1900.

PROCODECA Walker, 1855, List Specimens lepid. Insects
Colln Br. Mus. 4: 779 (key), 812, 919 (as ‡Prorodeca). LYMA
Type-species: Procodeca quadrata Walker, 1855, ibidem 4:
812 by subsequent designation by Collegette, 1935, Stylons 4:

812, by subsequent designation by Collenette, 1935, *Stylops* 4: 246. [AUSTRALIA] New Holland: New South Wales.

‡PRORODECA; Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 919.

An incorrect (of a multiple) original spelling of *Procodeca* Walker, 1855. ‡*Prorodeca* was used only in the Addenda and has not subsequently been applied to this taxon.

PROCOLAX Schaus, 1910, Ann. Mag. nat. Hist. (8) 6: 584.

Type-species: *Procolax apulana* Schaus, 1910, *ibidem* (8) **6**: 584, by monotypy. Costa Rica: Juan Vinas, Tuis.

PROCRIDIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 82, 355.

Type-species: Odozana metalica Dognin, 1897, Annls Soc. ent. Belg. 41: 243, by original designation (but cited as ‡metallica, an incorrect subsequent spelling). [ECUADOR]: Loja.

‡PROCERIDIA; Neave, 1940, Nomencl. zool. 3: 903.

An incorrect subsequent spelling of *Procridia* Hampson, 1900.

PROCRIMIMA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 84, 223.

Type-species: *Lithosia procris* Felder & Rogenhofer, 1875, *Reise öst. Fregatte Novara* (Zool.) **2** (Abt.2): pl.139, fig.12, by original designation. [COLOMBIA]: Bogota.

PROCTOCOPIS Draudt, 1915, in Seitz, Gross-Schmett. Erde 6: 132.

Type-species: *Proctocopis forficula* Draudt, 1915, *ibidem* 6: 132, pl.20, row f, by original designation. Colombia: W., Rio Aguaca valley.

The date of publication of the part of Seitz containing page 132 was cited on page 129 as 16.iv.1915 in both the German and English editions.

PROELYMIOTIS Schaus, 1901, Trans. ent. Soc. Lond. 1901: 259 (key), 273.

Type-species: Nystalea aequipars Walker, 1858, List Specimens lepid. Insects Colln Br. Mus. 15: 1742, by original designation. [Brazil]: Amazon region.

PROGONA Berg, 1882, An. Soc. cient. argent. 13: 182. LITH Type-species: Cydosia luridipennis Burmeister, 1878, Descr. phys. Répub. Argent. 5 (1): 423, by monotypy. ARGENTINA: [Cordoba] Cordova.

PROLOBOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxv, 688.

Type-species: *Procrimina schistacea* Rothschild, 1913, *Novit. zool.* 20: 224, by original designation. PERU: Carabaya, Santo Domingo.

PRONERICE Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 258 (key), 265.

Type-species: Nerice disjuncta Dognin, 1892, Naturaliste 14: 85, by original designation. [ECUADOR]: Loja.

PRONOLA Schaus, 1899, Jl N.Y. ent. Soc. 7: 214. LITH Type-species: Pronola magniplaga Schaus, 1899, ibidem 7: 214, by monotypy. [BRAZIL]: São Paulo.

The name *Pronola* probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson, 1900, as the name for a new genus.

PRONOLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 87, 294.

Type-species: *Pronola magniplaga* Schaus, 1899, by original designation.

A junior homonym of *Pronola* Schaus, 1899. The objective replacement name is *Pronola* Schaus, 1899.

‡PRONOLLA; Collins, 1962, Ann. Mag. nat. Hist. (13) 5: 124. An incorrect subsequent spelling of Pronola Schaus, 1899.

PRONOLA Hampson, 1900. See Pronola Schaus, 1899.

‡PRONOLLA; misspelling. See Pronola Schaus, 1899.

PRONYGMIA Toxopeus, 1948, Treubia 19: 474.

Type-species: Euproctis xanthopera Hampson, 1897, J. Bombay nat. Hist. Soc. 11: 295, by original designation. INDIA: [Assam], Khasis.

PROPYRIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xx, 27, 521.

Type-species: *Propyria ptychoglene* Hampson, 1898, *ibidem* 1: 521, fig.276, by original designation. MEXICO: Vera Cruz.

‡PRORODECA; misspelling. See Procodeca Walker, 1855.

PROSCHALIPHORA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xvi, 7, 454.

Type-species: *Proschaliphora citricostata* Hampson, 1901, *ibidem* 3: 454, fig.194, by original designation. [Kenya] Br. E. Africa: Athi River.

PROSICCIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxii, 581.

Type-species: *Pasteosia albescens* Rothschild, 1912, *Novit. zool.* 19: 228, by original designation. New Guinea: [Papua New Guinea], junction of Angabunga River and St Joseph's River.

PROSOPIDIA Bryk, 1953, Ark. Zool. (2) 5 (1): 246. CTEN Type-species: Hyaleucerea morosa Schaus, 1910, Ann. Mag. nat. Hist. (8) 6: 195, by original designation. Costa Rica: Juan Viñas.

PROSPHOROPLITIS Kiriakoff, 1955, Revue Zool. Bot. afr. 52: 343.

Type-species: *Prosphoroplitis aglaurus* Kiriakoff, 1955, *ibidem* **52**: 343, fig.16, pl.2, fig.16, by original designation. MALAWI: Zomba.

PROTOLITHOSIA Röber, 1925, Stettin. ent. Ztg 85: 189.

Type-species: *Protolithosia roseivenata* Röber, 1925, *ibidem* 85: 189, by original designation. New Guinea: [Irian Jaya] Deutsch-Neuguinea.

PROTOMOLIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 6, 182.

Type-species: Eucereon promathides Druce, 1894, Ann. Mag. nat. Hist. (6) 13:, 172, by original designation. MEXICO: Presidio.

PROTOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 92, 255.

Type-species: Apistosia terminalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 478, by original designation. MEXICO.

See also Pygoctenucha Grote, 1883.

PROTRACTA Gaede, 1925, Mitt. zool. Mus. Berl. 11: 251.

Type-species: *Pais illuminata* Maassen, 1890, *in* Reiss & Stübel, *Reisen in Süd-Amerika*: 98, 129, pl.4, fig.20, by monotypy. Bolivia: Sicasica.

PROXHYLE Toulgoët, 1959, Annls Soc. ent. Fr. 128: 138.

Type-species: Asura vadoni Toulgoët, 1953, Mém. Inst. scient. Madagascar (E) 4: 356, text-fig.10, pl.1, fig.10, by original designation. MADAGASCAR: N.E., Maroantsetra.

PRUMALA Schaus, 1896, Jl N.Y. ent. Soc. 4: 136. ARCT Type-species: Prumala jamaicensis Schaus, 1896, ibidem 4: 137, by monotypy. JAMAICA.

PRYTANIA Debauche, 1938, Bull. Mus. r. Hist. nat. Belg. 14
(9): 3. CTEN
Type-species: Prytania straeleni Debauche, 1938, ibidem

14 (9): 3, fig., by original designation. BRAZIL: [Minas Gerais] Minas Geraes.

PRYTERIA Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 335.

Type-species: *Pryteria costata* Möschler, 1883, *ibidem* 32 (Abh.): 335, by monotypy. Surinam: Paramaribo.

‡PSALIODES; misspelling. See Psalisodes Hampson, 1910.

PSALIS Hübner, 1823, Zuträge Samml. exot. Schmett. 2: 19.

Type-species: *Psalis securis* Hübner, 1823, *ibidem* 2: 19, pl.[51], figs 291, 292, by monotypy. Java.

PSALISODES Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 482.

Type-species: *Psalisodes atrifasciata* Hampson, 1910, *ibidem* (8) 5: 483, by original designation. [Kenya]: Athi River.

‡PSALIODES; Strand, 1912, Fauna exot. 2: 30.

An incorrect subsequent spelling of *Psalisodes* Hampson, 1910.

PSAPHARACIS Turner, 1899, Trans. R. Soc. S. Aust. 23: 14.

Type-species: *Psapharacis toxophora* Turner, 1899, *ibidem* 23: 15, by monotypy. Australia: Brisbane.

PSAPHIDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 447.

Type-species: *Psaphida resumens* Walker, 1865, *ibidem* 32: 448, by monotypy. [U.S.A.]: Georgia.

Psaphida was established in the Notodontidae; it was placed in the Noctuidae by Smith, 1893, Bull. U.S. natn. Mus. 44: 111 (but misspelt as ‡Psaphidia).

PSEGMAPHORA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 636.

Type-species: Psegmaphora tripunctata Gaede, 1930, ibidem 10: 637, pl.84 row f, by monotypy. SUMATRA: Mt Korintji.

The date of publication of the part of Seitz containing page 636 was cited on page 633 as 21.ii.1930 in the German edition; and as 17.v.1930 in the English edition.

PSEPHEA Billberg, 1820, Enumeratio Insect. Mus. G.J. Bill-

Type-species: Noctua caricae Fabricius, 1775, Syst. Ent.: 596, by original designation. [ASIA]: [S.E., tropical] Indiae orientalis.

See also Euimata Billberg, 1820.

DAMALIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 172.

Type-species: Noctua caricae Fabricius, 1775, by subsequent designation by Jordan, 1896, Novit. zool. 3: 203.

A junior homonym Damalis Fabricius, 1805, Syst. Antliatorum: 147, - Insecta, Diptera. The objective replacement name is Psephea Billberg, 1820.

AGANAIS Boisduval, 1832, in d'Urville, Voyage Découvertes de l'Astrolabe (Faune ent.) 1: 248.

Type-species: *Noctua caricae* Fabricius, 1775, by monotypy. A junior objective synonym of Psephea Billberg, 1820.

PSEUDACLYTIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 409.

Type-species: Pampa opponens Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 100, by original . designation. [BRAZIL]: [R. Amazon], [Tefé] Ega.

PSEUDAETHRIA Schaus, 1924, Proc. U.S. natn. Mus. 65 (7):

Type-species: Pseudaethria cessogae Schaus, 1924, ibidem 65 (7): 11, by original designation. BRAZIL: Joinville.

PSEUDAGOMA Kiriakoff, 1975, Bull. Annls Soc. r. ent. Belg. 111: 199.

Type-species: Pseudagoma pinheyi Kiriakoff, 1975, ibidem 111: 200, fig.2, by original designation. Holotype &, Moz-AMBIQUE: Amatonga forest, (NMR, Bulawayo).

Pseudagoma in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PSEUDALLATA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 244.

Type-species: Spatalia laticostalis Hampson, 1900, J. Bombay nat. Hist. Soc. 13: 43, pl.B, fig.15, by original designation. INDIA: Khasia [Hills].

PSEUDALUS Schaus, 1896, Jl N.Y. ent. Soc. 4: 135. Type-species: Pseudalus limona Schaus, 1896, ibidem 4: 135, by subsequent designation by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 253, (but cited as ‡limonia, an incorrect subsequent spelling by Hampson). BRAZIL: São Paulo.

PSEUDANTIORA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 566.

Type-species: Antiora contingata Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 347, pl.18, fig.35, by monotypy (of Antiora Möschler, 1883). Surinam: interior.

Pseudantiora was established as the objective replacement name for Antiora Möschler, 1883.

ANTIORA Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 346.

Type-species: Antiora contingata Möschler, 1883, by monotypy.

A junior homonym of Antiora Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1769, - Lepid., Notodontidae. The objective replacement name is Pseudantiora Kirby,

‡PSEUDAPICOMA: misspelling. See Pseudapiconoma Aurivillius, 1881. THYRET

PSEUDAPICONOMA Aurivillius, 1881, Ent. Tidskr. 2: 46.

Type-species: Pseudapiconoma testacea Aurivillius, 1881, ibidem 2: 46, fig.1, by monotypy. [GABON] Gaboon.

Pseudapiconoma was established in the Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson,

1898, Cat. Lepid. Phalaenae Br. Mus. 1: 149; and was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 10.

‡PSEUDAPICOMA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling of *Pseudapiconoma* Aurivillius, 1881.

PSEUDAPISTOSIA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 665.

Type-species: Phalaena umber Cramer, 1775, Uitlandsche Kapellen (Papillons exot.) 1: 24, 155, pl.15, fig.F, by monotypy. SURINAM.

P. umber is dated from the wrapper of the part and not from the title-page of the volume, 1779.

‡PSEUDOPISTOSIA; Dognin, 1894, Annls Soc. ent. Belg. 38: 682.

An incorrect subsequent spelling of Pseudapistosia Möschler, 1878.

PSEUDARCTIA Bethune-Baker, 1911, Ann. Mag. nat. Hist.

Type-species: Pseudarctia nivea Bethune-Baker, 1911, ibidem (8) 7: 540, by original designation. UGANDA: [Entebbe] N'tebi.

Pseudarctia was established in the Lymantriadae, now Lymantriidae; it was transferred to the Arctiidae by Collenette, 1955, Trans. R. ent. Soc. Lond. 107: 189.

PSEUDARGYROEIDES Klages, 1906, Proc. U.S. natn. Mus.

Type-species: Pseudargyroeides caurensis Klages, 1906, ibidem 29: 547, by original designation. VENEZUELA: S., Suapure.

PSEUDARGYROIDES Zerny, 1912, in Wagner, Lepid. Cat. **7**: 110.

An unjustified emendation of *Pseudargyroeides* Klages, 1906.

PSEUDARGYROIDES Zerny, 1912. See Pseudargyroeides Klages, 1906. **CTEN**

‡PSEUDEBESSA; misspelling. See Pseuderbessa Butler, 1876.

PSEUDERBESSA Butler, 1876, Cistula ent. 2: 107. Type-species: Dioptis umbrifera Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 326, by original designation. [BRAZIL]: Para.

‡PSEUDEBESSA; Prout, 1918, Novit. zool. 25: 401.

An incorrect subsequent spelling of Pseuderbessa Butler, 1876.

PSEUDEUCEREON Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 652.

Type-species: Phalaena eleuthera Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 159, pl.371, fig.A, by monotypy. SURINAM.

P. eleuthera is dated from the wrapper of the part and not from the title-page of the volume, 1782.

PSEUDHAPIGIA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 264 (key), 342.

Type-species: *Pseudhapigia brunnea* Schaus, 1901, *ibidem* 1901: 343, by original designation. MEXICO: Guadalajara.

PSEUDHEMICERAS Dognin, 1914, Hétérocères nouv. Am. Sud 7: 29.

Type-species: Pseudhemiceras lepidoides Dognin, 1914, ibidem 7: 29, by original designation. PERU: Yahuarmayo.

SCHAUSIADES Dognin, 1914, Hétérocères nouv. Am. Sud 8: 93.

Proposed unnecessarily as an objective replacement name for *Pseudhemiceras* Dognin, 1914, in the mistaken belief that it was preoccupied by *Pseudohemiceras* Möschler, 1890.

PSEUDHYPSA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 384.

Type-species: *Phalaena speciosa* Drury, 1773, *Illust. nat. Hist. exot. Insects* 2: 8, pl.5, fig.2, by subsequent designation by Hampson, 1905, *Ann. S. Afr. Mus.* 3: 413. SIERRA LEONE.

PSEUDILEMA Bethune-Baker, 1904, Novit. zool. 11: 418.

Type-species: *Pseudilema dinawa* Bethune-Baker, 1904, *ibidem* 11: 419, pl.5, figs 17, 18, by original designation. New GUINEA: [Papua New Guinea], Aroa R. and Dinawa.

PSEUDISCHNOCAMPA Rothschild, 1935, Novit. zool. 39: 243.

Type-species: *Ischnocampa nigrivena* Schaus, 1901, *Ann. Mag. nat. Hist.* (7) 7: 268, by original designation. [BOLIVIA]: Suapi.

PSEUDLEPISTA Hampson, 1910, Proc. zool. Soc. Lond. 1910: 392.

Type-species: *Pseudlepista atrizona* Hampson, 1910, *ibidem* **1910**: 392, pl.36, fig.7, by original designation. [ZAMBIA] N.E. Rhodesia: [Serenje] Serenji district.

PSEUDMELISA Hampson, 1910, Proc. zool. Soc. Lond. 1910: 391.

Type-species: *Pseudmelisa chalybsa* Hampson, 1910, *ibidem* **1910**: 391, pl.36, fig.5, by original designation. [ZAIRE] Congo: [Shaba], Kambove district.

Pseudmelisa was established in the Syntomidae, now Ctenuchidae; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 21.

PSEUDOMELISA Zerny, 1912, in Wagner, Lepid. Cat. 7: 47. An unjustified emendation of Pseudmelisa Hampson, 1910.

PSEUDOBAROBATA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 438.

Type-species: *Pseudobarobata angulata* Gaede, 1928, *ibidem* 14: 438, pl.72 row d, by monotypy. CAMEROUN, RWANDA.

The date of publication of the part of Seitz containing page 438 was cited on page 433 as 30.v.1928 in the German edition; and as 26.xi.1928 in the English edition.

PSEUDOBAZISA Bryk, 1934, in Strand, Lepid. Cat. 62: 110.

Type-species: Aroa perculta Distant, 1897, Ann. Mag. nat. Hist. (6) 20: 201, by original designation. [South Africa]: Transvaal, Pretoria.

Pseudobazisa was proposed as a 'nom. nov.' for 'Bazisa Janse (nec Walker)'. Bryk was not, however, proposing a replacement name for a homonym but was proposing a new genus for Bazisa Walker sensu Janse, 1915, Ann. Transv. Mus. 5: 7.

PSEUDOBLABES Zeller, 1853, Bull. Soc. imp. Nat. Moscou 26 (4): 512.

Type-species: *Pseudoblabes oophora* Zeller, 1853, *ibidem* **26** (4): 514, pl.4, fig.1, by monotypy. JAVA.

PSEUDOCALLIMORPHA Oberthür, 1879, Annali Mus. civ. Stor. nat. Giacomo Doria 15: 175.

Type-species: *Pseudocallimorpha doriae* Oberthür, 1879, *ibidem* **15**: 175, pl.1, fig.7, by monotypy. [Етнюріа] Abyssinia: Scioa, Mahal Uonz.

PSEUDOCERYX Rothschild, 1910, Novit. zool. 17: 440.

Type-species: *Pseudoceryx dohertyi* Rothschild, 1910, *ibidem* 17: 444, by monotypy. New Guinea: [Irian Jaya], Geelvink Bay, Biak Island.

PSEUDOCHARIDEA Druce, 1896, Ann. Mag. nat. Hist. (6) 18: 35.

Type-species: *Pseudocharidea eion* Druce, 1896, *ibidem* (6) **18**: 36, by original designation. PANAMA: Chiriqui.

PSEUDOCHARIS Druce, 1884, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 56.

Type-species: *Pseudocharis naenia* Druce, 1884, *ibidem* 1: 57, pl.7, fig.12, by original designation. GUATEMALA: San Geronimo.

PSEUDOCOARICA Holloway, 1979, Series Ent. 15: 516.

Type-species: *Pseudocoarica caledonica* Holloway, 1979, *ibidem* **15**: 516, text-figs 131, 141, 142, pl.85, figs 13, 14, by original designation. Holotype &, New Caledonia: Mt Aoupinié, (BMNH).

Pseudocoarica in the Noctuidae, Ophiderinae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PSEUDODIPTERA Kaye, 1918, Ann. Mag. nat. Hist. (9) 2: 229.

Type-species: *Pseudodiptera musiforme* Kaye, 1918, *ibidem* (9) 2: 230, by original designation. [CHAD] Congo: Oubanguichari, Tschad.

Pseudodiptera was established in the Syntomidae, now Ctenuchidae; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 48.

PSEUDODREATA Bethune-Baker, 1904, ibidem 11: 371.

ANTHELIDAE

Type-species: *Pseudodreata strigata* Bethune-Baker, 1904, *ibidem* 11: 371, pl.5, fig.45, by original designation. New Guinea: Papua New Guinea, Angabunga River, Dinawa.

Pseudodreata was established in the Eupterotidae; it was transferred to the Lymantriidae by Joicey, Noakes & Talbot, 1915, Trans. ent. Soc. Lond. 1915: 379. Swinhoe, 1922, Ann. Mag. nat. Hist. (9) 10: 451, placed Pseudodreata in the Lymantriidae, subfamily Anthelinae, which had been raised to family level by Turner, 1920, Trans. ent. Soc. Lond. 1919: 415.

CYCETHRA Bethune-Baker, 1904, Novit. zool. 11: 393.

Type-species: Cycethra aroa Bethune-Baker, 1904, ibidem 11: 393, by original designation. New Guinea: Papua New Guinea, Aroa River.

A junior homonym of Cycethra Bell, 1881, Proc. zool. Soc. Lond. 1881: 96, — Echinoderma. There is no objective replacement name, but C. aroa is congeneric with Pseudodreata strigata Bethune-Baker, 1904, the type-species of Pseudodreata Bethune-Baker, 1904; the latter is therefore available for use as a subjective replacement name.

Cycethra was established in the Lasiocampidae. It was transferred to the Lymantriidae by Joicey, Noakes & Talbot, 1915, Trans. ent. Soc. Lond. 1915: 379. Swinhoe, 1922, Ann. Mag. nat. Hist. (9) 10: 451, placed Cycethra in the Lymantri-

idae, subfamily Anthelinae which had been raised to family level by Turner, 1920, Trans. ent. Soc. Lond. 1919: 415.

PSEUDODRYAS Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 685.

Type-species: Pseudodryas olivacea Möschler, 1878, ibidem **27** (Abh.): 685, pl.10, fig.43, by monotypy. Surinam.

PSEUDODURA Strand, 1914, Ent. Mitt. 3: 332. LYMA Type-species: Pseudodura dasychiroides Strand, 1914, ibideni 3: 333, by original designation. [Taiwan] Formosa.

PSEUDOEDALEOSIA Strand, 1924, Dt. ent. Z. Iris 38: 137.

Type-species: Pseudoedaleosia scoparioides Strand, 1924, ibidem 38: 137, by original designation. AUSTRALIA: Western Australia, [Shark's Bay], Dirk Hartog [I.], Brown Station.

PSEUDOFENTONIA Strand, 1912, Fauna exot. 2: 40. NOTO Type-species: Heterocampa argentifera Moore, 1866, Proc. zool. Soc. Lond. 1865: 813, by original designation. [INDIA]: Darjeeling.

PSEUDOFENTONIA Kiriakoff, 1964. See Antifentonia Kiriakoff, 1967.

PSEUDOGALTARA Toulgoët, 1979, Bull. Soc. ent. Fr. 83: 221.

Type-species: Pseudogaltara inexpectata Toulgoët, 1979, ibidem 83: 221, by original designation. RWANDA: Nyungwe Forest, Route Delvaux, km 16.

PSEUDOGARGETTA Bethune-Baker, 1904, Novit. zool. 11: 373.

Type-species: Pseudogargetta diversa Bethune-Baker, 1904, ibidem 11: 373, pl.5, fig.4, by original designation. New GUINEA: [Papua New Guinea], Dinawa, Ekeikei.

PSEUDOGENUSA Rebel, 1914, Annln K. K. naturh. Hofmus. Wien 28: 285.

Type-species: Pseudogenusa opalescens Rebel, 1914, ibidem 28: 285, pl.24, fig.72, by monotypy. [TANZANIA]: [Tanganyika] Deutsch-Ostafrika, [between] Bukoba and [Bujumbura] Usumbura, Ukaika.

P. opalescens is a junior subjective synonym of Olapa makala Bethune-Baker, 1909, Ann. Mag. nat. Hist. (8) 3: 428.

Pseudogenusa was established in the Geometridae; it was transferred to the Lymantriidae by Collenette, 1931, Ann. Mag. nat. Hist. (10) 7: 510.

PSEUDOHEMIHYALEA Rego Barros, 1956, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 134: 33.

Type-species: Phaegoptera rhoda Druce, 1894, Ann. Mag. nat. Hist. (6) 13: 173, by original designation. GUATEMALA: [Guatemala City] in the city.

PSEUDOHOPLITIS Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 640.

Type-species: Pseudohoplitis vernalis Gaede, 1930, ibidem 10: 640, pl.84 row h, by original designation. [Sulawesi] Celebes: northern.

The date of publication of the part of Seitz containing page 640 was cited on page 633 as 21.ii.1930 in the German edition; and as 17.v.1930 in the English edition.

‡PSEUDOHYALEUCEREA; Rego Barros & Machado, 1971. See Pseudohyaleucerea Rego Barros & Machado nom. **CTEN** n.

PSEUDOHYALEUCEREA Rego Barros & Machado gen. n.

Type-species: Hyaleucerea vulnerata Butler, 1875, Ann. Mag.

nat. Hist. (4) 15: 399, by PRESENT DESIGNATION. [BRA-

ZIL]: [Espirito Santo] Espiritu Santo.

Pseudohyaleucerea Rego Barros & Machado is here established with the agreement of the authors, for the concept ‡Pseudohyaleucerea Rego Barros Machado, 1971, Arg. Mus. nac. Rio de J. 54: 173. When the name was originally proposed in 1971 for two species, it was accompanied by a statement 'that purports to give characters differentiating the taxon' (Article 13(a)(i)) but was not also 'accompanied by the definite fixation of a type-species' (Article 13(b)). Pseudohyaleucerea was therefore unavailable when originally proposed, but is here established.

‡PSEUDOHYALEUCEREA; Rego Barros & Machado, 1971, Arg. Mus. nac. Rio de J. 54: 173.

Pseudohyaleucerea was proposed for two included nominal species but was not originally accompanied by the fixation of a type-species and is therefore unavailable under the Code, Article 13(b).

PSEUDOHYPSOIDES Viette, 1960, Bull. mens. Soc. linn. Lyon 29: 71.

Type-species: Pseudohypsoides bicolor Viette, 1960, ibidem 29: 71, by original designation. MADAGASCAR: southern, Sakaraka district.

PSEUDOLABIS Hartig, 1963. See Uliolepis Warren, 1897. LYMA

PSEUDOMACROPTILA Fleming, 1951, Zoologica, N.Y. 36: 183.

Type-species: Pseudomacroptila argentea Fleming, 1951, ibidem 36: 183, fig., by original designation. VENEZUELA: Aragua, Rancho Grande.

PSEUDOMELISA Zerny, 1912. See Pseudmelisa Hampson, 1910. THYRET

PSEUDOMESA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 923.

Type-species: Pseudomesa quadriplagiata Walker, 1855, ibidem 4: 923, by monotypy. [BANGLADESH]: [Sylhet], Silhet.

PSEUDOMYA Hübner, [1819] 1816, Verz. bekannter Schmett.: 124.

Type-species: Glaucopis tipulina Hübner, [1812], Samml. exot. Schmett. 1: pl.[163], by subsequent designation by Butler, 1876, J. Linn. Soc. (Zool.) 12: 385. TYPE-LOCALITY: not stated, [South America]: [tropical].

‡PSEUDONACHA; misspelling. See Pseudonaclia Butler, 1876.

PSEUDONACLIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 351.

Type-species: Naclia puella Boisduval, 1847, in Delegorgue, Voyage Afr. australe 2: 596, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 106. TYPE-LOCALITY: not stated, [SOUTH AFRICA].

‡PSEUDONACHA; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 106.

An incorrect subsequent spelling of Pseudonaclia Butler, 1876.

PSEUDONERICE Bryk, 1950, Ark. Zool. 42 (A) 19: 40.

Type-species: Pseudonerice unidentata Bryk, 1950, ibidem 42 (A) 19: 40, pl.3, fig.10, by original designation. Burma: Kambaiti.

PSEUDONOTODONTA Möschler, 1887, Abh. senckenb. naturforsch. Ges. 15: 76. LYMA

Type-species: Pseudonotodonta virescens Möschler, 1887, ibidem 15: 77, fig.6, by monotypy. [GHANA] Goldküste: Aburi.

PSEUDOPHAIO Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xiii, 216.

Type-species: Poliopasta rosenbergi Rothschild, 1911, Novit. zool. 18: 40, by original designation. ECUADOR: W., Paramba.

PSEUDOPHALOE Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 429 (but cited as Pseudophaloë).

Type-species: Phaloe verania Druce, 1884, Biologia cent.am. (Zool.) Lepid.-Heterocera 1: 107, pl.11, fig.7, by original designation. GUATEMALA: Zapote.

The date of publication of the part of Seitz containing page 429 was cited on page 425 as 12.i.1925 in the German edition and as 4.iii.1925 in the English edition.

PSEUDOPHANES Turner, 1940, Proc. R. Soc. Qd 51: 104.

Type-species: Pseudophanes melanoptera Turner, 1940, ibidem 51: 105, by monotypy. AUSTRALIA: N. Queensland, Kuranda.

PSEUDOPHARUS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: ix, 3, 133.

Type-species: Opharus amata Druce, 1900, Ann. Mag. nat. Hist. (7) 6: 67, by original designation. VENEZUELA.

‡PSEUDOPISTOSIA; misspelling. See Pseudapistosia Möschler, 1878. ARCT

PSEUDOPOMPILIA Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 401.

Type-species: Pseudopompilia mimica Druce, 1898, ibidem

(7) 1: 401, by monotypy. [Brazil]: Amazon region.

The name Pseudopompilia probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson in 1898 [December] as the name for a new genus.

PSEUDOPOMPILIA Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xviii, 27, 425.

Type-species: Pseudopompilia mimica Druce, 1898, by orig-

inal designation.

A junior homonym of Pseudopompilia Druce, 1898 May. The objective replacement name is Pseudopompilia Druce, 1898 May.

PSEUDOPOMPILIA Hampson, 1898 [December]. See Pseudopompilia Druce, 1898 May.

PSEUDORETHONA Janse, 1920, Ann. Transv. Mus. 7: 168.

Type-species: Rethona albicans Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1043, by original designation. South Africa.

PSEUDORICIA Prout, 1918, Novit. zool. 25: 396 (key), 406.

Type-species: Josiodes sibyllae Druce, 1885, Proc. zool. Soc. Lond. 1885: 523, pl.32, fig.7, by original designation. ECUA-

In the original combination of the type-species, the generic name was published as ‡Josioides, an incorrect subsequent spelling.

PSEUDOSCAPTIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xix, 477.

Type-species: Scaptesyle bicolor Rothschild, 1912, Novit. zool. 19: 241 (but cited as ‡Scaptosyle, an incorrect subsequent spelling by Rothschild), by original designation (but cited as ‡Scaptosyle). [MALAYSIA]: Malay Peninsula, Padang Rengas. PSEUDOSCOLIA Hampson, 1914. See Parascolia Watson

PSEUDOSCRANCIA Strand, 1915, Ent. Mitt. 4: 174. NOTO Type-species: Maguza africana Holland, 1893, Psyche, Camb. 6: 565, fig., by monotypy. [GABON]: [Ogooué] Ogové

Pseudoscrancia was established as a subgenus of Scrancia; it was raised to generic level by Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 423.

PSEUDOSPHECOSOMA Strand, 1915, Dt. ent. Z. 1: 19.

Type-species: Pseudosphecosoma vespoides Strand, 1915, ibidem 1: 20, by original designation. Brazil: Maranhão.

PSEUDOSPHENOPTERA Butler, 1876, J. Linn. Soc. (Zool.) 12: 370.

Type-species: Euchromia basalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 251, by original designation. [Brazil]: Amazon R., [Tefé] Ega.

See also Chloropsinus Butler, 1876, and Enioche Druce, 1896.

PSEUDOSPHEX Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 12, 13.

Type-species: Pseudosphex zethus Hübner, 1818, ibidem 1: 13, pl.[9], figs 49, 50, by subsequent designation by Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1618.

‡SPHECOMORPHA Hübner, 1808, Erste Zuträge Samml. exot. Schmett.: 4.

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1966, Bull. zool. Nom. 23 (Opinion 789): 213. Placed on Official Index of Rejected and Invalid Generic Names in Zoology: Name No.1884. Two included species ‡Sphecomorpha polistes Hübner, 1808, ibidem: 4, later published and made available as Pseudosphex polistes Hübner, 1818, Zuträge Samml. exot. Schmett.1: 12 and ‡Sphecomorpha zethus Hübner, 1808, ibidem: 4, later published and made available as Pseudosphex zethus Hübner, 1818, ibidem 1: 13 (the typespecies of Pseudosphex Hübner, 1818).

‡SPHECOMORPHA Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 12, 13, 34.

Not nomenclaturally available as a genus-group name. Hübner used Sphecomorpha, on all three pages, in a suprageneric sense for one of the names of his tribes (Namen der Stämme), listed on pages [33] and [34]. Hübner listed his genus-group names on pages [35] and [36].

ABROCHIA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 101, 103.

Type-species: Pseudosphex zethus Hübner, 1818, by monotypy (but cited as ‡zethes, an incorrect subsequent spelling by Herrich-Schäffer).

Abrochia is a junior objective synonym of Pseudosphex Hübner, 1818.

Abrochia was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some

important Publications.

PSEUDOSTAUROPUS Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 631.

Type-species: *Pseudostauropus plagosus* Gaede, 1930, *ibidem* 10: 631, pl.84 row a, by monotypy. [MALAYSIA]: Penang, SUMATRA: Barisan Mts.

The date of publication of the part of Seitz containing page 631 was cited on page 625 as 11.iii.1930 in both the German and the English editions.

PSEUDOSTRACILLA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 124.

Type-species: *Pseudostracilla infausta* Hering, 1926, *ibidem* **14**: 124, pl.19, row b, by monotypy. SIERRA LEONE.

The date of publication of the part of Seitz containing page 124 was cited on page 121 as 9.x.1926 in both the German and the English editions.

Pseudostracilla was established in the Pterothysanidae; it was placed in the Lymantriidae by Collenette, 1955, Trans. R. ent. Soc. Lond. 107: 190.

PSEUDOTELECLITA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 219.

Type-species: *Pheosia centristicta* Hampson, 1898, *J. Bombay nat. Hist. Soc.* 11: 282, by original designation. [INDIA]: Sikkim.

PSEUDOTESSELLARCTIA Travassos, 1949, Revta bras. Biol. 9: 449. ARCT

Type-species: *Phaegoptera ursina* Schaus, 1892 August, *Proc. zool. Soc. Lond.* **1892**: 281, by original designation. BRAZIL: Rio de Janeiro.

P. ursina was redescribed by Schaus, 1892 November, Am. Lepid. 12: pl.3, fig.15.

PSEUDOTHYRETES Dufrane, 1945, Bull. Annls Soc. r. ent. Belg. 81: 127.

Type-species: *Pseudothyretes mariae* Dufrane, 1945, *ibidem* 81: 127, by original designation. [ZAIRE]: Kivu [province], Kamituga.

Pseudothyretes when established was not placed in a family; it was placed in the Thyretidae by Kiriakoff, 1960, in Wytsman, Genera Insect. 214 (E): 47.

PSEUDOTUERTA Kiriakoff, 1976, Tierreich 98: 113.

Type-species: *Tuerta argyrochlora* Carcasson, 1964, *Jl E. Afr. nat. Hist. Soc.* **24**: 72, figs 30, 59, by original designation. Holotype & UGANDA: Ankole, Kalinzu Forest, (BMNH).

Pseudotuerta in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PSEUDOTURNACA Kiriakoff, 1967, Tijdschr. Ent. 110: 41.

Type-species: *Pseudoturnaca samarinda* Kiriakoff, 1967, *ibidem* 110: 42, fig.5, by original designation. Borneo: southeastern, Samarinda.

PSICHOTOE Boisduval, 1828, Essai Monogr. Zygénides: 129.

Type-species: *Psichotoe duvaucelii* Boisduval, 1828, *ibidem*: 129, pl.8, figs 5, 6, by monotypy. [INDIA]: [Bengal] Bengale.

PSYCHOTHOE Zerny, 1912, in Wagner, Lepid. Cat. 7: 4. An unjustified emendation of Psichotoe Boisduval, 1828.

PSILACRON Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.22. NOTO

Type-species: *Psilacron luteovirens* Felder, 1874, *ibidem* 2 (Abt.2): pl.97, fig.22, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 8.

‡PSILAGON; Druce, 1911, Ann. Mag. nat. Hist. (8) 8: 144.
An incorrect subsequent spelling of Psilacron Felder, 1874.

‡PSILAGON; misspelling. See Psilacron Felder, 1874.

PSILOCHIRA Toxopeus, 1948, Treubia 19: 447.

Type-species: Lymantria lineata Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 874 (key), 875, by original designation. [BANGLADESH]: [E. Bengal], [Sylhet] Silhet.

PSILOPEPLA Turner, 1899, Trans. Proc. R. Soc. S. Aust. 23: 13.

Type-species: *Nudaria mollis* Lucas, 1894, *Proc. Linn. Soc. N.S.W.* (2) 8: 136, by monotypy. AUSTRALIA: near Brisbane.

PSILOPLEURA Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 407.

Type-species: *Psilopleura polia* Druce, 1898, *ibidem* (7) 1: 407, by monotypy. [BRAZIL]: [Espirito Santo] Espiritu Santo.

The name *Psilopleura* probably originated from Hampson but was used and made nomenclaturally available by Druce prior to its proposal by Hampson in 1898 December as the name for a new genus.

PSILOPLEURA Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xiv, 26, 267.

Type-species: *Pheia vittata* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 83, by original designation. [MEXICO]: Vera Cruz.

A junior homonym of *Psilopleura* Druce, 1898 May. There is no objective replacement name but the type-species of *Psilopleura* Hampson is currently considered to be congeneric with the type-species of *Psilopleura* Druce. The latter is therefore available for use as a subjective replacement name.

PSILOPLEURA Hampson, 1898 [December]. See Psilopleura Druce, 1898 May. CTEN

PSILURA Stephens, 1828. See Lymantria Hübner, [1819].

PSOLOPTERA Butler, 1876, J. Linn. Soc. (Zool.) 12: 369.

Type-species: *Euchromia thoracica* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 243, by original designation. [Brazil]: Amazon River, [Tefé] Ega.

PSOROCAMPA Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 260 (key), 286.

Type-species: *Psorocampa denticulata* Schaus, 1901, *ibidem* **1901**: 286, pl.11, fig.5, by original designation. [Brazil]: Parana, Castro.

PSYCHARIUM Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.80, fig.461; [1858], ibidem 1: 76. SOMABRACHYIDAE

Type-species: *Psycharium pellucens* Herrich-Schäffer, [1856] 1850-1858, *ibidem* 1 (1): wrapper, pl.80, fig.461; [1858], *ibidem* 1: 76, by monotypy. [SOUTH AFRICA] 'Afr.?'.

Psycharium when established was not placed in a family; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 498; was transferred to the Megalopygidae by Aurivillius, 1894, Dt. ent. Z. Iris 7: 189; was placed in the Arctiadae, now Arctiidae, by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 451; was transferred to the Zygaenidae by Jordan, 1928, Novit. zool. 34: 136; and was transferred to the Somabrachyidae by Tremewan, 1973, Bull. Br. Mus. nat. Hist. (Ent.) 28: 139.

PSYCHOPHASMA Butler, 1878, Trans. ent. Soc. Lond. 1878: 51.

Type-species: Halysidota albidator Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 313 (but cited as ‡Halesidota, an incorrect subsequent spelling by Walker),

by original designation (but cited as ‡Halesidota). [BRAZIL]: [Amazon River], [Tefé] Ega.

H. albidator is a junior subjective synonym of Halysidota erosa Herrich-Schäffer, [1858] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): 84, pl. [96], fig. 550, (but cited as ‡Halesidota, an incorrect subsequent spelling).

PSYCHOTHOE Zerny, 1912. See Psichotoe Boisduval, 1828.

PTEREDOA Hampson, 1905, Ann. S. Afr. Mus. 3: 411.

Type-species: Pteredoa plumosa Hampson, 1905, ibidem 3: 412, by original designation. [RHODESIA]: Mashonaland, Salisbury.

PTEROMA Staudinger, 1896. See Pterotes Berg, 1901. NOTO

PTEROODES Butler, 1877, Trans. ent. Soc. Lond. 1877: 376.

Type-species: Lithosia longipennis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 510, by original designation. MEXICO.

PTEROPHAEA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 51.

Type-species: Metarctia brunneipennis Hering, 1932, Revue Zool. Bot. afr. 22: 108, pl.2, fig.19, by original designation. [ZAIRE]: [Katanga], [Lubumbashi] Elisabethville.

Pterophaea was established as a subgenus of Metarctia Walker, 1855.

PTEROSTOMA Germar, 1812, Diss. sistens Bombycum Species (2): 42.

Type-species: Pterostoma salicis Germar, 1812, ibidem (2): 43, by monotypy. Type-locality: not stated, [Europe].

P. salicis is a junior subjective synonym of Phalaena palpina Clerck, 1759, Icon. Insect. rariorum 1: pl.9, fig.8.

See also Euchila Billberg, 1820.

PTEROTES Berg, 1901, Comun. Mus. nac. B. Aires 1: 311.

Type-species: Pteroma eugenia Staudinger, 1896, Dt. ent. Z. Iris 8: pl.5, fig.13; 1900, ibidem 12: 352, by monotypy (of Pteroma Staudinger, 1896). Mongolia: Uliastay.

Pterotes was established as the objective replacement name for Pteroma Staudinger, 1896.

PTEROMA Staudinger, 1896, Dt. ent. Z. Iris 8: pl.5, fig.13; 1900, ibidem 12: 352.

Type-species: Pteroma eugenia Staudinger, 1896, by monotypy.

A junior homonym of Pteroma Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 65, - Lepid., Psychidae. The objective replacement name is Pterotes Berg, 1901.

DANIELITA Kiriakoff, 1970, NachrBl. bayer. Ent. 19: 124.

Proposed unnecessarily as the objective replacement name for Pteroma Staudinger, 1896; Pterotes Berg, 1901, had already been established.

PTEROTHYSANUS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 401. PTEROTHYSANIDAE

Type-species: Pterothysanus laticilia Walker, 1854, ibidem 2: 401, by monotypy. [BANGLADESH]: [E. Bengal], [Sylhet] Silhet, [INDONESIA] East Indies.

Pterothysanus was established in the Lithosiidae, now Arctiidae; it was made the type-genus of the family Pterothysanidae Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 430. Bryk, 1934, in Strand, Lepid. Cat. 62: 351, treated this group as a subfamily of the Lymantriidae.

PTERYGOPTERUS Butler, 1876. See Timalus Watson nom.

PTERYGOSOMA Felder, 1874. See Oenosandra Walker,

‡PTILODON Hübner, [1806]. See Ptilodon Hübner, [1822].

‡PTILODON Hübner, 1808. See Gonodonta Hübner, 1818. NOCTUIDAE

PTILODON Hübner, 1822, Syst.-alphab. Verz.: 14-16, 18, 20.

Type-species: Phalaena camelina Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 507, by subsequent designation by Grote, 1895, Abh. naturw. Ver. Bremen 14: 48. Type-locality: not stated, [EUROPE].

P. camelina is a junior subjective synonym of Phalaena capucina Linnaeus, 1758, ibidem 1: 507.

‡PTILODON Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) Opinion 97: 19. Also idem, 1954, Opin. Decl. int. Commn zool. Nom. 6 Opinion 278:

Only included species: Phalaena camelina Linnaeus, 1758.

LOPHOPTERYX Stephens, 1828, Illust. Br. Ent. (Haustellata) **2**: 26.

Type-species: Phalaena camelina Linnaeus, 1758, by subsequent designation by Westwood, 1840, Introd. mod. Classif. Insects 2 (Synopsis Genera Br. Insects): 90.

A junior objective synonym of Ptilodon Hübner, 1822.

Invalid type-species designation: Phalaena camelina Linnaeus, 1758, was designated by Boisduval, 1836, Hist. nat. Insectes (Spec. gén. Lépid.) 1: 142.

In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different type-species for each generic name. In his 'Exposé de notre méthode', pages 155-690, no typespecies designation was made for any of the genera that he himself used. Under the Code, Article 69(a)(iii), the typespecies designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other authors. The acceptance of Boisduval's type-species designation in the case of Lophopteryx would make no difference to the generic concept.

Lophopteryx was again proposed by Stephens, 1829 [June], Nom. Br. Insects: 39; and 1829 [July], Syst. Cat. Br. Insects (2): 43.

PTILODONTELLA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 176.

Type-species: Bombyx cucullina [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 311, by original designation. [AUSTRIA]: Vienna district.

PTILODONTIS Stephens, 1828. See Euchila Billberg, 1820. NOTO

PTILODONTOSIA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 232.

Type-species: Lophopteryx crenulata Hampson, 1896, Fauna Br. India (Moths) 4: 460, by original designation. [INDIA]: Sikkim, Yatong.

PTILOMACRA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1098.

Type-species: Ptilomacra senex Walker, 1855, ibidem 5: 1098, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 862. Australia: New South Wales.

Ptilomacra was established in the Notodontidae; it was placed in the Zeuzeridae, now Cossidae, by Kirby, 1892, ibidem 1: 862.

‡PHILOMACRA; Moore, [1883] 1882-3, Lepid. Ceylon 2: 144. An incorrect subsequent spelling of Ptilomacra Walker, 1855.

‡PTILOPHOIDES; misspelling. See Ptilophoroides Matsumura, 1920.

PTILOPHORA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 29.

Type-species: *Phalaena variegata* Villers, 1789, *Caroli Linn. Ent.* 2: 160 (a junior primary homonym, see below), by monotypy. France: southern.

Phalaena variegata Villers is a junior primary homonym of Phalaena variegata Scopoli, 1763, Ent. Carniolica: 222, — Lepid., Geometridae. There is no objective replacement name, but P. variegata Villers is a junior subjective synonym of Bombyx plumigera [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 61; the latter is thus available for use as a subjective replacement name.

Ptilophora was again proposed by Stephens, 1829 [June], Nom. Br. Insects. 39; and 1829 [July], Syst. Cat. Br. Insects (2): 44.

PTILOPHOROIDES Matsumura, 1920, Zool. Mag. Tokyo 32: 149.

Type-species: *Ptilophoroides jezoensis* Matsumura, 1920, *ibidem* 32: 150, by original designation. JAPAN: [Sapporo], Jozankei.

Matsumura, 1920, *ibidem* 32: 150, spelt the generic name as ‡*Ptilophoides*, an incorrect (of a multiple) original spelling.

Ptilophoroides was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 821-823; on page 822 the name was incorrectly spelt as ‡Ptitophoroides.

‡PTILOPHOIDES; Matsumura, 1920, Zool. Mag. Tokyo 32: 150.

An incorrect (of a multiple) original spelling of *Ptilophoroides* Matsumura, 1920.

‡PTITOPHOROIDES; Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 822.

An incorrect subsequent spelling of *Ptilophoroides* Matsumura, 1920.

PTILURA Holland, 1893, Ent. News 4: 103. NOTO

Type-species: *Ptilura argyraspis* Holland, 1893, *ibidem* 4: 104, pl.7, fig.1, by original designation. [GABON]: [Ogooué] Ogové [R.] valley.

Ptilura was established in the Limacodidae; it was placed in the Notodontidae by Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 407.

PTILURODES Kiriakoff, 1964, Bonn. zool. Beitr. 14: 255.

Type-species: *Ptilurodes castor* Kiriakoff, 1964, *ibidem* 14: 256, abb.1, fig.10, pl.1, fig.9, by original designation. China: north Yunnan, Li-Kiang.

‡PTITOPHOROIDES; misspelling. See Ptilophoroides Matsumura, 1920.

PTOCHOSIPHLA Meyrick, 1933, Exot. Microlepid. 4: 438.
NOCTUIDAE

Type-species: *Ptochosiphla oedipus* Meyrick, 1933, *ibidem* 4: 438, by monotypy. Holotype &, Fiji: Vunidawa, (BMNH).

Ptochosiphla was established in the Pyralididae, now Pyralidae; on the advice of and with acknowledgements to our colleagues G.S. Robinson and M. Shaffer, it is here transferred to the Noctuidae, Hypeninae.

‡PTYCHOGLENE Boisduval, 1870. See Ptychoglene Felder, 1874.

PTYCHOGLENE Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.26. LITH Type-species: Ptychoglene erythrophora Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.26, by monotypy. Mexico: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 8.

‡PTYCHOGLENE Boisduval, 1870, Considérations Lépid. Guatemala: 95 (but attributed to Felder, an incorrect authorship).

An unavailable name under the *Code*, Article 11(d) as it was first published in the synonymy of *Lithosia* Fabricius, 1798, and not subsequently treated as an available name as from Boisduval, 1870. The only originally included nominal species was also unavailable but was later established as the type-species of *Ptychoglene* Felder, 1874. Boisduval presumably had access to Felder's plates and legends (to which Boisduval refers) prior to their publication by Felder in 1874.

‡PTYCHOTRICHOS; misspelling. See Ptychotricos Schaus, 1894. CTEN

PTYCHOTRICHUS Zerny, 1912. See Ptychotricos Schaus, 1894.

PTYCHOTRICOS Schaus, 1894, Proc. zool. Soc. Lond. 1894: 227.

Type-species: *Ptychotricos zeus* Schaus, 1894, *ibidem* **1894**: 228, by monotypy. VENEZUELA: Aroa.

‡PTYCHOTRICHOS; Zerny, 1912, in Wagner, Lepid. Cat. 7: 135.

An incorrect subsequent spelling of *Ptychotricos* Schaus, 1894.

PTYCHOTRICHUS Zerny, 1912, in Wagner, Lepid. Cat. 7: 135.

An unjustified emendation of *Ptychotricos* Schaus, 1894, (but cited as an emendation of ‡*Ptychotrichos*).

A junior homonym of *Ptychotrichus* Kuwert, 1896, *Novit.* zool 3: 227, – Insecta, Coleoptera. The objective replacement name is *Ptychotricos* Schaus, 1894.

PULIA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 406.

Type-species: *Peridea albimaculata* Okano, 1958, *Rep. Gakugei Fac. Iwate Univ.* 13 (2): 53, pl.1, fig.5, pl.2, figs 5-7, by original designation. TAIWAN: central, Puli.

HORAIA Nakamura, 1973, Ent. Rev. Japan 25: 55.

Type-species: *Peridea albimaculata* Okano, 1958, by original designation (but designated as ‡*albomaculata*, an incorrect subsequent spelling).

A junior homonym of *Horaia* Tonnoir, 1930, *Rec. Indian Mus.* 32: 193, – Insecta, Diptera. The objective replacement name is *Pulia* Kiriakoff, 1974.

PERIDOPSIS Nakamura, 1976, Ent. Rev. Japan 29: 50.

Peridonsis was established as an objective replacement name

Peridopsis was established as an objective replacement name for Horaia Nakamura, 1973.

‡PULODONTIS; misspelling. See Euchila Billberg, 1820.

PURIPLUSIA Chou & Lu, 1974, Acta ent. sin. 17: 71, 77.

Type-species: Plusia purissima Butler, 1878, Ann. Mag. nat.

Hist. (5) 1: 202, by original designation. Type(s) JAPAN:

Puriplusia in the Noctuidae, Plusiinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

PURIUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 719.

Type-species: Purius sordidus Walker, 1855, ibidem 3: 719,

by monotypy. [BRAZIL]: Para, VENEZUELA.

P. sordidus is a junior subjective synonym of Phalaena pilumnia Stoll, 1780, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 36 (dated from the wrapper of the part and not from the title-page of the volume, 1782).

PUSIOLA Wallengren, 1863, Wien. ent. Monatschr. 7: 146,

Type-species: Lithosia flavicosta Wallengren, 1860, ibidem 4: 45, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 326. [South Africa] Caffraria.

Pusiola was redescribed as new by Wallengren in 1865, K. svenska Vetensk Akad. Handl. 5 (4): 44.

PUSIOLANIA Strand, 1912, Arch. Naturgesch. 78 (A) 9: 93.

Type-species: Pusiola occidentalis Strand, 1912, ibidem 78 (A) 9: 92, by monotypy. CAMEROUN: Bibundi.

PYCNOGRAPHA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 66.

Type-species: Pycnographa tamarix Kiriakoff, 1958, ibidem 66: 66, pl.3, fig.16, pl.8, fig.12, pl.9, fig.40, by original designation. [ZAIRE]: Sankaru, Lusambo.

PYDNA Walker, 1856. See Eupydna Fletcher nom. n. NOTO

PYDNAODES Hampson, 1911, Ann. Mag. nat. Hist. (8) 8:

Type-species: Tessellarctia distincta Rothschild, 1909, Ann. Mag. nat. Hist. (8) 4: 207, by original designation. [BRAZIL]: Upper Amazon, Fonté Boa.

PYDNELLA Roepke, 1943, Natuurh. Maandbl. 32: 93. NOTO Type-species: Pydnella monticola Roepke, 1943, ibidem 32: 93, figs 11a, 11b, 11c, by monotypy. Java: eastern, Djunggo-

P. monticola is a subspecies of Pydna rosacea Hampson,

1896, Fauna Br. India (Moths) 4: 458.

PYGAERA Ochsenheimer, 1810, Schmett. Eur. 3: 224. NOTO Type-species: Bombyx timon Hübner, [1803] 1796, Samml. eur. Schmett. 3: pl.22, fig.86, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 611. TYPE-LOCALITY: not stated, [EUROPE].

Invalid type-species designation: Phalaena bucephala Linnaeus, 1758, a nominal species that Ochsenheimer included doubtfully in Pygaera. Under the Code, Article 67(h), a doubtfully included nominal species may not be validly desig-

nated as type-species.

Phalaena bucephala was designated by Curtis, 1835, Br. Ent. 12: 530; by Westwood, 1840, Introd. mod. Classif. Insects 2 (Synopsis Genera Br. Insects): 90; Stephens, 1845, in Smedley, Rose & Rose, Encycl. Metropolitana 23: 694 and by Blanchard, 1845, in Castelnau, Hist. nat. Insectes 3: 496.

Phalaena curtula Linnaeus, 1758, was designated by Boisduval, 1836, Hist. nat. Insectes (Spec. gén. Lépid.) 1: 129, 135. Phalaena bucephala Linnaeus, 1758, was also designated by

Boisduval, 1836, ibidem 1: 142.

In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different type-species for each generic name. In his 'Exposé de notre méthode', pages 155-690, no typespecies designation was made for any of the genera that he himself used. Under the *Code*, Article 69(a)(iii), the type-species designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other authors.

PYGAERIDIA Kiriakoff, 1967, Biologia gabon. 3: 339. NOTO Type-species: Pygaeridia cornuta Kiriakoff, 1967, ibidem 3: 340, fig.26, by original designation. GABON: Makokou.

PYGAERINA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14:

Type-species: Pygaerina lugubris Gaede, 1928, ibidem 14: 425, by subsequent designation by Kiriakoff, 1960, Revue Zool. Bot. afr. 61: 62. CAMEROUN, ZAIRE: [Kivu], Lowa [valley].

The date of publication of the part of Seitz containing page 425 was cited on that page as 30.iii.1928 in both the German and the English editions.

PYGARCTIA Grote, 1871, Can. Ent. 3: 124. ARCT Type-species: Pygarctia abdominalis Grote, 1871, ibidem 3: 124, by monotypy. U.S.A.: Alabama.

PYGETERA Kirby, 1892. See Cataphractes Felder, 1874.

PYGOCTENUCHA Grote, 1883, Trans. Kans. Acad. Sci. 8: 37, 46.

Type-species: Ctenucha harrisii Boisduval, 1869, Annls Soc. ent. Belg. 12: 72, by original designation. U.S.A.: California, near Los Angeles.

C. harrisii is a junior subjective synonym of Apistosia terminalis Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 478.

See also Protosia Hampson, 1900.

PYGOPTERYX Staudinger, 1887, in Romanoff, Mém. Lépid.

Type-species: Pygopteryx suava Staudinger, 1887, ibidem 3: 230, pl.13 [recte 17], fig.4, by monotypy. [U.S.S.R.]: Amur region, Sidemi and Vladivostok.

Pygopteryx was established in the Notodontidae; it was placed in the Noctuidae by Inoue, 1958, Check List Lepid. Japan (5): 497.

PYRALIDIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.23. Type-species: Pyralidia deserta Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.23, by monotypy. [U.S.A.]: Utah, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 8.

PYRALOPSIS Boisduval, 1870, Considérations Lépid. Guatemala: 94 (but attributed to Felder, an incorrect authorship). **PYRALIDAE**

Type-species: Pyralopsis divisa Boisduval, 1870, ibidem: 94,

by monotypy. Guatemala.

Pyralopsis and two of the three originally included nominal species, ‡P. homalochroa and ‡P. guatemalteca, were not described by Boisduval and were attributed to Felder, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105; but this plate, which was included in Heft 4 of Felder Rogenhofer's work, was not published until November 1874. The only described and therefore nomenclaturally available species included by Boisduval in Pyralopsis was P. divisa Boisduval, 1870. It is probable that Boisduval had obtained a proof copy of Felder & Rogenhofer's plate and its legend privately, prior to its publication in Vienna (see Fletcher, 1979, Generic Names Moths World 3: 'Dates of some important publications').

Pyralopsis was established in the Lithosides, now Arctiidae;

it was placed by Prout, 1918, Novit. zool. 25: 405, in the Dioptidae, with P. homalochroa Felder incorrectly cited as type-species; and is here transferred to the Pyralidae, Pyralinae as a junior subjective synonym of Mapeta Walker, 1863.

Pyralopsis divisa is a junior subjective synonym of Mapeta xanthomelas Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 27: 17.

PYRAMOCERA Butler, 1880, J. Linn. Soc. (Zool.) 15: 85.

Type-species: Pyramocera fuliginea Butler, 1880, ibidem (Zool.) 15: 85, fig., by original designation. MADAGASCAR: Fianarantsoa.

PYRODESMA Boisduval, 1870. See Calodesma Hübner, [1820].

PYRRHARCTIA Packard, 1864, Proc. ent. Soc. Philad. 3: 120.

Type-species: Phalaena isabella Smith, 1797, in Smith & Abbot, Nat. Hist. rarer lepid. Insects Georgia 2: 131, pl.66, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 244. [U.S.A.]: Georgia.

PYRRHOPTERYX Hering, 1941, Revue Zool. Bot. afr. 35: 81.

Type-species: Pyrrhopteryx amabilis Hering, 1941, ibidem 35: 82, pl.9, fig.11, by original designation. [ZAIRE] Congo: [Shaba], [Lubumbashi] Elisabethville.

PYRSOPSYCHE Kiriakoff, 1968, Biologia gabon. 4: 45. NOTO

Type-species: Pyrsopsyche pyrrhias Kiriakoff, 1968, ibidem 4: 46, text-fig.58, pl.2, fig.19, by original designation. GABON: Belinga.

QUADRICALCARIFERA Strand, 1916, Arch. Naturgesch. 81 (A) 12: 160. NOTO

Type-species: Stauropus subgeneris Strand, 1916, ibidem (A) 12: 160, by monotypy. TAIWAN: Kosempo.

Quadricalcarifera was established as a subgenus of Stauropus Germar, 1812.

QUISTA Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 180.

Type-species: Quista cota Kiriakoff, 1959, ibidem 95: 180, fig. 25, by original designation. ZAIRE: Sankuru, Lusambo.

RABTALA Draeseke, 1926, Dt. ent. Z. Iris. 40: 105. Type-species: Trabala cristata Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 480, by original designation. JAPAN: Yokohama.

LAMPRONADATA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 23.

Type-species: Trabala cristata Butler, 1877, by original designation.

A junior objective synonym of Rabtala Draeseke, 1926.

RACHIA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 70. NOTO

Type-species: Rachia plumosa Moore, 1879, ibidem (1): 70, by monotypy. INDIA: Darjeeling.

‡RHACHIA; Gaede, 1934, in Strand, Lepid. Cat. 59: 190. An incorrect subsequent spelling of Rachia Moore, 1879.

RACIIIADES Kiriakoff, 1967, in Wytsman, Genera Insect. 217

Type-species: Semidonta lichenicolor Oberthür, Etudes Lépid. comp. 5 (1): 336, pl.83, fig.776 [CHINA]: [western], Ta-tsien-Lou.

RADAMARIA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80:

Type-species: Dasychira miselioides Kenrick, 1914, Trans. ent. Soc. Lond. 1913: 595, pl.32, fig.28, by original designation. MADAGASCAR: Central.

RAHONA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 228.

Type-species: Dasychira albilunula Collenette, 1936, Novit. zool. 40: 161, pl.12, fig.17, by original designation. MADA-GASCAR: E., Périnet (149km E. of Tananarivo).

RAJACOA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 434.

Type-species: Cypra forbesi Druce, 1899, Ann. Mag. nat. Hist. (7) 3: 469, by original designation. New Guinea.

RAJENDRA Moore, 1879, in Hewitson & Moore, Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson (1): 43.

Type-species: Rajendra lativitta Moore, 1879, ibidem (1): 43, by original designation. [INDIA]: [Bihar] Behar, Parisnath

R. lativitta is a junior subjective synonym of Chelonia perrottetii Guérin-Méneville, [1844], Icon. Règne anim. Cuvier (Insectes): 515.

RAMADRA Nye nom. n. for Mardara Walker, 1865. Type-species: Mardara calligramma Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 402, by monotypy (of Mardara Walker, 1865). [INDIA]: [W. Bengal], Darjeeling.

MARDARA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 402.

Type-species: Mardara calligramma Walker, 1865, by monotypy.

A junior homonym of Mardara Walker, [1859] 1858, ibidem 16: 157, - Lepidoptera, Noctuidae. The objective replacement name is Ramadra Nye, nom. n.

RAMESA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1016.

Type-species: Ramesa tosta Walker, 1855, ibidem 5: 1017, by monotypy. INDIA: northern.

RANGHANA Moore, 1878, Proc. zool. Soc. Lond. 1878: 22.

Type-species: Ranghana punctata Moore, 1878, ibidem 1878: 22, pl.2, fig.12, by monotypy. [INDIA]: Calcutta.

‡RHANGHANA; Hampson, 1894, Fauna Br. India (Moths) **2**: 122, 124.

An incorrect subsequent spelling of Ranghana Moore, 1878.

‡RHANGANA; Draudt, 1914, in Seitz, Gross-Schmett. Erde

An incorrect subsequent spelling, in both the German and the English editions of Seitz, of Ranghana Moore, 1878.

Neave, 1940, Nomencl. zool. 4: 35 incorrectly listed ‡Rhangana as a misspelling of Rhangena Moore, 1886, Lepid. Ceylon 3: 375, - Lepid., Noctuidae.

RAPHIPEZA Butler, 1880, Ann. Mag. nat. Hist. (5) 5: 386. LASIOCAMPIDAE

Type-species: Gogane turbata Butler, 1879, ibidem (5) 4:

237, by original designation.

Raphipeza was established in the Liparidae, now Lymantriidae; it was placed in the Bombycidae by Saalmüller, 1884, Lepid. Madagascar (1): 240; and was transferred from the 'Liparidae' to the Lasiocampidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497.

RASEMIA Kiriakoff, 1962, Revue Zool. Bot. afr. 66: 33. NOTO Type-species: Ramesa macrodonta Hampson, 1909, Trans. zool. Soc. Lond. 19: 119, pl.4, fig.38, by original designation. UGANDA: S.E. Ruwenzori, Mokia.

REBITA Kiriakoff, 1962, *Bull. Annls Soc. r. ent. Belg.* **98**: 151 (key), 211.

Type-species: *Pydna indica* Moore, 1879, in Hewitson & Moore, *Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson* (1): 67, by original designation. INDIA: Calcutta.

Rebita was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 64.

REDOA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 826.

Type-species: *Redoa submarginata* Walker, 1855, *ibidem* 4: 826, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 435. [BANGLADESH]: [Sylhet], Silhet, EAST INDIES, [SRI LANKA] Ceylon.

‡REGMATOPHORA; misspelling. See Rhegmatophila Standfuss, 1888.

REGOBARROSIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 59.

Type-species: Automolis aureogrisea Rothschild, 1910, Novit. zool. 16: 38, pl.5, fig.36 PERU: Carabaya Mts, Santo Domingo.

REPA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1898.

Type-species: *Repa cana* Walker, 1866, *ibidem* 35: 1898, by monotypy. U.S.A.

R. cana is a junior subjective synonym of Clemensia albata Packard, 1864, Proc. ent. Soc. Philad. 3: 101.

See also Clemensia Packard, 1864, and Uxia Walker, 1866.

REPENA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 799.

Type-species: Repena cervina Walker, 1855, ibidem 4: 800, by monotypy. EAST INDIES.

RETHONA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1042.

Type-species: *Rethona strigosa* Walker, 1855, *ibidem* 5: 1043, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 592. SOUTH AFRICA.

Rethona was established in the Notodontidae; it was placed in the Cossidae by Gaede, 1929, in Seitz, Gross-Schmett. Erde 14: 550.

RETILA Boisduval, 1870. See Ephialtias Hübner, [1819].

REZIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 149.

Type-species: Glaucopis erythrarchos Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 164, by monotypy. Brazil.

Rezia was established as the objective replacement name for Lagaria Walker, 1854; it was used as a valid genus but retained in the Zygaenidae by Kirby, 1892, ibidem 1: 149; and was included in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 225.

LAGARIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 164.

Type-species: Glaucopis erythrarchos Walker, 1854, by monotypy. Brazil.

Lagaria was established as a subgenus of Glaucopis Fabricius, 1807, in the Zygaenides, now Zygaenidae.

A junior homonym of Lagaria Dallas, 1852, List Specimens

hemipt. Insects Colln Br. Mus. 2: 442, – Insecta, Hemiptera. The objective replacement name is Rezia Kirby, 1892.

‡RIZIA; Zerny, 1912, in Wagner, Lepid. Cat. 7: 156.
An incorrect subsequent spelling of Rezia Kirby, 1892.

RHABDATOMIS Dyar, 1907, Jl N.Y. ent. Soc. 15: 227. LITH Type-species: Rhabdatomis zaba Dyar, 1907, ibidem 15: 227, by original designation. MEXICO: Orizaba.

RHABDOMARCTIA Kiriakoff, 1953, Annls Mus, r. Congo Belge (8vo Sci. zool.) 26: 30. THYRET

Type-species: *Metarctia rubrilineata* Bethune-Baker, 1911, *Ann. Mag. nat. Hist.* (8) 7: 533, by original designation. ANGOLA: N., [Vila Salazar] N'Dalla Tando.

RHABDOTURNACA Kiriakoff, 1967, Zool. Meded. Leiden 42: 192. NOTO

Type-species: *Turnaca subcarnea* Warren, 1903, *Novit. zool.* 10: 120, by original designation. New Guinea: [Papua New Guinea].

‡RHACHIA; misspelling. See Rachia Moore, 1879. NOTO

RHAGONIS Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 270.

ARCT

Type-species: *Rhagonis bicolor* Walker, 1862, *ibidem* (3) 1: 270, by monotypy. [U.S.A.]: California.

R. bicolor is a junior subjective synonym of Arctia vagans Boisduval, 1852, Annls Soc. ent. Fr. (2) 10: 322.

RHAGOPHANES Zeller, 1853, Bull. Soc. imp. Nat. Moscou 26 (3): 508.

Type-species: Rhagophanes tortriciformis Zeller, 1853, ibidem 26 (3): 511, fig., by monotypy. JAVA.

‡RHANGANA; misspelling. See Ranghana Moore, 1878.

‡RHANGHANA; misspelling. See Ranghana Moore, 1878.

RHANIDOPHORA Wallengren, 1858, Ofvers. K. Vetensk-Akad. Förh. Stockh. 15: 213.

NOCTUIDAE

Type-species: *Phalaena phedonia* Stoll, 1781, in Cramer, *Uitlandsche Kapellen (Papillons exot.)* 4: 110, pl.347, fig.C, by original designation. [SOUTH AFRICA]: [Cape of Good Hope] Kaap de Goede Hoop.

Rhanidophora was established in the 'Chelonides', now Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 462; and was transferred to the Noctuidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498.

RHAPIGIA Schaus, 1928, Proc. U.S. natn Mus. 73: 70. NOTO Type-species: Hapigia accipiter Schaus, 1892, Proc. zool. Soc. Lond. 1892: 340 (but designated as ‡accipites, an incorrect subsequent spelling), by original designation. [BRAZIL]: Rio Janeiro.

RHEGMATOPHILA Standfuss, 1888, Berl. ent. Z. 32: 239.

Type-species: *Clostera alpina* Bellier, 1881, *Annls Soc. ent. Fr.* (5) **10**: 367, pl.11, figs 10, 10a, by monotypy. France: Basses Alpes, Digne.

‡REGMATOPHORA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 429.

An incorrect subsequent spelling of Rhegmatophila Standfuss, 1888.

RHENEA Saalmüller, 1884, Lepid. Madagascar (Abt.1): 242.

Type-species: Notodonta circumcincta Saalmüller, 1880, Ber. naturf. Ges. 1879-80: 268, by senckenb. monotypy. MADAGASCAR.

RHENEADES Kiriakoff, 1962, Tijdschr. Ent. 105: 252.

Type-species: Rheneades flavescens Kiriakoff, 1962, ibidem 105: 252, fig.18, by original designation. UGANDA: Entebbe.

RHENEINA Kiriakoff, 1971, Bull. Annls Soc. r. ent. Belg. 107: 314.

Type-species: *Rheneina buceros* Kiriakoff, 1971, *ibidem* 107: 316, text-fig.1, by original designation. [ZIMBABWE-]RHODE-SIA: Vumba.

‡RHINA; misspelling. See Rhipha Walker, 1854. ARCT

RHIPHA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 273.

Type-species: Euchromia strigosa Walker, 1854, ibidem 1:

273, by monotypy. Brazil: Rio de Janeiro.

Rhipha was established as a subgenus of Euchromia Hübner, [1819], in the Zygaenides, now Zygaenidae; it was placed, as a valid genus, in the Arctiidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 171.

‡RHINA; Travassos, 1954, Revta bras. Ent. 1: 213. An incorrect subsequent spelling of Rhipha Walker, 1854.

RHIPIDARCTIA Kiriakoff, 1953, Annls Mus. r. Congo Belge (8vo Sci. zool.) 26: 28.

Type-species: Elsa rubrosuffusa Kiriakoff, 1953, ibidem 26: 28, pl.2, fig.22, pl.6, fig.73, by original designation. [ZAIRE] Congo Belge: Tshuapa, Eala.

Rhipidarctia was established as a subgenus of Elsa Kiriakoff,

RHODAREAS Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 254.

Type-species: Arctia melanopsis Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 280, by PRESENT DESIGNATION. [SRI LANKA] Ceylon.

Rhodareas was attributed to Butler (as a manuscript name) by Kirby.

RHODESANA Bethune-Baker, 1908, Ann. Mag. nat. Hist. (8) LYMA

Type-species: Rhodesana crenulata Bethune-Baker, 1908, ibidem (8) 2: 261, by original designation. [ZAMBIA] Rhodesia, N.E.: Fort Jameson.

RHODOCLEPTRIA Hampson, 1903. See Chazaria Moore, 1881. NOCTUIDAE

RHODOGASTRIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 172.

Type-species: Phalaena amasis Cramer, 1779, Uitlandsche Kapellen (Papillons exot.) 3: 23, pl.206, fig.B, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 223. TYPE-LOCALITY: stated in error to be Surinam; the species does not occur in the New World, but is widespread in SOUTH AFRICA.

P. amasis is dated from the wrapper of the part and not from the title-page of the volume, 1782.

See also Munychia Wallengren, 1858.

DIONYCHOPUS Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 94, 96.

Type-species: Phalaena amasis Cramer, 1779, by subsequent designation by Hampson, 1900, Ann. S. Afr. Mus. 2: 58.

A junior objective synonym of *Rhodogastria* Hübner, [1819]. Dionychopus was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

RHODOGRAPHA Schaus, 1899, Jl N.Y. ent. Soc. 7: 215.

Type-species: Rhodographa phaeoplaga Schaus, 1899, ibidem 7: 215, by monotypy. Brazil: S.E., São Paulo.

The name Rhodographa probably originated from Hampson but was used and made available by Schaus prior to its proposal by Hampson, 1900, as the name for a new genus.

RHODOGRAPHA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 87, 295.

Type-species: Rhodographa phaeoplaga Schaus, 1899, by original designation.

A junior homonym of Rhodographa Schaus, 1899. The objective replacement name is Rhodographa Schaus, 1899.

RHODOGRAPHA Hampson, 1900. See Rhodographa Schaus, 1899. LITH

RHOGALIA Hübner, [1825] 1816, Verz. bekannter Schmett.: 396.

Type-species: Phalaena epigena Stoll, 1790, in Cramer, Uitlandsche Kapellen (Papillons exot.) (Aanhangsel): 72, pl.16, figs 1, 1A, 1B (but included by Hübner as ‡epigenana, an incorrect subsequent spelling), by monotypy. SURINAM.

P. epigena is dated from the wrapper of the part, not from the title-page of the volume which is dated 1791.

RHOPTROPHALAENA Hampson, 1918. See Cocytia Boisduval, 1828.

RHUDA Walker, 1857, List Specimens lepid. Insects Colln Br. Mus. 11: 611 (key), 621.

Type-species: Rhuda basifera Walker, 1857, ibidem 11: 621,

by monotypy. [Brazil]: Rio de Janeiro.

R. basifera is a junior subjective synonym of Phalaena focula Stoll, 1782, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 187, pl.383, figs G, H.

Rhuda was established in the Noctuidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 617.

ORTHOSOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.18.

Type-species: Orthosoma diffusum Felder, 1874, ibidem 2 (Abt.2): pl.97, fig.18, by monotypy. VENEZUELA, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 7.

A junior homonym of Orthosoma Audinet-Serville, 1832, Annls Soc. ent. Fr. 1: 155, - Insecta, Coleoptera. There is no objective replacement name, but O. diffusum is currently treated as congeneric with Rhuda basifera Walker, 1857, the typespecies of Rhuda Walker, 1857; the latter is therefore available for use as a subjective replacement name.

RHYNCHOPHALERA Aurivillius, 1904, Ark. Zool. 2 (4): 1. NOTO

Type-species: Rhynchophalera signata Aurivillius, 1904, ibidem 2 (4): 2, text-fig.9, by monotypy. CAMEROUN.

RHYNCHOPHALERINA Kiriakoff, 1969, Faune Madagascar **29**: 84. NOTO

Type-species: Rhynchophalerina inexpectata Kiriakoff, 1969, ibidem 29: 86, text-figs 53, 54, pl.3, fig.28, by original designation. MADAGASCAR: western, 64 km east of Tuléar, Andranovory forest.

RHYNCHOPYGA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.30. CTEN

Type-species: Rhynchopyga ichneumonea Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.30, by monotypy. [COLOMBIA]: Bogotà, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 8.

RHYPARIA Hübner, [1820] 1816, Verz. bekannter Schmett.: 183.

Type-species: Phalaena purpurea Linnaeus, 1767, Syst. Nat. (Edn 12) 1 (2): 828, by subsequent designation by Kirby, 1892, in Synonymic Cat. Lepid. Heterocera 1: 260, but designated as Phalaena purpurata Linnaeus; see below. Type-locality: not stated, [EUROPE].

When Kirby designated as type-species Phalaena purpurata Linnaeus, 1758, Syst. nat. (Edn 10) 1: 505, a nominal species not originally included in Rhyparia, he also placed purpurea, a nominal species originally included in Rhyparia, as a junior synonym of purpurata. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species.

P. purpurea Linnaeus, 1767, may be regarded either as an emendation and hence a junior objective synonym, or else a junior subjective synonym of P. purpurata Linnaeus, 1758.

RHYPARIOIDES Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 395.

Type-species: Rhyparioides nebulosa Butler, 1877, ibidem (4) 20: 396, by original designation. JAPAN: Yokohama; Hakodate.

‡RHYPOPTERIX; misspelling. See Rhypopteryx Aurivillius, 1879. LYMA

RHYPOPTERYX Aurivillius, 1879, Ofvers. K. Vetensk Akad. Förh. Stockh. 36 (7): 57.

Type-species: Rhypopteryx sordida Aurivillius, 1879, ibidem 36 (7): 58, by monotypy. [Namibia]: Damara[land].

Rhypopteryx was established in the Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 491.

‡RHYPOPTERIX; Hering, 1927, in Seitz, Gross-Schmett. Erde 14: 198.

An incorrect subsequent spelling of Rhypopteryx Aurivillius, 1879.

‡Rhypopterix was the spelling used as the generic heading, but Rhypopteryx was used for the title of the page. In the English edition published on the same day Rhypopteryx was used as the generic heading as well as for the title of the page.

RHYPOTOSES Collenette, 1931, Zool. Meded. Leiden 14: 182.

Type-species: Rhypotoses atima Collenette, 1931, ibidem 14: 182, by original designation. JAVA: W., Malabar Geb., Gouv. Kina-Ondern, Tjinjiroean.

RIBALDIA Dyar, 1916, Proc. U.S. natn. Mus. 51: 22. NOTO Type-species: Ribaldia amatame Dyar, 1916, ibidem 51: 22, by original designation. MEXICO: Hidalgo, Guerrero Mill.

RICCIA Travassos, 1953, Papéis Dep. Zool. S. Paulo 11: 280.

Type-species: Trichura aliaria Druce, 1890, Proc. zool. Soc. Lond. 1890: 494, pl.42, fig.3, by original designation. [PERU]: Amazon R., Pebas.

RICINE Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 780 (kev), 824.

Type-species: Ricine suffusa Walker, 1855, ibidem 4: 824, by monotypy. JAVA.

RIFARGIA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 258.

Type-species: Rifargia xylinoides Walker, 1862, ibidem (3) 1: 259, by monotypy. [BRAZIL]: Rio de Janeiro.

RIGEMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1079.

Type-species: Rigema vittata Walker, 1855, ibidem 5: 1079, by monotypy. SIERRA LEONE.

RILIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1075.

Type-species: Rilia lanceolata Walker, 1855, ibidem 5: 1075. by monotypy. [BANGLADESH]: [Sylhet], Silhet, [SRI LANKA] Ceylon.

Rilia was established in the Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 488.

RINCODES Schaus, 1901, Trans. ent. Soc. Lond. 1901: 262 (key), 313.

Type-species: Crinodes minuta Druce, 1900, Ann. Mag. nat. Hist. (7) 5: 516, by original designation. [Brazil]: Amazons, Manaos.

RIPISA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1082.

Type-species: Ripisa viridescens Walker, 1855, ibidem 5: 1083, by monotypy. [BRAZIL]: Rio de Janeiro.

RIVOTRA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 228.

Type-species: Dasychira viridipicta Kenrick, 1914, Trans. ent. Soc. Lond. 1913: 594, pl.32, fig.26, by original designation. MADAGASCAR: Central.

‡RIZIA; misspelling. See Rezia Kirby, 1892. CTEN

ROBINSONIA Grote, 1865, Proc. ent. Soc. Philad. 5: 241.

Type-species: Robinsonia formula Grote, 1865, ibidem 5: 241, pl.4, fig.3, by monotypy. CUBA.

RODNEYA Kiriakoff, 1967, Tijdschr. Ent. 110: 63. Type-species: Rodneya caudata Kiriakoff, 1967, ibidem 110: 63, fig. 12, by original designation. SUMATRA: western, Lebong Tandai.

ROEPKEELLA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 40.

Type-species: Pseudogargetta fuscicollis Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 618, pl.81 row f, by original designation. MALAYSIA, BORNEO, JAVA, [SULAWESI] Celebes.

ROLEPA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 770.

Type-species: Rolepa delineata Walker, 1855, ibideni 3: 770, by monotypy. BRAZIL: Rio Janeiro.

Rolepa was established in the Arctiidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 851; and was placed in the Lymantriidae by Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 557.

See also Olenoptera Herrich-Schäffer, [1856], and Athrula Walker, 1862.

ROMALEOSTAURA Kiriakoff, 1960, Revue fr. Ent. 27: 173.

NOTO

Type-species: Romaleostaura insularis Kiriakoff, 1960, ibideni 27: 174, fig.1, by original designation. MADAGASCAR: eastern, Km. 57 route d'Anosibe.

ROMALINA Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 23.

Type-species: *Eurystaura simplex* Gaede, 1928, *in* Seitz, *Gross-Schmett. Erde* 14: 429, pl. 71 row c, by original designation. CAMEROUN.

ROMANOFFIA Heylaerts, 1884, Annls Soc. ent. Belg. 28 (Bull.): xciv.

Type-species: Romanoffia imperialis Heylaerts, 1884, ibidem 28 (Bull.): xcv, by monotypy. [Panama]: [Chiriqui] Chirequi.

ROMUALDIA Rego Barros, 1957, Revta bras. Biol. 17: 89.

Type-species: Amastus opharina Schaus, 1921, Insecutor Inscit. menstr. 9: 169, by original designation. BRAZIL: Nova Friburgo.

ROMUALDISCA d'Andretta & Travassos, 1946, Livro de Homenagem a R. Ferreira d'Almeida: 19.

Type-species: Romualdisca dalmeidai d'Andretta & Travassos, 1946, ibidem: 19, fig., by original designation. BRAZIL: São Paulo state, São Paulo.

ROPPA Kiriakoff, 1961, Bull. Annls Soc. r. ent. Belg. 97: 114.

Type-species: *Pydna rhabdophora* Hampson, 1910, *Ann. Mag. nat. Hist.* (8) **5**: 487, by original designation. NIGERIA: Lokoja, Mt Patti.

‡ROPTROPHALAENA; misspelling. See Cocytia Boisduval, 1828.

ROSAMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1066.

Type-species: *Rosama strigosa* Walker, 1855, *ibidem* 5: 1066, by monotypy. Type-locality: not known, [Java].

ROSEMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1159 (key), 1168.

Type-species: Rosema dorsalis Walker, 1855, ibidem 5: 1169, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 581. BRAZIL.

R. dorsalis is a form of Phalaena zelica Stoll, 1790, in Cramer, Uitlandsche Kapellen (Papillons exot.) (Aanhangsel): 73, pl.16, figs 2-2D.

Rosema was established in the Drepanidae; it was placed in the Notodontidae by Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 696.

See also Zelica Hübner, [1825].

ROSINELLA Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 342.

Type-species: *Chadisra rosinaria* Hampson, 1910, *Ann. Mag. nat. Hist.* (8) **5**: 461, by original designation. [South Africa]: Natal, Durban.

ROSIORA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 151 (key), 160.

Type-species: *Pydna bela* Swinhoe, 1904, *Trans. ent. Soc. Lond.* 1904: 159, by original designation. [INDIA]: [Assam], Khasia Hills, Cherrapunji.

ROUGEOTIA Laporte, 1974, Bull. mens. Soc. linn. Lyon 43: 218. NOCTUIDAE

Type-species: Rougeotia aethiopica Laporte, 1974, ibidem 43: 218, figs 2, 26, by original designation. Holotype 3, ETHIOPIA: marais de Dinsho, (MNHN, Paris).

Rougeotia in the Noctuidae, Hadeninae is cited here as it

was not included in Nye, 1975, Generic Names Moths World

ROUGEOTIANA Laporte, 1974, Bull. Soc. ent. Fr. 79: 33.
NOCTUIDAE

Type-species: Sphingomorpha xanthoperas Hampson, 1926, Descr. new Genera Species Lepid. Phalaenae Subfamily Noctuinae Br. Mus.: 102, by original designation. Holotype Q, [Ghana] Gold Coast: Kumasi, (BMNH).

Rougeotiana in the Noctuidae, Ophiderinae is cited here as it was not included in Nye, 1975, Generic Names Moths World

RUANDA Strand, 1909, Dt. ent. Z. Iris 22: 111. LYMA Type-species: Ruanda aetheria Strand, 1909, ibidem 22: 113, by original designation. [RWANDA] Ruanda: N.E.

‡RUSCINA; misspelling. See Ruscino Walker, 1854. LITH

RUSCINO Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 563.

Type-species: *Phalaena menea* Drury, 1782, *Illust. nat. Hist. exot. Insects* 3: 4, pl.3, fig.2, by monotypy. BRAZIL.

‡RUSCINA; Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus, 31: 264.

An incorrect subsequent spelling of Ruscino Walker, 1854.

SACADA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 136.
PYRALIDAE

Type-species: Sacada decora Walker, 1862, ibidem (Zool.) 6: 136, by monotypy. BORNEO: Sarawak.

Sacada was established in the Notodontidae; it was placed in the Pyralidae by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 435.

SAENURA Wallengren, 1858, Ofvers. K. VetenskAkad. Förh. Stockh. 15: 214.

Type-species: Saenura flava Wallengren, 1858, ibidem 15: 214, by original designation. [SOUTH AFRICA]: Caffraria.

‡SENURA; Wallengren, 1873, Ofvers. K. VetenskAkad. Förh. Stockh. 29 (3): 50.

An incorrect subsequent spelling of Saenura Wallengren, 1858.

SAGAMORA Kiriakoff, 1967, Tijdschr. Ent. 110: 61. NOTO Type-species: Sagamora umbrina Kiriakoff, 1967, ibidem 110: 62, fig.29, by original designation. SUMATRA: Fort de Kock.

SAGARIS Walker, 1854. See Phavaraea Walker, 1854.

SAGAROPSIS Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 434.

Type-species: Sagaris horae Druce, 1885, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 144. pl.13, fig.25, by original designation. PANAMA: Bugaba, Chiriqui Volcano.

The date of publication of the part of Seitz containing page 434 was cited on page 433 as 9.i.1925 in the German edition and as 11.iii.1925 in the English edition.

SALAMBORIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 33: 855.

Type-species: Salamboria deornata Walker, 1865, ibidem 33: 855 (a junior secondary homonym, see below), by monotypy. [Colombia]: Bogota.

Salamboria was established in the Noctuidae; it was placed in the Notodontidae by Schaus, 1901, Trans. ent. Soc. Lond. 1901: 333, where it was included as a junior subjective synonym of Hemiceras Guenée, 1852. At the same time Schaus included both Epicoria deornata Walker, 1865, ibidem 33: 853, and

Salamboria deornata Walker, 1865, ibidem 33: 855, in Hemiceras; the junior name, Salamboria deornata, thus became a junior secondary homonym for which Schaus, 1901, ibidem 1901: 334 proposed Hemiceras walkeri as the objective replace-

Under the Code, Article 59(b)(i), a junior secondary homonym replaced before 1961 is permanently rejected and cannot be reinstated as a valid name unless the use of the replacement name is contrary to existing usage.

SALAPOLA Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 525.

Type-species: Salapola argentea Walker, 1863, ibidem 28: 525, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 920. [BRAZIL]: Rio [de] Janeiro.

‡SALOPOLA; Schaus, 1894, Proc. zool. Soc. Lond. 1894: 241. An incorrect subsequent spelling of Salapola Walker, 1863.

SALIOCLETA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 124.

Type-species: Saliocleta nonagrioides Walker, 1862, ibidem (Zool.) 6: 124, by monotypy. Borneo: Sarawak.

Saliocleta was established in the Arctiidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 930.

SALLAEA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.101, fig.10.

Type-species: Sallaea ochrosterna Felder, 1874, ibidem 2 (Abt.2): pl.101, fig.10, by monotypy. MEXICO: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 8.

S. ochrosterna is a junior subjective synonym of Turuptiana obliqua Walker, 1869, Characters undescr. Lepid. Heterocera:

See also Turuptiana Walker, 1869.

SALLUCA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 260 (key), 294.

Type-species: Salluca moruma Schaus, 1901, ibidem 1901: 294, pl.11, fig.11, by original designation. Mexico: Orizaba.

‡SALOPOLA; misspelling. See Salapola Walker, 1863. LITH

SALVATGEA Griveaud, 1977, Faune Madagascar 43: 221.

Type-species: Suarezia bipuncta Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 151, pl.21, row c, by monotypy (of Suarezia Hering, 1926). MADAGASCAR: Diego Suarez.

Salvatgea was proposed as the objective replacement name for Suarezia Hering, 1926.

SUAREZIA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 151.

Type-species: Suarezia bipuncta Hering, 1926, by monotypy. A junior homonym of Suarezia Budde-Lund, 1904, Revision Crustacea Isopoda terrestria (2-3): 41, - Crustacea.

The objective replacement name is Salvatgea Griveaud, 1977.

SAMERA Wallengren, 1863. See Pelosia Hübner, [1819].

SANEGA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1085.

Type-species: Sanega orcus Walker, 1855, ibidem 5: 1085, by monotypy. [BRAZIL]: Rio de Janeiro.

SANGATA Walker, 1860, List Specimens lepid. Insects Colln Br. Mus. 20: 265.

Type-species: Sangata rosea Walker, 1860, ibidem 20: 265, by monotypy. Type-locality,: stated to be Cape, without further detail. The species is now known to occur in NORTH AMERICA.

S. rosea is a junior subjective synonym of Phalaena aurora Smith, 1797, in Smith & Abbot, Nat. Hist. rarer lepid. Insects Georgia 2: 173, pl.87.

Sangata was established in the Geometridae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 585.

See also Hyparpax Hübner, [1825].

‡SUNGATA; Kirby, 1892, Synonymic Cat. Lepid. Heterocera

An incorrect subsequent spelling of Sangata Walker, 1860.

SANKURUA Collenette, 1960. See Kunusara Nye nom. n.

SANSAPTERA Schaus, 1896, Jl N.Y. ent. Soc. 4: 132. CTEN Type-species: Sansaptera cocho Schaus, 1896, ibidem 4: 132, by monotypy. PERU: Amazons, Cavallo-cocho.

SAOZANA Dyar, 1914, Proc. U.S. natn. Mus. 47: 172. LITH Type-species: Odozana leucota Hampson, 1901, Ann. Mag. nat. Hist. (7) 8: 184, by original designation. PANAMA: Cana Mines.

SAPELIA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 389.

Type-species: Sapelia limpida Swinhoe, 1903, ibidem 1903: 389, by original designation. [NIGERIA]: River Niger, Sapele.

SARAPUS Walker, 1855. See Asparus Watson nom. n. LITH

SARBENA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 137. NOLIDAE

Type-species: Sarbena lignifera Walker, 1862, ibidem (Zool.) 6: 137, by monotypy. Borneo: Sarawak.

Sarbena was established in the Notodontidae; it was placed in the Nolidae by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 589.

SARBISSA Kiriakoff, 1976, Tierreich 97: 152. AGARISTIDAE Type-species: Seudyra bostrychonota Tams, 1929, Entomologist 62: 252, pl.5, fig.3, by original designation. Holotype ♀, Fiji: Lautoka, (BMNH).

Sarbissa in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

SARIMARAIS Kiriakoff, 1962, Revue Zool. Bot. afr. 66: 28.

Type-species: Chadisra bicolor Distant, 1899, Ann. Mag. nat. Hist. (7) 4: 360, by original designation. [South Africa]: Transvaal, Pretoria.

SAROSA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 151.

Type-species: Glaucopis sesiiformis Walker, 1854, ibidem 1: 151, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 133. VENEZUELA.

Sarosa was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 185.

SAROTHROPYGA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.100, fig.23. pl.101, fig.13.

Type-species: Sarothropyga rhodopepla Felder, 1874, ibidem 2 (Abt.2): pl.100, fig.23, pl.101, fig.13, by monotypy. [South Africa]: Knysna, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 8.

S. rhodopepla is a junior subjective synonym of Morasa

modesta Walker, 1855, the type-species of Morasa Walker, 1855.

See also Morasa Walker, 1855.

‡SAROTHRORYGA: Felder, 1874, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.100, fig.23.

An incorrect (of a multiple) original spelling of *Sarothropyga* Felder, 1874.

‡Sarothroryga was a lapsus in the engraved script of the legend to pl.100. The name was spelled as Sarothropyga on pl.101 and in the Erklärung to pls 75-107: 8.

‡SAROTHRORYGA; misspelling. See Sarothropyga Felder, 1874.

SARSINA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 779 (key), 800.

Type-species: Sarsina purpurascens Walker, 1858, ibidem 4: 800, by monotypy. [Mexico]: Yucatan.

Sarsina was established in the Liparidae, now Lymantridae; it was transferred from the Liparidae, but not placed elsewhere, by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497; it is currently retained in the Lymantriidae.

SATARA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 160.

Type-species: Satara aequata Walker, [1865] 1864, ibidem 31: 160, by monotypy. [SULAWES] Celebes.

See also Arctioneura Felder, 1874.

SAURITA Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 101, 103.

Type-species: *Sphinx cassandra* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 494, by monotypy (but attributed to Cramer, an incorrect authorship). AMERICA: [tropical].

Saurita was attributed to Boisduval, an incorrect authorship, by Herrich-Schäffer.

Saurita was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855; the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

SAURITINIA Dyar, 1905, Proc. U.S. natn. Mus. 29: 173.

Type-species: *Sauritinia dubiosa* Schaus, 1905, *ibidem* **29**: 188, by original designation (in anticipation of Schaus's description of *dubiosa*). FRENCH GUIANA: Maroni River, St Jean.

Sauritinia was available but was without an available included species between its date of publication 1905 October 3rd and that of its type-species 1905 October 11th by Schaus.

SAVARA Walker, [1865]. See Bucaea Walker, 1866. ARCT

SAWIA Kiriakoff, 1967, Zool. Meded. Leiden 42: 211. NOTO Type-species: Sawia undulata Kiriakoff, 1967, ibidem 42: 212, fig.14, by original designation. New Guinea: [Irian Jaya], [Sukarnapura] Hollandia.

SCAEODORA Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 691, 731.

Type-species: Scaeodora omophanes Meyrick, 1886, ibidem (2) 1: 731, by monotypy. AUSTRALIA: Victoria, Melbourne.

SCAEOPTERYX Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 342.

Type-species: *Hoplitis curvatula* Rothschild, 1917, *Novit. zool.* **24**: 251, pl.6, fig.11, by original designation. [ZAIRE] Congo: Kasai river.

SCALMICAUDA Holland, 1893, Psyche, Camb. 6: 489.
NOTO

Type-species: Scalmicauda benga Holland, 1893, ibidem 6: 489, by original designation. [GABON]: [Ogooué] Ogové R. valley.

SCAPHIDRIOTIS Turner, 1899, Trans. Proc. R. Soc. S. Aust. 23: 14.

Type-species: Scaphidriotis xylogramma Turner, 1899, ibidem 23: 14, by monotypy. AUSTRALIA: Brisbane.

Scaphidriotis and S. xylogramma were both attributed to Meyrick, an incorrect authorship, by Turner.

SCAPHOCERA Saalmüller, 1884, Lepid. Madagascar (Abt.1): 181.

Type-species: Cypra marginepunctata Saalmüller, 1878, Ber. senckenb. naturf. Ges. 1877-1878: 92, by monotypy. MADAGASCAR.

SCAPTESYLE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 378.

Type-species: *Scaptesyle tricolor* Walker, 1854, *ibidem* 2: 378, by monotypy. [BANGLADESH]: [Sylhet] Silhet.

‡SCAPTEZYLE; Hampson, 1901, J. Bombay nat. Hist. Soc. 13: 501.

An incorrect subsequent spelling of Scaptesyle Walker, 1854.

‡SCAPTOSYLE; Rothschild, 1912, Novit. zool. 19: 240. An incorrect subsequent spelling of Scaptesyle Walker, 1854.

‡SCAPTEZYLE; misspelling. See Scaptesyle Walker, 1854.

SCAPTIUS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 642.

Type-species: *Scaptius ditissimus* Walker, 1855, *ibidem* 3: 643, by monotypy. [Brazil]: [Amazon R.], [Tefé] Ega.

‡SCAPTOSYLE; misspelling. See Scaptesyle Walker, 1854.

SCARNICA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 42.

Type-species: *Scarnica olivina* Kiriakoff, 1962, *ibidem* (8vo Sci. zool.) **114**: 43, fig.55, by original designation. CAMEROUN: Onyanga.

SCARPONA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 123.

Type-species: Scarpona ennomoides Walker, 1862, ibidem 6:

123, by monotypy. Borneo: Sarawak. Scarpona was established in the Arct[i]idae; it was placed

in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 456.

SCEA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 293.

Type-species: *Josia auriflamma* Geyer, [1827] 1806, *in* Hübner, *Samml. exot. Schmett.* 2: pl.[186], by monotypy. Type-LOCALITY: not stated, [Brazil].

Scea was established as a subgenus of Josia Hübner, [1819].

SCEARCTIA Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 451.

Type-species: Gangamela figulina Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: xii, 55, pl.19,

fig.8, by original designation. [BRAZIL]: [Espirito Santo]

Espiritu Santo.

The date of publication of the part of Seitz containing page 451 was cited on page 449 as 20.iv.1925 in both the German and English editions.

SCEDROS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 316.

Type-species: Josia dilatata Walker, 1854, ibidem 2: 316, by

monotypy. BRAZIL.

Scedros was established as a subgenus of Josia Hübner, [1819], in the Lithosiidae, now Arctiidae; it was placed in the Cyllopodidae, now Geometridae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 401; and was subsequently placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 426.

‡SCEDROSA; Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 137.

An incorrect subsequent spelling of Scedros Walker, 1854.

APOCINESIA Bryk, 1930, in Strand, Lepid. Cat. 42: 55.

Type-species: Josia dilatata Walker, 1854, by original designation.

A junior objective synonym of Scedros Walker, 1854.

‡SCEDROSA; misspelling. See Scedros Walker, 1854. DIOP

SCELILASIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: x, 112.

Type-species: Lasiosceles erythrozonata Hampson, 1909, Ann. Mag. nat. Hist. (8) 4: 345, by original designation (for Lasiosceles Hampson, 1909). PANAMA: Chiriqui.

Scelilasia was established as an objective replacement name

for Lasiosceles Hampson, 1909.

LASIOSCELES Hampson, 1909, Ann. Mag. nat. Hist. (8) 4: 345.

Type-species: Lasiosceles erythrozonata Hampson, 1909,

ibidem (8) 4: 345, by original designation.

A junior homonym of Lasiosceles Bethune-Baker, 1906, Novit. zool. 13: 204, - Lepid., Noctuidae. The objective replacement name is Scelilasia Hampson, 1914.

SCENA Walker, 1854, List Specimens lepid. Insects Colln Br.

Type-species: Scena propylea Druce, 1894, Ann. Mag. nat. Hist. (6) 13: 170 (but included by Walker as 'Zygaena Styx? Fabr.', a misidentification; see below), by monotypy. Mexico.

Druce, 1894 (loc.cit.) considered that Walker had misidentified Zygaena styx Fabricius, 1775, Syst. Ent.: 556. Druce therefore established a new nominal species Scena propylea for this taxon.

Scena is a nominal genus having a misidentified type-species. Under the Code, Article 70(a), such cases should be referred to the Commission.

Scena was established as a subgenus of Euchromia Hübner, [1819] in the Zygaenides, now Zygaenidae; it was placed in the Syntomidae, now Ctenuchidae, by Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus.1: 281.

SCEPSIS Walker, 1854. See Cisseps Franclemont, 1936.

SCEVESIA Dyar, 1916, Proc. U.S. natn. Mus. 51: 22. NOTO Type-species: Scevesia broidricci Dyar, 1916, ibidem 51: 22, by original designation. Mexico: Hidalgo, Guerrero Mill.

SCHALIDOMITRA Strand, 1911, Dt. ent. Z. 1911: 584. NOCTUIDAE

Type-species: Schalidomitra ambages Strand, 1911, ibidem 1911: 584, by original designation. [TANZANIA] Deutsch-Ost-Afrika: [Lake] Nyassa, Kilondo-Mündung.

Schalidomitra was established in the Lymantriidae; it was transferred to the Noctuidae by C.L. Collenette and W.H.T. Tams in the collection of the BMNH; and was placed in the Noctuidae, Ophiderinae, by Nye, 1975, Generic Names Moths World 1: 441.

‡SCHALLOTOMIS; misspelling. See Schalotomis Hampson, 1920.

SCHALODETA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxviii, 801.

Type-species: Caulocera fasciolata Rothschild, 1913, Novit. zool. 20: 218, by original designation. New Guinea: [Papua New Guineal, N.E., Kumusi River.

SCHALOTOMIS Hampson, 1920, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 2: x, 123.

Type-species: Diarhabdosia roseothorax Rothschild, 1913, Novit. zool. 20: 217, by original designation. PERU: Rio Inambari, La Oroya; Carabaya, Rio Huacamayo, La Union.

‡SCHALLOTOMIS; Seitz, 1921, in Seitz, Gross-Schmett, Erde **6**: 362.

An incorrect subsequent spelling of Schalotomis Hampson, 1920.

SCHASIURA Butler, 1877, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 1: x, 37.

Type-species: Schasiura mimica Butler, 1877, ibidem 1: x, 37, pl. 16, fig. 6, by original designation. [BRAZIL]: [Amazonas], Rio Solimões, Barreiras das Araras.

SCHAUSIA Dyar, 1897. See Euschausia Dyar, 1898. ARCT

SCHAUSIADES Dognin, 1914. See Pseudhemiceras Dognin,

‡SCHAUSIANA; misspelling. See Schausilla Kiriakoff, 1974. AGARISTIDAE

SCHAUSILLA Kiriakoff, 1974, in Kiriakoff & Viette, Faune Madagascar 39: 5 (key), 97.

Type-species: Eusemia obryzos Mabille, [1879] 1878, Bull. Soc. zool. Fr. 3: 89, by original designation. Lectotype 9, MADAG[ASCAR], (BMNH) designated (as holotype) by Viette & Fletcher, 1968, Bull. Br. Mus. nat. Hist. (Ent.) 21: 417.

Schausilla in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡SCHAUSIANA; Kiriakoff, 1976, Tierreich 98: viii, 1, 125, 141, 164.

An incorrect subsequent spelling of Schausilla Kiriakoff,

SCHEDOSTAUROPUS Kiriakoff, 1960, Revue fr. Ent. 27:

Type-species: Schedostauropus elegans Kiriakoff, 1960, ibidem 27: 183, figs 9, 10, by original designation. MADAGASCAR: central, Ambatolampy district, Ampolomita.

‡SCHISTOPHLEBS; misspelling. See Schistophleps Hampson, 1891.

SCHISTOPHLEPS Hampson, 1891, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 8: 5, 53.

Type-species: Schistophleps bipuncta Hampson, 1891, ibidem 8: 54, pl.140, fig.23, by monotypy. INDIA: Madras, Nilgiri Hills.

‡SCHISTOPHLEBS; Strand, 1922, in Wagner, Lepid. Cat.

An incorrect subsequent spelling of Schistophleps Hampson, 1891.

SCHIZURA Doubleday, 1841, Entomologist 1: 59. NOTO Type-species: Schizura impomaeae Doubleday, 1841, ibidem 1: 60, by monotypy. U.S.A.: East Florida, St John's Bluff.

SCHOUTENIA Kiriakoff, 1970, Tijdschr. Ent. 113: 108.

Type-species: *Schoutenia bivittata* Kiriakoff, 1970, *ibidem* 113: 109, fig. 5, by original designation. [New Guinea]: [Irian Jaya], Waigeo Island, Camp Nok.

SCIATTA Walker, 1869, in Chapman, Proc. nat. Hist. Soc. Glasgow 1: 337.

NOCTUIDAE

Type-species: *Sciatta inconcisa* Walker, 1869, *ibidem* 1: 338, by monotypy. Type(s) &, [ZAIRE] Congo, (UM, Oxford).

Sciatta was not placed in a family when it was established; it was placed in the Arctiidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 233; it is here transferred to the Noctuidae, Ophiderinae and placed near Schalidomitra Strand, 1911.

SCIOPSYCHE Butler, 1876, J. Linn. Soc. (Zool.) 12: 426.

Type-species: *Euchromia tropica* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 236, by original designation. HONDURAS, BRAZIL.

SCLEROGENIA Ichinosé, 1973, Kontyu 41: 135. NOCTUIDAE Type-species: Plusia jessica Butler, 1878, Ann. Mag. nat. Hist. 5: 201, by original designation. Syntypes &, Japan, (BMNH).

Sclerogenia was established as a subgenus of Autographa Hübner, [1821].

Sclerogenia in the Noctuidae, Plusiinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

SCOLIACMA Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 692, 695.

Type-species: Lithosia bicolora Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 211, pl.3, fig.9, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 288. [AUSTRALIA]: Nouvelle-Hollande.

SCOLIOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xix, 453.

Type-species: Scoliacma brunnescens Rothschild, 1912, Novit. zool. 19: 215, by original designation. New Guinea: [Papua New Guinea], Angabunga River.

SCOTODONTA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 147.

Type-species: *Phalera tenebrosa* Moore, 1866, *Proc. zool. Soc. Lond.* **1865**: 815, by original designation. [INDIA]: Darjeeling.

SCOTOGRAMMA Edwards, 1887 August, Can. Ent. 19: 146.

Type-species: Scotogramma stretchii Edwards, 1887, ibidem 19: 146, by monotypy. Syntypes 28, 19, [U.S.A.]: Colorado

Scotogramma was used and made nomenclaturally available by Edwards before its proposal by Smith, [1888 January 6] 1887, not 1897 as cited in Nye, 1975, Generic Names Moths World 1: 444. Scotogramma Smith, [1888], is therefore a junior homonym; it has no objective replacement name but its type-species Anarta submarina Grote is congeneric with Scotogramma stretchii Edwards, the type-species of Scotogramma Edwards. The latter is therefore available for use as a subjective replacement name.

Scotogramma Edwards, in the Noctuidae, Hadeninae, is cited here as it was not included in Nye, 1975, *ibidem* 1.

SCOTURA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 524.

Type-species: Scotura pyraloides Walker, 1854, ibidem 2:

525, by monotypy. [BRAZIL]: Para.

S. pyraloides is a junior subjective synonym of Atolmis flavicapilla Hübner, [1819] 1806, Samml. exot. Schmett. 2: pl.[180].

SCOTUROPSIS Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 508.

Type-species: Scoturopsis basilinea Hering, 1925, ibidem 6: 508, pl.68 row k, by original designation. BOLIVIA: Rio Tanampaya.

The date of publication of the part of Seitz containing page 508 was cited on page 505 as 24.x.1925 in both the German and English editions.

SCRANCIA Holland, 1893, Psyche, Camb. 6: 536. NOTO Type-species: Scrancia modesta Holland, 1893, ibidem 6: 537, pl.17, fig.33, by original designation. [GABON]: [Ogooué] Ogové R. valley.

SCRANCIANA Strand, 1912, Arch. Naturgesch. 78 (A) 6: 161.

Type-species: Scranciana caudatissima Strand, 1912, ibidem 78 (A) 6: 162, by original designation. [EQUATORIAL GUINEA]: [Rio Muni], Benito region, Alen.

SCRANCIELLA Kiriakoff, 1967, Biologia gabon. 3: 343.

Type-species: Scranciella phryganea Kiriakoff, 1967, ibidem 3: 344, fig.30, by original designation. GABON: Bélinga.

SCRANCIOLA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 425.

Type-species: Scranciola quadripunctata Gaede, 1928, ibidem 14: 425, pl.70 row i, by monotypy. [ZAIRE] Congo: [Western Kivu, near Masisi, Middle Lowa Valley] Lowa. This additional type-locality data are taken from the label on the type specimen in the BMNH.

The date of publication of the part of Seitz containing page 425 was cited on that page as 30.iii.1928 in both the German

and English editions.

SCYTALE Felder, 1874. See Clystea Watson nom. n. CTEN

SCYTHROPHANES Turner, 1926, Proc. Linn. Soc. N.S.W. 51: 413.

Type-species: Scythrophanes stenoptera Turner, 1926, ibidem 51: 413, by monotypy. AUSTRALIA: Victoria, Inverloch.

SEBASTIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 383.

Type-species: Grotea elegans Moore, 1866, Proc. zool. Soc. Lond. 1865: 797, pl.43, fig.1, by monotypy (of Grotea Moore, 1866). [INDIA]: [N.E.], N.E. Bengal.

Sebastia was established as the objective replacement name for Grotea Moore, 1866.

G. elegans is a junior subjective synonym of Hypercompe argus Walker, 1862, Trans. ent. Soc. Lond. (3) 1:73 (but cited as ‡Hypercompa, an incorrect subsequent spelling).

GROTEA Moore, 1866, Proc. zool. Soc. Lond. 1865: 797.

Type-species: Grotea elegans Moore, 1866, by monotypy. A junior homonym of Grotea Cresson, 1864, Proc. ent. Soc. Philad. 3: 397, – Insecta, Hymenoptera. The objective replacement name is Sebastia Kirby, 1892.

MOOREA Hampson, 1894, Fauna Br. India (Moths) 2: vi, 2,

Proposed, unnecessarily, as an objective replacement name for *Grotea* Moore, 1866.

Moorea Hampson is a junior homonym of Moorea Jones & Holland, 1869, Ann. Mag. nat. Hist. (4) 3: 225, - Crustacea.

SECUSIO Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 558.

Type-species: Secusio strigata Walker, 1854, ibidem 2: 559, by monotypy. [South Africa]: [Durban] Port Natal.

‡SEIARCTIA; misspelling. See Seirarctia Packard, 1864.

SEILERIA Dognin, 1923, Hétérocères nouv. Am. Sud 23: 9.
PERI

Type-species: Seileria eucyaniformis Dognin, 1923, ibidem 23: 9, by original designation. Colombia: Villavicencio.

SEIRARCTIA Packard, 1864, Proc. ent. Soc. Philad. 3: 119.
ARCT

Type-species: *Phalaena echo* Smith, 1797, in Smith & Abbot, *Nat. Hist. rarer lepid. Insects Georgia* 2: 135, pl.68, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 242. U.S.A.: Georgia.

‡SEIARCTIA; Stretch, 1872, Illust. Zygaenidae Bombycidae N. Am. 1: 81.

An incorrect subsequent spelling of Seirarctia Packard, 1864.

SEIRODONTA Grote & Robinson, 1868, List Lepid. N. Am.: xi.

Type-species: Cecrita bilineata Packard, 1864, Proc. ent. Soc. Philad. 3: 359 (but included as 'bilineata Grote & Robinson', an incorrect authorship), by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 569. U.S.A.

SELENARCTIA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 25: 44.

Type-species: *Automolis elissa* Schaus, 1892, *Proc. zool. Soc. Lond.* **1892**: 277, by original designation. [BRAZIL]: Rio [de] Janeiro.

SELLOTA Breyer, 1956, Revta Soc. ent. argent. 19: 11. ARCT Type-species: Halysidota trifasciata Burmeister, 1878, Descr. phys. Répub. Argent. 5 (1): 44 (but cited as ‡Halesidota, an incorrect subsequent spelling), by original designation (but cited as ‡Halesidota). ARGENTINA: Buenos Aires.

SEMIDONTA Staudinger, 1892, in Romanoff, Mém. Lépid. 6: 358.

Type-species: *Drymonia biloba* Oberthür, 1880, *Etudes d'Entomologie* 5: 63, pl.8, fig.1, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

Semidonta was established as a subgenus of Allodonta Staudinger, 1887.

\$SESMODONTA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 429.

An incorrect subsequent spelling of *Semidonta* Staudinger, 1892.

SENIA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.):

Type-species: *Phalaena astur* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 35, [147], pl.120, fig.B, by monotypy. SURINAM.

P. astur is dated from the wrapper of the part and not from the title-page of the volume, 1779.

SENTANA Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 620.

Type-species: Sentana violascens Gaede, 1930, ibidem 10: 620, pl.82 row a, by monotypy. [INDIA]: North Kanara.

The date of publication of the part of Seitz containing page

620 was cited on page 617 as 10.iii.1930 in both the German and the English editions.

‡SENURA; misspelling. See Saenura Wallengren, 1858.

‡SERASANA; misspelling. See Cerasana Walker, 1862.

SERGIUSIA Nye nom. n. for Pentelia Kiriakoff, 1976.

Type-species: *Eusemia pentelia* Druce, 1887, *Proc. zool. Soc. Lond.* **1887**: 668, by original designation (for *Pentelia* Kiriakoff, 1976). Syntypes [MOZAMBIQUE]: Delagoa Bay, (BMNH).

Prof. Dr Sergius G. Kiriakoff was informed of the homonymy and has permitted me to propose the replacement name.

PENTELIA Kiriakoff, 1976, Tierreich 98: 38.

Type-species: Eusemia pentelia Druce, 1887, by original designation.

A junior homonym of *Pentelia* Brenske, 1891, *Ent. Nachr.* 17: 314, – Insecta, Coleoptera. The objective replacement name is *Sergiusia* Nye nom.n.

Pentelia in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

SERICANAPHE Kiriakoff, 1964, Bonn. zool. Beitr. 14: 292.
GEOMETRIDAE

Type-species: Sericanaphe lutea Kiriakoff, 1964, ibidem 14: 292, abb.6, figs 53, 54, pl.5, fig.56, by original designation. China: Chekiang, west Tien-mu-Shan.

Sericanaphe was established in the Notodontidae; it was transferred to the Geometridae by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 2.

SERICARIA Berthold, 1827. See Porthetria Hübner, [1819].

SERICOCHROA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.21. NOTO Type-species: Phalaena politia Stoll, 1780, in Cramer, Uit-

landsche Kapellen (Papillons exot.) 4: 41, pl.309, fig.F, (a junior primary homonym, see below), by monotypy. SURINAM.

Blera rogenhoferi Schaus, 1901, Trans. ent. Soc. Lond. 1901: 311, was established as the objective replacement name for Phalaena politia Stoll, 1780, a junior primary homonym of Phalaena politia Cramer, 1777, ibidem 2: 65, pl.139, fig.E, – Lepid., Geometridae.

BLERA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 255.

Type-species: Blera ceruroides Walker, 1862, ibidem (3) 1: 256, by monotypy. [Brazil]: Rio de Janeiro.

A junior homonym of *Blera* Billberg, 1820, *Enumeratio Insect. Mus. G.J. Billberg*: 118, – Insecta, Diptera. There is no objective replacement name, but *B. ceruroides* is a junior

no objective replacement name, but *B. ceruroides* is a junior subjective synonym of *Phalaena politia* Stoll, 1780, the typespecies of *Sericochroa* Felder, 1874; the latter is therefore available for use as a subjective replacement name.

‡BIERA; Neave, 1939, Nomencl. zool. 1: 428.

An incorrect subsequent spelling of *Blera* Walker, 1862. Neave attributed this misspelling to Dognin, 1908, *Annls Soc. ent. Belg.* **52**: 172, but at this reference *Blera* was spelt correctly.

SERINCIA Dyar, 1914, Proc. U.S. natn. Mus. 42: 168. LITH Type-species: Serincia metallica Dyar, 1914, ibidem 42: 169, by original designation. Panama: Canal Zone, Tabernilla.

SERIPHA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 441.

Type-species: Seripha caelicolor Walker, 1854, ibidem 2: 441, by monotypy. VENEZUELA.

SERMYLA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 461.

Type-species: Sermyla transversa Walker, 1854, ibidem 2: 461, by monotypy. [BRAZIL]: Rio [de] Janeiro.

SESAPA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 547.

Type-species: Sesapa inscripta Walker, 1854, ibidem 2: 547, by monotypy. China: northern.

SESIURA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xvii, 27, 371.

Type-species: Eunomia smaragdina Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 92, by original designation. [Brazil]: [Amazon R.], [Tefé] Ega.

‡SESMODONTA; misspelling. See Semidonta Staudinger, 1892.

SETINA Schrank, 1802, Fauna Boica 2 (2): 165. LITH

Type-species: Phalaena irrorella Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 535, but included as irrorea [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 68, an unjustified emendation of irrorella Linnaeus, by subsequent designation by Westwood, 1840, Introd. mod. Classif. Insects 2 (Synopsis Genera Br. Insects): 93, but designated as 'irrorella F.', an incorrect authorship.

Incorrect type-species designation: 'Irrorea' cited by Boisduval, 1836, Hist. nat. Insectes (Spec. gén. Lépid.) 1: 143. In the Introduction to the volume, pages 1-154, Boisduval reviewed earlier classifications of Lepidoptera and designated as many as three different type-species for each generic name. In his 'Exposé de notre méthode', pages 155-690, no type-species designation was made for any of the genera that he himself used. Under the Code, Article 69(a)(iii), the type-species designation of an author is eligible for consideration if he states that it is the type '... and if it is clear that he himself accepts it as the type-species'. Boisduval's type-designations, although clearly stated, do not fulfil the last requirement and so are invalid. Even though Boisduval's 1836 work was well known to lepidopterists, the type-designations contained in it have not been accepted, either by Hemming or by other authors. Acceptance of Boisduval's designation in this case would make no difference to the generic concept.

PHILEA Zetterstedt, [1839], Insecta Lapponica: 931, (but attributed to Dalman, an incorrect authorship).

Type-species: Phalaena irrorella Linnaeus, 1758 (but included as irrorea [Denis & Schiffermüller], 1775), by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 418.

A junior objective synonym of Setina Schrank, 1802.

SETINOCHROA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.16.

Type-species: Setinochroa infumata Felder, 1874, ibidem 2 (Abt.2): pl.106, fig.16, by monotypy. HIMALAYAS: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 9.

SETORA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 1069.

Type-species: Setora nitens Walker, 1855, ibidem 5: 1069,

by monotypy. JAVA.

Setora was established in the Notodontidae; it was placed in the Limacodidae by Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 412.

SEVASTOPULOA Collenette, 1960, Ann. Mag. nat. Hist. (13) 3: 91.

Typc-species: Sevastopuloa celaenocera Collenette, 1960, ibidem (13) 3: 91, by original designation. Kenya: Mombasa. SEYDELIA Kiriakoff, 1952, Bull. Annls Soc. r. ent. Belg. 88:

Type-species: Seydelia celsicola Toulgoët, 1976, Nouv. Rev. Ent. 6: 295, fig.5 (but cited as Callarctia ellioti Butler, a see below), by original designation. misidentification; RWANDA: Nyungwé forest, Wincka.

Kiriakoff, 1952, ibidem 88: 40 cited Callarctia ellioti Butler, 1895, Proc. zool. Soc. Lond. 1895: 739, pl.43, fig.6, as typespecies, but this was a misidentification of a then undescribed taxon as shown by Toulgoët, 1976, ibidem 6: 296, who established it as Seydelia turlini celsicola. This is a case of a misidentified type-species. Under the Code, Article 70(a), such cases should be referred to the Commission.

S. celsicola was established as and is currently placed as a subspecies of Seydelia turlini Toulgoët, 1976, ibidem 6: 294,

SHACHIA Matsumura, 1919, Zool. Mag. Tokyo 31: 75.

Type-species: Shachia subrosea Matsumura, 1919, ibidem 31: 75, by monotypy. JAPAN: Sapporo.

S. subrosea is a junior subjective synonym of Drymonia circumscripta Butler, 1885, Cistula ent. 3: 125.

Shachia was again proposed by Matsumura, 1920, ibidem 32: 143.

See also Microhoplitis Marumo, 1920.

SHACHIHOKA Matsumura, 1925, Zool. Mag. Tokyo 37:

Type-species: Shachihoka formosana Matsumura, 1925, ibidem 37: 403, pl.6, fig.1, by original designation. Talwan.

SHAKA Matsumura, 1920 May 15, Zool. Mag. Tokyo 32: 143.

Type-species: Asteroscopus atrovittatus Bremer, 1861, Bull. Acad. imp. Sci. St. Pétersb. 3: 483, by monotypy. [U.S.S.R.]: Amur river between confluence of Ussuri river and the Noor. See also Nagandopsis Matsumura, 1934.

BRACHIONYCOIDES Marumo, 1920 November, J. Coll. Agric. imp. Univ. Tokyo 6: 282 (key), 316.

Type-species: Asteroscopus atrovittatus Bremer, 1861, by original designation.

A junior objective synonym of Shaka Matsumura, 1920 May 15.

SHENSIPLUSIA Chou & Lu, 1974, Acta ent. sin. 17: 67, 76. NOCTUIDAE

Type-species: Shensiplusia nigribursa Chou & Lu, 1974, *ibidem* 17: 68, 76, fig.16, by original designation. Holotype ♀, CHINA: Shensi province, Wukung (Northwestern College of Agric., Shensi).

Shensiplusia in the Noctuidae, Plusiinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

‡SHIRONEA; misspelling. See Shironia Matsumura, 1925.

SHIRONIA Matsumura, 1925, J. Coll. Agric. Hokkaido imp. Univ. 15: 110.

Type-species: Shironia nivea Matsumura, 1925, ibidem 15: 110, pl.10, fig.3, by original designation. [U.S.S.R.]: [Sakhalin I.], South Saghalien, Ichinosawa.

S. nivea is a junior subjective synonym of Bombyx bicoloria [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 49.

Sec also Leucodonta Staudinger, 1892.

‡SHIRONEA; Gaede, 1934, in Strand, Lepid. Cat. 59: 202. An incorrect subsequent spelling of Shironia Matsumura,

1925.

SHISA Strand, 1917, Arch. Naturgesch. 82 (A) 1: 146.

Type-species: *Shisa excellens* Strand, 1917, *ibidem* **82** (A) 1: 146, by monotypy. [TAIWAN] Formosa: Shisa.

SICCIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 539.

Type-species: Siccia caffra Walker, 1854, ibidem 2: 539, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 368. SOUTH AFRICA.

MELANIA Wallengren, 1863, Wien. ent. Monatschr. 7: 145, 146.

Type-species: Lithosia nigropunctata Wallengren, 1860, ibidem 4: 46, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 389, but cited as Siccia caffra Walker, 1854, a nominal species not originally included in Melania. Hampson, on page 390, placed caffra as the senior synonym of nigropunctata, a nominal species originally included in Melania. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included nominal species as the type-species. [SOUTH AFRICA]: Caffraria.

A junior homonym of *Melania* Lamarck, [1799], *Mém. Soc. Hist. nat. Paris* [1]: 75. There is no objective replacement name but *L. nigropunctata* Wallengren, the type-species of *Melania* Wallengren, is a junior subjective synonym of *S. caffra* Walker, the type-species of *Siccia* Walker; the latter is therefore available for use as a subjective replacement name.

Melania was redescribed by Wallengren in 1865, K. svenska Vetensk Akad. Handl. 5 (4): 40.

SICCIAEMORPHA van Eecke, 1920, Zool. Meded. Leiden 5: 129.

Type-species: Sicciaemorpha ivyalba van Eecke, 1920, ibidem 5: 129, fig., but proposed as ‡i.v.y.-alba, an incorrect original spelling, by monotypy. Java: Preanger.

SICULIFER Hampson, 1896, Fauna Br. India (Moths) 4: 24, 497.

Type-species: Siculifer bilineatus Hampson, 1896, ibidem 4: 497, fig., by original designation. INDIA: Assam, [Khasia Hills] Khasis.

SIDANA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 1087.

Type-species: Sidana bifacies Walker, 1855, ibidem 5: 1087, by monotypy. [BRAZIL]: Rio de Janeiro.

‡SIDERIDIA; misspelling. See Sideridis Hübner, [1821].
NOCTUIDAE

SIDERIDIS Hübner, [1821] 1816, Verz. bekannter Schmett.: 232. NOCTUIDAE

Type-species: Noctua evidens Hübner, [1808], Samml. eur. Schmett. 4: pl.79, fig.369 (a primary homonym, replaced by Sideridis anapheles Nye, 1975, Generic Names Moths World 1: 450), by subsequent designation by Hampson, 1905, Cat. Lepid. Phalaenae Br. Mus. 5: 436.

Further information in Nye, 1975, ibidem 1: 449, 450.

‡SIDERIDIA; Nye, 1975, Generic Names Moths World 1: 514.

An incorrect subsequent spelling of Sideridis Hübner, [1821]. Sideridis and Sideridis anapheles were spelt correctly in the main catalogue entry on pages 449 and 450, but misspelt as ‡'Sideridia anaphelea' in the summary of new names proposed on page 514.

SIDISCA Kiriakoff, 1958, Annls Mus. r. Congo Belge (8vo Sci. zool.) 66: 64.

Type-species: Sidisca hyhochloe Kiriakoff, 1958, ibidem (8vo Sci. zool.) 66: 64, pl.3, fig.15, pl.5, fig.14, pl.9, fig.39, by original designation. [ZAIRE]: Sankuru, Lusambo.

SIDYMA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1686.

Type-species: Sidyma albifinis Walker, 1856, ibidem 7: 1686, by monotypy. [INDIA] Hindostan.

SIMAREEA Moore, 1878, Proc. zool. Soc. Lond. 1878: 14.

Type-species: *Lithosia basinota* Moore, 1866, *ibidem* **1865**: 798, by monotypy. INDIA: Darjeeling.

SIMESIA Kiriakoff, 1955, Revue Zool. Bot. afr. 52: 340.

Type-species: Stauropus dasychiroides Butler, 1898, Proc. zool. Soc. Lond. 1898: 433, pl.32, fig.12, by original designation. [Kenya]: Maziwa Mitatu [between Maungu and Buchuma, approx. 3°40'S 38°50'E].

CRITOBULUS Kiriakoff, 1963, Bull. Annls Soc. r. ent. Belg. 99: 350.

Type-species: Stauropus critobulus Fawcett, 1916, Proc. zool. Soc. Lond. 1916: 726, pl.1, fig.12, by original designation. [Kenya]: Kedai.

A junior homonym of *Critobulus* Distant, 1903, *Fauna Br. India* (Rhynchota) 2 (1): 59, 77, – Insecta, Hemiptera. There is no objective replacement name, but *S. critobulus* Fawcett, 1916, is a junior subjective synonym of *Stauropus dasychiroides* Butler, 1898, the type-species of *Simesia* Kiriakoff, 1955; the latter is thus available for use as a subjective replacement name.

SINODONTA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 5, 107 (but attributed to Matsumura, an incorrect authorship).

Type-species: Semidonta bidens Oberthür, 1914, Etud. Lépid. comp. 9 (2): 59, pl.257, fig.2163, by original designation. China: Tibet, Ta-tsien-lou district.

SIOPASTEA Schaus, 1922, Insecutor Inscit. menstr. 10: 27.

Type-species: Siopastea bakeri Schaus, 1922, ibidem 10: 28, by original designation. PHILIPPINES: Los Baños.

SIRANA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 232.

Type-species: Dasychira lygropis Collenette, 1954, Mém. Inst. scient. Madagascar (E) 5: 127, pl.6, fig.4, by original designation. MADAGASCAR: parc de Tsimbazaza.

SIRENOPYGA Wallengren, 1858, Ofvers. K. VetenskAkad. Förh. Stockh. 15: 210.

Type-species: Sirenopyga ephippiata Wallengren, 1858, ibidem 15: 210, 211, by original designation. [SOUTH AFRICA]: Caffraria.

S. ephippiata is a junior subjective synonym of Antheua simplex Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 687.

S. ephippiata was again described as a new species by Wallengren, 1860, Wien. ent. Monatschr. 4: 164.

See also Antheua Walker, 1855.

SITINA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 854.

LASIOCAMPIDAE

Type-species: *Phalaena vitulina* Donovan, 1805, *Epitome nat. Hist. Insects New Holland, New Zealand . . . Pacific Oceans*: Lepidoptera, pl.[35], fig.1, by monotypy. [AUSTRALIA]: [New South Wales], Sydney.

Sitina was established in the Liparidae, now Lymantriidae; it was placed in the Lasiocampidae by Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 267.

SITVIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 387.

Type-species: Sitvia denudata Walker, 1865, ibidem 32: 388, by monotypy. [Malaysia]: [Malaya], Malacca.

SIZALISCA Walker, 1869, in Chapman, Proc. nat. Hist. Soc. Glasgow 1 (2): 339.

Type-species: Sizalisca graminosa Walker, 1869, ibidem 1

(2): 339, by monotypy. Congo.

Sizalisca was established in the Liparidae, now Lymantriidae; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 559.

SKAPHITA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 260 (key), 295.

Type-species: *Phya salona* Druce, 1894, *Ann. Mag. nat. Hist.* (6) 13: 356, by original designation. MEXICO: Coatepec.

SMINTHOPSES Hampson, 1926, Descr. new Genera Species Lepid. Phalaenae Subfamily Noctuinae Br. Mus.: 528. LYMA Type-species: Sminthopses aeschra Hampson, 1926, ibidem: 528, by original designation. UGANDA: Entebbe.

Sminthopses was established in the Noctuidae; it was placed in the Lymantriidae by Collenette, 1935, Stylops 4: 242.

SNELLENITA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 60.

Type-species: *Phalera divaricata* Gaede, 1930, in Seitz, *Gross-Schmett. Erde* 10: 614, pl.81 row e (but attributed to Snellen, an incorrect authorship), by original designation. [INDIA]: Khasia Hills, JAVA, SUMATRA.

SNELLENOPSIS van Eecke, 1920, Zool. Meded. Leiden 5: 132.

Type-species: Snellenopsis mimetica van Eecke, 1920, ibidem 5: 132, fig., by monotypy. JAVA: W., Preanger.

SOGANACLIA Griveaud, 1964, Faune Madagascar 17: 105.

Type-species: Soganaclia viridisparsa Griveaud, 1964, ibidem 17: 107, text-figs 255-257, pl.1, fig.72, by original designation. MADAGASCAR: E., Andapa, Ambatosoratra.

SOLOE Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 557. AGAN

Type-species: *Soloe trigutta* Walker, 1854, *ibidem* **2**: 557, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* **1**: 429. [ZAIRE] Congo, SIERRA LEONE.

Soloe was established in the Lithosiidae, now Arctiidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 429; and transferred from the 'Liparidae' to the 'Aganaidae' by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497.

SOLOELLA Gaede, 1926, in Seitz, Gross-Schmett. Erde 14: 90.

Type-species: Soloe guttivaga Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 558, by monotypy. [GHANA]: Ashanti.

The date of publication of the part of Seitz containing page 90 was cited on page 81 as 9.vi.1926 in both the German and English editions.

SOLWEZIA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 439.

Type-species: Solwezia multilineata Gaede, 1928, ibidem 14: 439, pl.72 row d, by monotypy. [ZAMBIA]: Solwezi.

The date of publication of the part of Seitz containing page 439 was cited on page 433 as 30.v.1928 in the German edition; and as 26.xi.1928 in the English edition.

SOMATOXENA Aurivillius, 1904, Ark. Zool. 2 (4): 66.

Type-species: Xenosoma lasea Druce, 1899, Ann. Mag. nat. Hist. (7) 3: 468, by original designation. SIERRA LEONE.

‡PIRGALASEA; Swinhoe, 1923, Ann. Mag. nat. Hist. (9) 11: 86.

‡*Pirgalasea* was an incorrect subsequent spelling of *Pirga lasea* (Druce) but was cited as a generic name by Bryk, 1934, *in* Strand, *Lepid. Cat.* 62: 302, and by Neave, 1940, *Nomencl. zool.* 3: 768.

SOMATRICHIA Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 274.

Type-species: *Phalaena parasita* Hübner, 1790, *Beitr. Gesch. Schmett.* **2** (2): 42, pl.2, fig.I, by subsequent designation of *Trichosoma* Rambur, 1832. Type-locality: not stated, [Europe].

Somatrichia was established as the objective replacement name for Trichosoma Rambur, 1832.

TRICHOSOMA Rambur, 1832, Annls ent. Soc. Fr. (1) 1: 272. Type-species: Phalaena parasita Hübner, 1790, by subsequent designation by Blanchard, 1840, in Castelnau, Hist. nat. Insectes 3: 487, but cited for ‡Trichosomum Rambur, an incorrect subsequent spelling.

A junior homonym of *Trichosoma* Rudolphi, 1819, *Entozoorum Synopsis*: 13, – Nematoda. The objective replacement

name is Somatrichia Kirby, 1892.

Trichosoma Rambur was placed by Ferguson, 1978, in Dominick, Moths Am. N. of Mexico 22 (2): 58, as a junior subjective synonym of Orgyia Ochsenheimer, 1810, in the Lymantriidae, on the basis of an incorrect type-species.

‡TRICHOSOMUM; Blanchard, 1840, in Castelnau, Hist. nat.Insectes 3: 487.

An incorrect subsequent spelling of Trichosoma Rambur, 1832.

‡TRICHIOSOMA; Neave, 1940, Nomencl. zool. 4: 538.

An incorrect subsequent spelling of *Trichosoma* Rambur, 1832.

Neave attributed this misspelling to Lucas, 1849, Explor. scient. Algérie (Zool.) 3: 376, but no error had been made by Lucas.

SOMENA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1734.

Type-species: Somena scintillans Walker, 1856, ibidem 7: 1734, by monotypy. INDIA: North.

SOMERA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 882.

Type-species: Somera viridifusca Walker, 1855, ibidem 4: 882, by monotypy. [BANGLADESH]: Sylhet.

Somera was established in the Liparidae, now Lymantriidae; it was placed in the Notodontidae by Butler, 1880, Ann. Mag. nat. Hist. (5) 6: 67.

SOMERINA Kiriakoff, 1965, Bull. Annls Soc. r. ent. Belg. 101: 325.

Type-species: *Somerina arcuata* Kiriakoff, 1965, *ibidem* **101**: 325, fig.17, by monotypy. CAMEROUN: Kribi.

SOMEROPSIS Strand, 1912, Arch. Naturgesch. 78 (A) 6: 160.

Type-species: Someropsis viriditincta Strand, 1912, ibidem 78 (A) 6: 160, by original designation. [EQUATORIAL GUINEA]: [Rio Muni], Benito region, Makomo, Alen.

SOMMERIA Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 13.

Type-species: Sommeria culta Hübner, [1831] 1825, ibidem 3: 13, pl. [75], figs 433, 434, by monotypy. AFRICA: [southern].

SONTIA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1641.

Type-species: *Phalaena neuroptera* Sepp, [1848], *Surinaamsche Vlinders* (1): 83, pl.38, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 201. SURINAM.

P. neuroptera is a junior subjective synonym of Sphinx arontes Stoll, 1782, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 184, 248, pl.382, fig. E.

See also Pelochyta Hübner, [1819].

SORAMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 980 (key), 1034.

Type-species: *Sorama bicolor* Walker, 1855, *ibidem* 5: 1034, by monotypy. [Australia]: Tasmania.

SOREMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1064.

LASIOCAMPIDAE

Type-species: Sorema nubila Walker, 1855, ibidem 5: 1065, by PRESENT DESIGNATION. [AUSTRALIA]: [New South Wales], Sydney.

Sorema was established in the Notodontidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 841.

SOREUTONEURA Collenette, 1930, Trans. ent. Soc. Lond. 78: 68.

Type-species: Soreutoneura daedala Collenette, 1930, ibidem 78: 68, pl.6, fig.9, by original designation. NEW GUINEA: [Irian Jaya] Dutch N.G., Weyland Mts, Mt Kunupi, Menoo valley.

SORITENA Schaus, 1924, Proc. U.S. natn. Mus. 65 (7): 32.

Type-species: Soritena habanera Schaus, 1924, ibidem 65 (7): 32, by original designation. CUBA: Havana.

SOROCABA Moore, 1882, in Jones, Proc. lit. phil. Soc. Lpool 36: 331, 354.

Type-species: Sorocaba anomala Moore, 1882, ibidem 36: 331, 354, pl.6, fig.15, by monotypy. BRAZIL: [São Paulo] Santo Paulo.

S. anomala is a junior subjective synonym of Ceratonyx carmelitaria Guenée, 1857, the type-species of Ceratophora Guenée, 1858. See below.

Sorocaba was established in the Sphingidae; it was placed in the Lymantriidae by Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 559.

CERATOPHORA Guenée, 1858, in Boisduval & Guenée, Hist. nat. Insectes (Spec. gén. Lépid.) 9 (Atlas): 3, pl.3.

Type-species: Ceratonyx carmelitaria Guenée, 1857, ibidem 9: 194; 1858, ibidem 9 (Atlas): pl.3, fig.6, by monotypy. BRAZIL: [Nova Friburgo] Nouvelle-Fribourg.

A junior homonym of *Ceratophora* Gray, [1835] 1833-1834, *Illust. Indian Zool.* 2: pl.68, fig.2, — Reptilia. There is no objective replacement name, but *C. carmelitaria* is a senior subjective synonym of *Sorocaba anomala* Moore, 1882, the type-species of *Sorocaba* Moore, 1882; the latter is therefore available for use as a subjective replacement name.

Ceratophora Guenée, 1858, was established in the Geometridae; and was transferred to the Lymantriidae by Fletcher, 1979, Generic Names Moths World 3: 192.

SOZUSA Wallengren, 1863, Wien. ent. Monatschr. 7: 146.

Type-species: *Lithosia scutellata* Wallengren, 1860, *ibidem* 4: 45, by original designation. [SOUTH AFRICA]: Caffraria, eastern.

Sozusa was redescribed as new by Wallengren in 1865, K. svenska Vetensk Akad. Handl. 5 (4): 42.

‡SOZUZA; Butler, 1878, Ann. Mag. nat. Hist. (5) 2: 457.
An incorrect subsequent spelling of Sozusa Wallengren, 1863.

‡SOZUZA; misspelling. See Sozusa Wallengren, 1863.

SPATALIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 145. NOTO

Type-species: *Bombyx argentina* [Denis & Schiffermüller], 1775, *Ankündung syst. Werkes Schmett. Wienergegend*: 62, 249, pl.1a, fig.2, pl.2a, fig.2, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 604. [Austria]: Vienna district.

HETERODONTA Duponchel, [1845] 1844, Cat. méth. Lépid. Eur. (2): 92.

Type-species: *Bombyx argentina* [Denis & Schiffermüller], 1775, by monotypy.

A junior objective synonym of Spatalia Hübner, 1822.

SPATALINA Bryk, 1950, Ark. Zool. 42 (A) 19: 34. NOTO Type-species: Spatalina birmalina Bryk, 1950, ibidem 42 (A) 19: 35, pl.2, fig.5, by original designation. Burma: Kambaiti.

S. birmalina was established as a subspecies of Lophopteryx argentata Moore, (1879), in the combination Spatalina argentata (Moore) ssp. birmalina Bryk. Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 233 raised birmalina to specific rank.

XEROPTERYX Kiriakoff, 1964, Bonn. zool. Beitr. 14: 289.

Type-species: Xeropteryx desiccata Kiriakoff, 1964, ibidem 14: 289, text-fig.50, pl.5, fig.3, by original designation. China: northern Yunnan, Li-Kiang.

A junior homonym of Xeropteryx Butler, 1883, J. Linn. Soc. (Zool.) 17: 197, 203, — Lepid., Geometridae. There is no objective replacement name, but X. desiccata was shown by Kiriakoff, 1968, in Wytsman, Genera Insect. 217°(C): 234, to be congeneric with Spatalina birmalina Bryk, 1950, the typespecies of Spatalina Bryk, 1950; the latter is thus available for use as a subjective replacement name.

SPATALOIDES Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 36.

Type-species: Spatalia dives Oberthür, 1884, Etudes d'Entomologie 10: 15, pl.1, fig.1, by original designation. [U.S.S.R.]: [S. of Vladivostok], Sidemi.

SPATULOSIA Toulgoët, 1965, Bull. Soc. ent. Fr. 70: 226.

'Type-species: Spatulosia malgassica Toulgoët, 1965, ibidem 70: 227, text-fig.5, pl.5, fig.7, by original designation. MADAGASCAR: E., near Perinet, Analamazaotra forest.

SPHAEROMACHIA Grote, 1867, Proc. ent. Soc. Philad. 6: 304.

Type-species: *Pericopis cubana* Herrich-Schäffer, 1866, *CorrespBl. zool.-min. Ver. Regensburg* **20**: 131, by monotypy. Cuba.

‡SPHECODES Herrich-Schäffer, [1858]. See Eumenogaster Herrich-Schäffer, 1855.

SPHECOMIMAX Bryk, 1953, Ark. Zool. (2) 5: 255. CTEN Type-species: Pseudosphex aurifera Klages, 1906, Proc. U.S. natn. Mus. 29: 531, by original designation. Venezuela: S., Suapure.

‡SPHECOMORPHA Hübner, [1806]. See Isanthrene Hübner, [1819].

‡SPHĖCOMORPHA Hübner, 1808. See Pseudosphex Hübner, 1818.

‡SPHECOMORPHA Hübner, 1818. See Pseudosphex Hübner, 1818. CTEN

This name has not been used again in the Lepidoptera. It was later established by Newman, 1838, *Ent. Mag.* 5: 396, in the Coleoptera.

SPHECOPS Orfila, 1935, Revta Soc. ent. argent. 7: 181.

Type-species: *Pseudosphex arctata* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 95, by original designation. [BRAZIL]: [Amazon R.], [Tefé] Ega.

SPHECOPSIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.29. CTEN

Type-species: *Sphecopsis hyalozona* Felder, 1874, *ibidem* 2 (Abt.2): pl.102, fig.29, by monotypy. [Colombia]: Bogota, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 9.

SPHECOPSYCHE Dognin, 1898, Annls Soc. ent. Belg. 42: 344.

Type-species: Sphecopsyche julumito Dognin, 1898 October 5, ibidem 42: 344, by monotypy. [ECUADOR]: Loja.

S. julumito is a junior subjective synonym of Chrysostola sanguiceps Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 405.

SPHECOSOMA Butler, 1876, J. Linn. Soc. (Zool.) 12: 381.

Type-species: Sphecosoma fasciolatum Butler, 1876, ibidem 12: 381, by original designation. [BRAZIL]: [Matto Grosso], [Serra de] Santa Martha.

S. fasciolatum is a junior subjective synonym of Pseudosphex cognata Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1619.

SPHENOPTERA Felder, 1874. See Asphenoptera Watson nom. n. CTEN

SPHETTA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 457.

Type-species: *Sphetta apicalis* Walker, 1865, *ibidem* **32**: 457, by monotypy. [SRI LANKA] Ceylon.

Sphetta was established in the Notodontidae; it was placed in the Noctuidae by Hampson, 1894, Fauna Br. India (Moths) 2: 166

SPHRAGIDIUM Butler, 1887, Ann. Mag. nat. Hist. (5) 19: 218.

Type-species: *Sphragidium miles* Butler, 1887, *ibidem* (5) 19: 218, by original designation. Solomon Islands: Alu.

SPHRAGISTA Collenette, 1934, Ann. Mag. nat. Hist. (10) 13: 219.

Type-species: *Laelia kitchingi* Bethune-Baker, 1909, *Ann. Mag. nat. Hist.* (8) 3: 429, by original designation. [UGANDA]: Acholi Country, [Fatiko] Patigo.

SPILARCTIA Butler, 1875, Cistula ent. 2: 39. ARCT

Type-species: *Phalaena lutea* Hufnagel, 1766, *Berlin. Mag.* 2: 412, (but included by Butler as *lubricipeda* Linnaeus, a misidentification; see below), by subsequent designation by Kirby, 1877, *in* Rye, *Zool. Rec.* 12: 431, (but cited as *P. lubricipeda* Linnaeus). GERMANY: Berlin.

Butler, 1875, included *Phalaena lubricipeda* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 505, in *Spilarctia*, and Kirby, 1877, designated it as the type-species. Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 227, indicated that the true *lubricipeda* Linnaeus, 1758, was the senior subjective synonym of *Phalaena lubricipeda alba* Hufnagel, 1776, and also of *Bombyx men-*

thastri [Denis & Schiffermüller], 1775, (the type-species of Spilosoma Curtis, 1825). Kirby, 1892, ibidem 1: 229, indicated that Phalaena lubricipeda lutea Hufnagel, 1766, was the valid name for lubricipeda Linnaeus, 1758, var. beta, and for lubricipeda sensu Butler, 1875.

Spilarctia Butler is a case of a genus that has a misidentified type-species and so should be referred to the Commission under the Code, Article 70(a).

P. lutea was established as a subspecies of Phalaena lubricipeda Linnaeus.

See also Olistheria Hübner, [1820].

SPILARCTIA Staudinger, 1891. See Nebrarctia Watson nom.

SPILOBOTYS Butler, 1887, Ann. Mag. nat. Hist. (5) 20: 123.

AGAN

Type-species: *Spilobotys arctioides* Butler, 1887, *ibidem* (5) **20**: 123, by monotypy. Solomon Islands: [Guadalcanal] Guadalcanar.

SPILOSOMA Curtis, 1825, Br. Ent. 2: pl.92.

Type-species: Bombyx menthastri [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 54, by original designation. [AUSTRIA]: Vienna district.

B. menthastri is a junior subjective synonym of Phalaena lubricipeda Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 505.

The name *Spilosoma* was attributed to Stephens but was used and made nomenclaturally available by Curtis prior to its proposal by Stephens, 1827

SPILOSOMA Stephens, 1827, in Anonymous, Retrospective Review (2) 1: 242.

Type-species: *Phalaena lubricipeda* Linnaeus, 1758, *Syst. Nat.* 1 (Edn 10): 505, by monotypy. Type-Locality: not stated, [EUROPE].

A junior homonym of *Spilosoma* Curtis, 1825. There is no objective replacement name but *P. lubricipeda* is the senior subjective synonym of *Bombyx menthastri* [Denis & Schiffermüller], 1775, the type-species of *Spilosoma* Curtis, 1825; the latter is therefore available for use as a subjective replacement name.

Spilosoma was proposed again by Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 55, 74; 1829 [June], Nom. Br. Insects: 40; and 1829 [July], Syst. Cat. Br. Insects (2): 56.

SPILOSOMA Stephens, 1827. See Spilosoma Curtis, 1825.

SPIRIS Hübner, [1819] 1816, Verz. bekannter Schmett.: 169.

Type-species: *Phalaena grammica* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 822, by monotypy. GERMANY.

P. grammica is a junior subjective synonym of Phalaena striata Linnaeus, 1758, ibidem 1: 502.

CALLOPIS Billberg, 1820, Enumeratio Insect. Mus. G.J. Billberg: 91.

Type-species: *Phalaena grammica* Linnaeus, 1758, by PRES-ENT DESIGNATION.

A junior objective synonym of Spiris Hübner, [1819].

Invalid type-species designation by Tams, 1939, Entomologist 72: 136 who cited Phalaena striata Linnaeus a nominal species not originally included in Callopis and not linked in synonymy by Tams with the originally included nominal species.

EULEPIA Curtis, 1825, Br. Ent. 2: 56.

Type-species: *Phalaena grammica* Linnaeus, 1758, by original designation.

A junior objective synonym of Spiris Hübner, [1819].

CTENIA Le Peletier, 1825, Encycl. méth. 10 (Insectes): 650. Type-species: Phalaena grammica Linnaeus, 1758, by monotypy (but attributed to Latreille, an incorrect authorship).

A junior objective synonym of Spiris Hübner, [1819].

EMYDIA Boisduval, 1828, Eur. Lepid. Index meth.: 39.

Type-species: *Phalaena grammica* Linnaeus, 1758, by susequent designation by Blanchard, 1840, *in* Castelnau, *Hist. nat. Insectes* 3: 489.

A junior objective synonym of Spiris Hübner, [1819].

SPODARCTIA Dyar, 1925, Insecutor Inscit. menstr. 13: 3.

Type-species: *Spodarctia superpulverea* Dyar, 1925, *ibidem* 13 RP 3, by original designation. MEXICO: Colima.

SPODIOSOMERA Kiriakoff, 1960, Revue fr. Ent. 27: 183.

Type-species: *Spodiosomera oculata* Kiriakoff, 1960, *ibidem* 27: 184, figs 11, 12, by original designation. MADAGASCAR: eastern, Sambava district, Marojejy massif.

STAETHERINIA Butler, 1878, Trans. ent. Soc. Lond. 1878: 73.

Type-species: Oxytenis semilutea Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1942, by original designation. [BRAZIL]: Para.

STAUROPIDA Kiriakoff, 1962, Tijdschr. Ent. 105: 249.

Type-species: Stauropida griseola Kiriakoff, 1962, ibidem 105: 250, fig.15, by original designation. [Zambia]: Ndola.

STAUROPLITIS Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 639.

Type-species: Stauroplitis annulata Gaede, 1930, ibidem 10: 639, pl.84 row g, by monotypy. SUMATRA: Barisan Mts.

The date of publication of the part of Seitz containing page 639 was cited on page 633 as 21.ii.1930 in the German edition; and as 17.v.1930 in the English edition.

STAUROPODOPSIS Roepke, 1944, Natuurh. Maandbl. 33: 20.

Type-species: Stauropodopsis grisescens Roepke, 1944, ibidem 33: 21, figs 22a-22d, by original designation. JAVA: eastern, Djunggo-Ardjuno.

STAUROPUS Germar, 1812, Diss. sistens Bombycum Species (2): 45.

Type-species: *Phalaena fagi* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 508, by monotypy. TYPE-LOCALITY: not stated, [EUROPE].

TERASION Hübner, [1819] 1816, Verz. bekannter Schmett.: 147.

Type-species: *Phalaena fagi* Linnaeus, 1758, by monotypy. A junior objective synonym of *Stauropus* Germar, 1812.

‡TERASIA; Agassiz, 1846, Nomencl. zool. (Nom. syst. Generum Anim.) Lepid.: 64; 1847, ibidem (Index univl.): 364.

An incorrect subsequent spelling of *Terasion* Hübner, [1819]. The Lepidoptera part of *Nomencl. zool*. (Nom. syst. Generum Anim.) was included in fascicles 9 and 10 issued in one wrapper dated 1846. The 'Index universalis' formed fascicle 12 of the work; the wrapper is dated 1847, the title-page is dated 1846.

STAUROPUSSA Strand, 1912, Arch. Naturgesch. 78 (A) 6: NOTO

Type-species: Stauropussa viridipennis Strand, 1912, ibidem 78 (A) 6: 170, by monotypy. Cameroun: Bibundi.

S. viridipennis is a junior subjective synonym of Somera chloe Holland, 1893, Psyche, Camb. 6: 487.

STEMATOCAMPA Packard, 1895, Mem. natn. Acad. Sci. 7 (Bombycine Moths Am. N. of Mexico 1): 223.

Type-species: *Lochmaeus unicolor* Packard, 1864, *Proc. ent. Soc. Philad.* 3: 373, by monotypy. U.S.A.: Massachusetts, Seekonk, Cambridge, Rhode Island.

Stematocampa was established as a subgenus of Heterocampa Doubleday, 1841.

See also Misogada Walker, 1865.

STEMMATOPHALERA Aurivillius, 1910 [December 31], in Sjöstedt, Wiss. Ergebn. schwed. zool. Exped. Kilimandjaro, Meru . . . 1905-1906 (9): 20.

Type-species: *Stemmatophalera sjostedti* Aurivillius, 1910 [December 31], *ibidem* (9): 20, pl.1, fig.3, by monotypy. [TANZANIA]: Kilimandjaro.

S. sjostedti is a junior subjective synonym of Chadisra semiflava Hampson, 1910 May, Ann. Mag. nat. Hist. (8) 5: 462.

S. sjostedti is based on the Swedish name Sjöstedt, not on a German word and does not require more than the deletion of the diacritic mark under the Code, Article 32(c)(i).

STEMONOCERAS Karsch, 1895, Ent. Nachr. Berlin 21: 370.

Type-species: *Phaegorista leucomela* Herrich-Schäffer, [1854] 1850-1858, *Samml. neuer oder wenig bekannter aussereur. Schmett.* 1 (1): wrapper, pl.5, figs 22, 23, by original designation (but cited as *leucomelas*, a mandatory change in spelling). [SENEGAL]: Casamanca.

STENADONTA Hampson, 1895, Trans. ent. Soc. Lond. 1895: 281.

Type-species: *Stenadonta cyttarrosticta* Hampson, 1895, *ibidem* 1895: 281, fig., by original designation. BHUTAN.

STENARCHA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 89, 264.

Type-species: Chiriphe stenopa Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 732, by original designation. AUSTRALIA: Western Australia, Albany.

STENARCTIA Aurivillius, 1899, Ent. Tidskr. 20: 236, 244.

Type-species: *Stenarctia quadripunctata* Aurivillius, 1899, *ibidem* **20**: 244, fig., by monotypy. [Gabon]: [Ogooué] Ogowe, Africa: 'Kuilu'.

STENAROA Hampson, 1910, Ann. Mag. nat. Hist. (8) 5: 444.

Type-species: Stenaroa ignepicta Hampson, 1910, ibidem (8) 5: 444, by original designation. MADAGASCAR: Betsileo.

STENAULIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: ix, 81, 124.

Type-species: Lithosia discalis Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 108, by original designation. BORNEO: Sarawak.

STENELOPSIS Butler, 1877, Trans. ent. Soc. Lond. 1877: 375.

Type-species: Stenelopsis exposita Butler, 1877, ibidem 1877: 375, by original designation. [BRAZIL]: Para.

STENEUGOA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxiv, 639.

Type-species: Steneugoa nubilosa Hampson, 1914, ibidem (Suppl.) 1: 639, fig.196, by original designation. Syntypes 28, 29, New Guinea: [Irian Jaya] Dutch New Guinea, Wataikwa River, (BMNH).

Steneugoa was established in the Arctiadae, now Arctiidae; it was transferred to the Noctuidae, Sarrothripinae by Holloway, 1979, Series Ent. 15: 448.

Steneugoa in the Noctuidae, Sarrothripinae was not included in Nye, 1975, Generic Names Moths World 1.

STENILEMA Hampson, 1909, Ann. Mag. nat. Hist. (8) 4:

Type-species: Stenilema aurantiaca Hampson, 1909, ibidem (8) 4: 353, by original designation. ETHIOPIA.

STENOCYTTARA Hulstaert, 1923. See Onymapata Hulstaert, 1924.

‡STENOGNATHA; misspelling. See Sthenognatha Felder,

STENOPHAEA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xi, 6, 215.

Type-species: Ctenucha pollinia Boisduval, 1870, Considérations Lépid. Guatemala: 83, by original designation. GUATEMALA.

STENOPLASTIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, figs 16, 17.

Type-species: Stenoplastis satyroides Felder, 1874, ibidem 2 (Abt.2): pl.105, fig.16, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 342. [COLOMBIA]: Bogota, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 9.

Stenoplastis when established was not placed in a family; it was placed in the Lithosiidae, now Arctiidae, by Kirby, 1892, ibidem 1: 342; and was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 412.

STENOPTEROSIA Joannis, 1928, Annls Soc. ent. Fr. 97: 266.

Type-species: Stenopterosia angustipennis Joannis, 1928,

ibidem 97: 266, by original designation. [VIETNAM]: Tonkin, Cho ganh.

STENOPYDNA Roepke, 1943, Natuurh. Maandbl. 32: 94.

Type-species: Stenopydna lanceolata Roepke, 1943, ibidem 32: 94, fig.12, by monotypy. Type-locality: not known. Thought by Roepke to be JAVA: western, ?Perbawattee.

Stenopydna was established in the Notodontidae; it was transferred to the Noctuidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2; and was transferred to the Pyralidae, Crambinae by Nye, 1975, Generic Names Moths World 1: 459, as a junior subjective synonym of Schoenobiodes Hampson, 1917, Novit. zool. 24: 57.

STENOSCAPTIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 83, 280.

Type-species: Mosoda venusta Lucas, 1890, Proc. Linn. Soc. N.S.W. (2) 4: 1078, by original designation. AUSTRALIA: Brisbane.

STENOSHACHIA Matsumura, 1925, Zool. Mag. Tokyo 37:

Type-species: Stenoshachia bipartita Matsumura, 1925, ibidem 37: 398, pl.6, fig.7, by original designation. TAIWAN.

STENOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: viii, 82, 92. LITH

Type-species: Stenosia rufeola Hampson, 1900, ibidem 2: 93, fig.26, by original designation. [INDONESIA]: Bali [I.].

STENOSPATALIA Matsumura, 1924, Trans. Sapporo nat. Hist. Soc. 9: 36.

Type-species: Spatalia jezoensis Wileman & South, 1916, Entomologist 49: 133, by original designation. JAPAN: Hokkaido, Toshima Province, Tobetsu.

STENOSTAURA Hampson, 1909, Trans. zool. Soc. Lond. 19: 119. NOTO

Type-species: Cossus impeditus Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 583, by monotypy. [South Africa]: Cape [Province].

STENOSTAURIDIA Strand, 1912, Arch. Naturgesch. 78 (A) 6: 171.

Type-species: Stenostauridia comma Strand, 1912, ibidem 78 (A) 6: 172, by original designation. CAMEROUN: Bibundi.

STENUCHA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xvi, 8, 450.

Type-species: Ctenucha dolens Druce, 1897, Biologia cent.am. (Zool.) Lepid.-Heterocera 2: 360, pl.73, fig.4, by original designation. MEXICO.

See also Zamolis Dyar, 1910, and Ardonipsa Dyar 1912.

STERRHOSIA Joannis, 1928, Annls Soc. ent. Fr. 97: 268.

Type-species: Sterrhosia zonata Joannis, 1928, ibideni 97: 268, by original designation. [VIETNAM]: Tonkin, Cho ganh.

STHENADELPHA Turner, 1922, Proc. Linn. Soc. N.S.W. 47:

Type-species: Trichetra isabella White, 1841, in Grey, Jls two Exped. Disc. N.W. and W. Australia . . . 1837, 38, and 39 2 (Appendix F): 479, figs 10, 11, by monotypy. AUSTRALIA: [Western Australia], King George Sound.

STHENOGNATHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.101, fig.9.

Type-species: Sthenognatha gentilis Felder, 1874, ibidem 2 (Abt.2): pl.101, fig.9, by monotypy. TYPE-LOCALITY: not stated by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 9, but a female syntype in the BMNH is labelled 'Brasilia' [BRAZIL].

‡STENOGNATHA; Felder, 1875, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): Erklärung to pls 75-107: 9.

An incorrect subsequent spelling of Sthenognatha Felder, 1874.

STICTANE Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiii, 88, 258.

Type-species: Pitane fractilinea Snellen, [1880], in Veth, Midden-Sumatra 4 (2): 38, by original designation. SUMATRA:

STICTOGARGETTA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 27.

Type-species: Gargetta albimacula Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 136, by original designation. [INDIA]: Sikkim.

‡STICTONACLES; misspelling. See Stictonaclia Hampson, 1898. CTEN

STICTONACLIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xi, 23, 125.

Type-species: Naclia anastasia Oberthür, 1893, Etudes d'Entomologie 18: 2, pl.1, fig.8, by original designation. [MAD-AGASCAR]: Antsianaka.

‡STRICTONACLES; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling (cited as ‡Stictonacles by Neave, 1940, Nomencl. zool. 4: 313) of Stictonaclia Hampson,

‡STICTONACLES; Neave, 1940, Nomencl. zool. 4: 313.

An incorrect subsequent spelling (see ‡Strictonacles; Pagenstecher, 1909) of Stictonaclia Hampson, 1898.

STICTOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 88, 293.

Type-species: Conchylis flexilisana Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 362, by original designation. BORNEO: Sarawak.

STIDZAERAS Druce, 1905, Ann. Mag. nat. Hist. (7) 15: 467.

Type-species: Stidzaeras strigifera Druce, 1905, ibidem (7) 15: 467, by monotypy. VENEZUELA: Caura Valley, Corosita.

STIGMATOPHORA Staudinger, 1881, Stettin. ent. Ztg 42: 399.

Type-species: Setina micans Bremer & Grey, 1853, in Motschulsky, Etudes entomologiques 1: 63, by monotypy. [CHINA]: [Peking] Pekin region.

STIGMATOPHORINA Mell, 1922, Dt. ent. Z. 1922: 122.

Type-species: Stigmatophorina hammamelis Mell, 1922, ibidem 1922: 122, by subsequent designation by Gaede, 1930, in Seitz, Gross-Schmett. Erde 10: 608. China: northern Kwangtung province.

STILPNAROMA Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 133.

Type-species: Stilpnaroma venosa Hering, 1926, ibidem 14: 133, pl.26, row h, by original designation. [MALAWI] Britisch Zentral-Afrika: Blantyre, Luchonza.

The date of publication of the part of Seitz containing page 133 was cited on page 129 as 1.xi.1926 in both the German and the English editions.

STILPNOTIA Westwood, 1843. See Leucoma Hübner, 1822.

STONIA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 187.

Type-species: Stonia bipars Walker, [1865] 1864, ibidem 31: 187, by monotypy. [AUSTRALIA]: Moreton Bay.

STRACENA Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 388.

Type-species: Stracena fuscivena Swinhoe, 1903, ibidem 1903: 388, by original designation. [NIGERIA]: River Niger, Sapele.

STRACILLA Aurivillius, 1910, in Sjöstedt, Wiss. Ergebn. schwed. zool. Exped. Kilimandjaro, Meru . . . 1905-1906 (9): 22.

Type-species: *Leucoma translucida* Oberthür, 1880, *Annali Mus. civ. Stor. nat. Genova* 15: 177, pl.1, fig.6, by monotypy. [Етнюріа] Abissinia.

Stracilla was incorrectly attributed to 'Hering, 1926, in Seitz' by Bryk, 1934, in Strand, Lepid. Cat. 62: 187.

STREPSIMANES Meyrick, 1930, Exot. Microlepid. 4: 10.

Type-species: Strepsimanes scieropis Meyrick, 1930, ibidem 4: 10, by monotypy. [INDIA]: [Uttar Pradesh], [Kumaun] Kumaon, [Mukteswar] Muktesar.

Strepsimanes was established as the type-genus of the Strepsimanidae Meyrick, 1930, in the Microlepidoptera. The family was transferred to the Noctuoidea by Hodges, 1978, in Dominick et al., Moths Am. N. of Mexico 6 (1): 11.

STREPTOPHLEBIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: [ix], 22, 52.

Type-species: Streptoplebia obliquistria Hampson, 1898, ibidem 1: 52, fig.23, by original designation. Borneo: [Sabah], Kina Balu.

‡STRICTONACLES; misspelling. See Stictonaclia Hampson, 1898.

STRIOSIA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xix, 452.

Type-species: *Calamidia irrorata* Rothschild, 1912, *Novit. zool.* 19: 223, by original designation. New Guinea: [Irian Jaya], central, Mt Goliath.

STROPHOCERUS Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 344.

Type-species: Strophocerus flocciferus Möschler, 1883, ibidem 32 (Abh.): 345, pl.18, fig.33, by monotypy. Surinam: interior.

STRUBA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 152 (key), 169.

Type-species: Bireta argenteodivisa Kiriakoff, 1962 [July 31], ibidem 98 (10): 170; 1962 October 20, Bonn. zool. Beitr. 13: 225, abb.1, fig.7, pl.1, fig.6 (as Bireta argenteodivisa sp.nov.), by original designation. China: north Yunnan, [Li-Kiang]. Additional data cited by Kiriakoff, 1962 October 20, ibidem 13: 225.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr. to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper B. argenteodivisa was described as a new species. As the dates show, the Belgian paper appeared first.

In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

Struba was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 42.

SUAREZIA Hering, 1926. See Salvatgea Griveaud, 1977.

SUBNIGANDA Kiriakoff, 1962 [July 31], Bull. Annls Soc. r. ent. Belg. 98 (10): 153 (key), 170; 1962 October 20, Bonn. zool. Beitr. 13: 222.

Type-species: Bireta aurantiistriga Kiriakoff, 1962 [July 31], ibidem 98 (10): 171; 1962 October 20, ibidem 13: 223, abb.1, fig.4, pl.1, fig.4 (as Subniganda aurantiistriga sp.nov.), by original designation. China: south Shensi, [Tapaishan im Tsinling]. Additional data cited by Kiriakoff, 1962 October 20, ibidem 13: 223.

Kiriakoff evidently expected his paper in the Bonn. zool. Beitr. to appear before that in the Bull. Annls Soc. r. ent. Belg., for in the Bonn paper B. aurantiistriga was described as a new species in the combination Subniganda aurantiistriga, and Subniganda was proposed as a new genus. As the dates show, the Belgian paper appeared first.

In a personal communication, the secretary of the Société Royale d'Entomologie de Belgique stated that fascicle 10 of volume 98 of the Society's journal was despatched to them from the printer on 30 June 1962; he added that fascicles are available in the Society's library and in the Royal Library of Belgium about 15 days after leaving the printer. In the absence of a precise date of publication, I have adopted the date 31 July 1962 for fascicle 10 of volume 98.

Subniganda was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1962 October 20, ibidem 13: 222, and again in 1967, in Wytsman, Genera Insect. 217 (B): 42.

SUBSCRANCIA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 425.

Type-species: Scrancia nigra Aurivillius, 1904, Ark. Zool. 2 (4): 7, by monotypy. Togo.

The date of publication of the part of Seitz containing page 425 was cited on that page as 30.iii.1928 in both the German and the English editions.

SUBWILEMANUS Kiriakoff, 1964, Bonn. zool. Beitr. 14: 281.

Type-species: Subwilemanus pictus Kiriakoff, 1964, ibidem 14: 282, abb.5, figs 41, 42, pl.4, fig.44, by original designation. CHINA: Chekiang, west Tien-mu-Shan.

SUMERIA Tams, 1938, Proc. R. ent. Soc. Lond. (B) 7: 144.

Type-species: Sumeria dipotamica Tams, 1938, ibidem (B) 7: 144, figs 1-7, by original designation. [IRAQ]: [Al Faw] Fao.

‡SUNGATA; misspelling. See Sangata Walker, 1860. NOTO

SURETA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1073.

Type-species: Sureta tripars Walker, 1855, ibidem 5: 1074, by monotypy. [BRAZIL]: Rio de Janeiro.

SURIGA Kiriakoff, 1962, Bull. Annls Soc. r. ent. Belg. 98: 153 (key), 208.

Type-species: Turnaca suriga Schaus, 1928, Proc. U.S. natn. Mus. 73 (19): 88, by original designation. Philippines: Mindanao I., Surigao.

Suriga was established as a subgenus of Bireta Walker, 1856; it was raised to generic level by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 94.

SUTONOCREA Butler, 1876 September, J. Linn. Soc. (Zool.) 12: [vii] (Corrigenda et Addenda); 1876 [May], ibidem: 419.

Type-species: Creatonotos lobifer Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.88, fig.503 (but cited as ‡Creatonotus, an incorrect subsequent spelling by Herrich-Schäffer), by monotypy (but included by Butler, 1876 September, as see below). [SOUTH AMERICA]: [tropical] Am. aequin., cited by Herrich-Schäffer, [1856], ibidem 1 (1): wrapper to pl.88.

Herrich-Schäffer, [1856], illustrated as fig.503 and named, on the wrapper of the Heft containing this figure, his new species lobifer. Later, in the text to the same work, [1858], ibidem 1: 72, he placed lobifer as a junior synonym of Malabus incertus Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 716, but these have since been treated as distinct species.

Butler, 1876 September, in a Corrigenda et Addenda established Sutonocrea for the species he had cited on page 419 as 'Creatonotus incertus, Herrich-Schäffer, Auss. Schmett. i. fig. 503'. This misidentification incertus Walker sensu Herrich-Schäffer, [1858], had already been named by Herrich-Schäffer himself as lobifer.

This is a case of a misidentified type-species which under the Code, Article 70(a), should be referred to the Commission.

Sutonocrea was established in the Zygaenidae; it was placed in the Arctiidae by Druce, 1884, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 79.

SUZUKIA Matsumura, 1920. See Suzukiana Sugi, 1976. NOTO

SUZUKIANA Sugi, 1976, Kontyu 44: 287. NOTO

Type-species: Suzukia suzukii Matsumura, 1920, Zool. Mag. Tokyo 32: 140, by original designation (for Suzukia Matsumura, 1920). JAPAN: Takao-san, Kyoto, [Nagano pref.], Oiwake, Hokkaido.

S. suzukii is a junior subjective synonym of Peridea cinerea Butler, 1879, Ann. Mag. nat. Hist. (5) 4: 353.

Suzukiana was established as the objective replacement name for Suzukia Matsumura, 1920.

SUZUKIA Matsumura, 1920, Zool, Mag. Tokyo 32: 140.

Type-species: Suzukia suzukii Matsumura, 1920, by original designation.

A junior homonym of Suzukia Okamoto, 1913, Trans. Sapporo nat. Hist. Soc. 4 (2): 109, - Insecta, Plecoptera. The objective replacement name is Suzukiana Sugi, 1976.

Suzukia was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 768.

SYBRIDA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 465. PYRALIDAE.

Type-species: Sybrida inordinata Walker, 1865, ibidem 32:

466, by monotypy. [INDIA]: Darjeeling.

Sybrida was established in the Notodontidae; it was placed in the Pyralidae by Snellen, 1890, Trans. ent. Soc. Lond. 1890:

SYCHESIA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 653.

Type-species: Sychesia fimbria Möschler, 1878, ibidem 27 (Abh.): 654, by monotypy. Surinam: interior of.

S. fimbria is a junior subjective synonym of Phalaena dryas Cramer, 1775, Uitlandsche Kapellen (Papillons exot.) 1: 110, pl.70, fig.C.

SYCHNACEDES Collenette, 1953, Ann. Mag. nat. Hist. (12) LYMA

Type-species: Sychnacedes idiopis Collenette, 1953, ibidem (12) 6: 562, pl.13, fig.22, by original designation. MADAGAS-CAR: Antsianaka.

SYLESCAPTIA van Eecke, 1920, Zool. Meded. Leiden 5: 134.

Type-species: Sylescaptia tigrina van Eecke, 1920, ibidem 5: 134, fig., by monotypy. JAVA: Ambarawa; and Semarang.

‡SYMIRE; misspelling. See Lymire Walker, 1854.

SYMMERISTA Hübner, [1821] 1816, Verz. bekannter Schmett.: 248.

Type-species: Noctua albicosta Hübner, [1809] 1796, Samml. eur. Schmett. 4: pl.93, fig.440, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 572. [NORTH AMERICA]: mistakenly included by Hübner as an European species; N. albicosta occurs in Canada, the U.S.A. and Mexico.

N. albicosta is a form of Phalaena albifrons Smith, 1797, in Smith & Abbot, Nat. Hist. rarer lepid. Insects Georgia 2: 159, pl.80.

See also Edema Walker, 1855.

SYMMETRODES Meyrick, 1887, Proc. Linn. Soc. N.S.W. (2) 1: 692, 703.

Type-species: Symmetrodes sciocosma Meyrick, 1888, ibidem (2) 2: 920, by monotypy (but cited as Lithosia nitens Walker, a misidentification; see below). AUSTRALIA: Queensland.

Symmetrodes was established for one species Lithosia nitens Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 231. After his redescription Meyrick added 'In this instance I have identified Walker's species from description only, and it is possible I may be mistaken'. The following year Meyrick, 1888, established S. sciocosma as a new species for his misidentification L. nitens Walker sensu Meyrick, 1887, and added 'I propose this name for the species which I identified from Walker's description as his Lithosia nitens; having since seen his specimens, I find the identification to be erroneous . . .'.

This is a case of misidentified type-species which under the *Code*, Article 70(a), should be referred to the Commission.

SYMPHLEBIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.1. ARCT

Type-species: Symphlebia lophocampoides Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.1, by monotypy. BRAZIL: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 9.

SYMPHLEBOMIS Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 425.

Type-species: *Syntomoides antipolo* Semper, 1898, *Reisen Archipel Philipp. Semper* (2) **6**: 418, pl.53, figs 2, 3, by original designation. Philippines: central Luzon.

SYNALAMIS Dognin, 1912, Hétérocères nouv. Am. Sud 6: 18. NOCTUIDAE

Type-species: *Synalamis brunneoviridans* Dognin, 1912, *ibidem* 6: 18, by monotypy. Syntypes 18, 19, COLOMBIA: near San Antonio, Alto de las Cruces, (USNM, Washington).

Synalamis in the Noctuidae, Ophiderinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

1.

SYNETE Kiriakoff, 1959, Bull. Annls Soc. r. ent. Belg. 95: 173.

Type-species: Synete albipunctella Kiriakoff, 1959, ibidem 95: 173, fig.19, by original designation. [ZAIRE] Congo Belge: Sankuru, Lusambo.

SYNOGDOA Aurivillius, 1904, Ark. Zool. 2 (4): 52. LYMA Type-species: Synogdoa simplex Aurivillius, 1904, ibidem 2 (4): 53, fig.40, by monotypy. [Cameroun] Kamerun: N'dian.

SYNORTHODES Franclemont, 1976, Proc. ent. Soc. Wash. 78: 409.

Type-species: Synorthodes auriginea Franclemont, 1976, ibidem 78: 411, figs 2, 6, 7, 12, 19, 20, by original designation. Holotype 3, [U.S.A.]: Arizona, Cochise Co., Chiricahua Mts, Pinery Canyon, (Colln J.G. Franclemont, Ithaca).

Synorthodes, in the Noctuidae, Hadeninae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

SYNTARCTIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: ix, 4, 98.

Type-species: *Halysidota aenone* Butler, 1878, *Trans. ent. Soc. Lond.* 1878: 50 (but cited as ‡*Halesidota*, an incorrect subsequent spelling), by original designation (but cited as *H. testacea* Möschler, a misidentification; see below). South America: Amazon Basin.

The original designation by Hampson of Halysidota testacea Möschler, 1877, Verh. zool.-bot. Ges. Wien (Abh.) 27: 668 (but cited as ‡Halisidota), as type-species of Syntarctia, was an error as S. testacea Möschler sensu Hampson, 1901, was a misidentification of Halysidota aenone Butler, 1878, as shown by Hampson, 1920, ibidem (Suppl.) 2: 211 (but cited as ‡oenone, an incorrect subsequent spelling). This is a case of a misidentified type-species which should be referred to the Commission under Article 70(a) of the Code.

‡SYNTAXIS Hering, 1926. See Leucoperina Aurivillius, 1909.

‡SYNTHOMIS; misspelling. See Syntomis Ochsenheimer, 1808.

SYNTOMA Billberg, 1820. See Syntomis Ochsenheimer, 1808.

‡SYNTOMEDIA; misspelling. See Syntomeida Harris, 1839.

SYNTOMEIDA Harris, 1839, Am. J. Sci. 36: 316. CTEN Type-species: Glaucopis ipomoeae Harris, 1839, ibidem 36: 316, by monotypy. [U.S.A.]: Georgia, Savannah.

Syntomeida was established as a subgenus of Glaucopis Fabricius, 1807, in the 'Zygaenides', now Zygaenidae; it was placed in the Euchromiidae, now Ctenuchidae by Neumoegen & Dyar, 1893, Jl N.Y. ent. Soc. 1: 101.

See also Hippola Walker, 1854.

‡SYNTOMEDIA; Druce, 1884, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 42.

An incorrect subsequent spelling of Syntomeida Harris, 1839.

SYNTOMEREA Strand, 1912, Arch. Naturgesch. 78 (A) 6: CTEN

Type-species: Syntomerea typica Strand, 1912, ibidem 78 (A) 6: 177, by original designation. [EQUATORIAL GUINEA]: Benito [R.] region.

Seitz, 1926, Gross-Schmett. Erde 14: 48 cited S. typica Strand as an aberration (and therefore a junior subjective synonym) of Epitoxis myopsychoides Strand, 1912, ibidem 78 (A) 6: 182.

SYNTOMIDOPSIS Hering, 1925, in Seitz, Gross-Schmett. Erde 6: 452.

Type-species: Eudule variegata Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 380, by monotypy. JAMAICA.

The date of publication of the part of Seitz containing page 452 was cited on page 449 as 20.iv.1925 in both the German and English editions.

SYNTOMIMORPHA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xii, 92, 254.

Type-species: Syntomimorpha caerulescens Hampson, 1900, ibidem 2: 255, fig.170, by original designation. [SULAWESI] Celebes.

SYNTOMIS Ochsenheimer, 1808, Schmett. Eur. 2: 104.

Type-species: *Sphinx phegea* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 494, by monotypy. Germany, Syria.

‡GLAUCOPIS Hübner, [1806], Tentamen determinationis digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) Opinion 97: 19, also ibidem, 1954, Opin. Decl. int. Commn zool. Nom. 6(Opinion 278): 140.

Only included species Sphinx phegea Linnaeus, 1758.

SYNTOMA Billberg, 1820, in Enumeratio Insect. Mus. G.J. Billberg, 82.

An unjustified emendation of Syntoniis Ochsenheimer, 1808.

‡SYNTHOMIS; Mabille, 1892, Bull. Soc. ent. Fr. 1892: cxxxix.
An incorrect subsequent spelling of Syntomis Ochsenheimer, 1808.

SYNTOMOIDES Hampson, [1893] 1892, Fauna Br. India (Moths) 1: xiv, 209.

Type-species: *Sphinx imaon* Cramer, 1779, *Uitlandsche Kapellen (Papillons exot.)* 3: 94, 175, pl.248, fig.E, by original designation. Surinam.

S. imaon is dated from the wrapper of the part and not from the title-page of the volume, 1782.

SYNTOMOSTOLA Dognin, 1912, Hétérocères nouv. Am. Sud 5: 11. ARCT

Type-species: Syntomostola xanthosoma Dognin, 1912, ibidem 5: 11, by original designation. Colombia: eastern, Medina.

SYNTRICHURA Butler, 1876, J. Linn. Soc. (Zool.) 12: 405.

Type-species: Syntrichura virens Butler, 1876, ibidem 12: 405, by original designation. [Brazil]: [São Paulo] St Paulo.

SYNTYPISTIS Turner, 1907, Proc. Linn. Soc. N.S.W. 31: 679.

Type-species: Syntypistis chloropasta Turner, 1907, ibidem 31: 679, by monotypy. AUSTRALIA: Queensland, Kuranda.

SYRRUSOIDES Laporte, 1972, Bull. Soc. ent. Fr. 77: 243.

Type-species: Syrrusoides viettei Laporte, 1972, ibidem 77: 244, figs 7, 9, by original designation. Holotype & MADAGASCAR: Ambatofinandrahana, (MNHN, Paris).

Syrrusoides in the Noctuidae, Acronictinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

SYSTROPHA Hübner, [1819]. See Eilema Hübner, [1819].

TACHUDA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 259 (key), 278.

Type-species: Lochmaeus albosigma Druce, 1887, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 236; 1898, ibidem 2: pl.90, fig.17, by original designation. Guatemala: Volcan de Atitlan.

TADANA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 982 (key), 990.

Type-species: *Tadana cinerascens* Walker, 1855, *ibidem* 5: 991, by monotypy. Type-locality: not known. The species is now known to occur in Canada and the eastern U.S.A.

T. cinerascens is a junior subjective synonym of Lochmaeus manteo Doubleday, 1841, Entomologist 1: 58.

See also Lochmaeus Doubleday, 1841.

TAENIOPTERYX Janse, 1920. See Metopteryx Fletcher nom.

TAGELA Schaus, 1901, *Trans. ent. Soc. Lond.* **1901**: 258 (key), 265.

Type-species: *Symmerista dentata* Schaus, 1892, *Proc. zool. Soc. Lond.* **1892**: 335, by original designation. [BRAZIL]: Rio de Janeiro.

TAIWA Kiriakoff, 1967, Tijdschr. Ent. 110: 51. NOTO Type-species: Stauropus confusa Wileman, 1910, Entomologist 43: 289, by original designation. TAIWAN: Rantaizan.

TAKADONTA Matsumura, 1920, Zool. Mag. Tokyo 32: 147.

Type-species: *Takadonta takamukui* Matsumura, 1920, *ibidem* 32: 147, by monotypy. JAPAN: [Nagano pref.], Kamikochi.

Takadonta was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 797, but was incorrectly spelt as ‡Fusadonta.

‡FUSADONTA; Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 797.

An incorrect subsequent spelling of *Takadonta* Matsumura, 1920.

TAKASHACHIA Matsumura, 1929, Insecta matsum. 4: 46.

Type-species: *Takashachia maculosa* Matsumura, 1929, *ibidem* 4: 46, pl.1, fig.13, by original designation. [TAIWAN] Formosa: Horisha.

Takashachia was established in the Notodontidae; it was transferred to the Lymantriidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

TAKWA Kiriakoff, 1957, Tijdschr. Ent. 100: 96. THYRET Type-species: Takwa xenops Kiriakoff, 1957, ibidem 100: 96, by original designation. [GHANA] Gold Coast: Takwa.

TALARA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1891.

Type-species: *Talara megaspila* Walker, 1866, *ibidem* 35: 1891, by monotypy. [Brazil]: [Amazon R.], [Tefé] Ega.

TALMECA Schaus, 1905, Proc. U.S. natn. Mus. 29: 262.

Type-species: *Talmeca perplexa* Schaus, 1905, *ibidem* **29**: 262, by original designation. French Guiana: Maroni River, West Indies: Trinidad.

TALMENIA Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 345.

Type-species: *Talmenia arsilonchoides* Möschler, 1883, *ibidem* **32** (Abh.): 346, pl.18, fig.34, by monotypy. Surinam: Paramaribo.

‡TALYCOPTERA; misspelling. See Thylacoptera Hampson, 1898.

TAMPEA Snellen, 1898, Tijdschr. Ent. 40: 226. LITH Type-species: Tampea lithosioides Snellen, 1898, ibidem 40: 226, by original designation. JAVA: W., Tjisolok.

T. lithosioides is a junior subjective synonym of Lithosia reversa Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 105.

See also Ocrosia Hampson, 1900.

TAMSEALE Nye, 1975, Generic Names Moths World 1: 470.

Type-species: *Eleale plusioides* Walker, 1862, *J. Proc. Linn. Soc.* (Zool.) 6: 135, by monotypy (of *Eleale* Walker, 1862). Borneo: Sarawak.

ELEALE Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 135.

Type-species: *Eleale plusioides* Walker, 1862, by monotypy. A junior homonym of *Eleale* Newman, 1841, *Entomologist* 1: 26, – Insecta, Coleoptera. The objective replacement name is *Tamseale* Nye, 1975.

Eleale was established in the Notodontidae; it was placed in the Noctuidae by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 85.

TAMSITA Kiriakoff, 1954, Revue Zool. Bot. afr. 50: 185.

Type-species: *Turuptiana ochthoeba* Hampson, 1920, *Cat. Lepid. Phalaenae Br. Mus.* (Suppl.) 2: 474, pl.69, fig.7, by original designation. [ZAMBIA]: N.W. Rhodesia, Solwezi.

TANADA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1745.

Type-species: *Tanada antica* Walker, 1856, *ibidem* 7: 1745, by monotypy. [U.S.A.]: Georgia.

T. antica is a junior subjective synonym of Cycnia tenera Hübner, 1818, Zuträge Samml. exot. Schmett. 1: 7, pl.[1], figs 3, 4.

See also Cycnia Hübner, 1818.

TANAOSTYLA Warren, 1897, Novit. zool. 4: 422. DIOP Type-species: Tanaostyla dilucida Warren, 1897, ibidem 4: 422, by original designation. COLOMBIA: Popayan.

T. dilucida is a junior subjective synonym of Monocreagra pheloides Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.105, fig.8.

Tanaostyla was established in the Geometridae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 415.

See also Monocreagra Felder, 1874.

TANCREA Püngeler, 1898, Societas ent. 13: 57.

Type-species: *Tancrea pardalina* Püngeler, 1898, *ibidem* 13: 57, by monotypy. [U.S.S.R.]: [Kazakhstan], Ili.

TANYSTOLA Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 370.

Type-species: Cnethocampa ochrogutta Herrich-Schäffer, [1856] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.80, fig.460, by monotypy. AUSTRALIA.

Turner, 1922, *ibidem* 47: 370, suggested that the type-locality was probably in Western Australia.

TARIKA Moore, 1878, Proc. zool. Soc. Lond. 1878: 14. LITH Type-species: Lithosia varana Moore, 1865, ibidem 1865: 797, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 322. INDIA: Darjeeling.

TARSOLEPIS Butler, 1872, Ann. Mag. nat. Hist. (4) 10: 125.

Type-species: *Tarsolepis remicauda* Butler, 1872, *ibidem* (4) 10: 125, pl.8, by original designation. JAVA: Batavia.

T. remicauda is a junior subjective synonym of Crino sommeri Hübner, [1821] 1806, Samml. exot. Schmett. 2: pl.[197].

CRINO Hübner, [1821] 1806, Samml. exot. Schmett. 2: pl.[197].

Type-species: Crino sommeri Hübner, [1821] 1806, ibidem 2: pl.[197], by monotypy. Type-locality: not stated. The series of the nominate subspecies in the BMNH includes specimens from India, Burma, Malaysia, Indonesia and the Philippine Islands.

A junior homonym of *Crino* Lamarck, [1798], in Virey, J. Phys. Chim. Hist. nat. 4 [47]: 429; 1801, Syst. Anim. sans Vert.: 339, — Vermes. There is no objective replacement name, but C. sommeri is a senior subjective synonym of Tarsolepis remicauda Butler, 1872, the type-species of Tarsolepis Butler, 1872; the latter is therefore available for use as a subjective replacement name.

PIKA Matsumura, 1929, Insecta matsum. 4: 79 (key), 80.

Type-species: Tarsolepis taiwana Wileman, 1910, Entomologist 43: 138, by original designation. TAIWAN: Rantaizan.

A junior homonym of *Pika* Lacepède, [1799], *Discours d'Ouverture et de Cloture du Cours d'Hist. nat. . . . l'An VII de la République* (Tableau des Divisions, Sous-divisions, Ordres et Genres des Mammifères): 9, – Mammalia. There is no objective replacement name but *T. taiwana* is congeneric with *Tarsolepis remicauda* Butler, 1872, the type-species of *Tarsolepis* Butler, 1872; the latter is thus available for use as a subjective replacement name.

TARSOLEPISOIDES Nakamura, 1976, Trans. lepid. Soc. Japan 27: 35.

Type-species: Tarsolepis japonica Wileman & South, 1917, Entomologist 50: 29, by original designation. Japan: Tokyo. Tarsolepisoides was established as a subgenus of Tarsolepis

Butler, 1872.

TATARGINA Butler, 1877, Trans. ent. Soc. Lond. 1877: 366.

Type-species: *Deiopeia picta* Walker, [1865] 1864, *List Specimens lepid. Insects Colln Br. Mus.* 31: 263, by original designation. [Burma]: [Moulmein] Moulmeine.

TAXILA Walker, 1855. See Dysschema Hübner, 1818. PERI

TEARA Walker, 1855 [August], List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 846.

Type-species: *Teara contraria* Walker, 1855 [August], *ibidem* 4: 849, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 468. AUSTRALIA.

T. contraria is a junior subjective synonym of Ochrogaster lunifer Herrich-Schäffer, [1855 June], the type-species of Ochrogaster Herrich-Schäffer, [1855 June].

Teara was established in the Liparidae, now Lymantriidae; it was transferred to the Eupterotidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 498; and was placed in the Cnethocampinae, now Thaumetopoeidae, by Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 371.

See also Ochrogaster Herrich-Schäffer, [1855 June].

TEARINA Kiriakoff, 1970, in Wytsman, Genera Insect. 219 (E): 40.

Type-species: *Epicoma nigrolineata* Joicey, Noakes & Talbot, 1916, *Trans. ent. Soc. Lond.* 1915 (3, 4): 385, pl.61, fig.14, by original designation. New Guinea: [Irian Jaya], Arfak Mts, Angi Lakes.

TEAROSOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt. 2): pl.100, fig.6. LYMA Type-species: Tearosoma aspersum Felder, 1874, ibident 2 (Abt.2): pl.100, fig.6, by monotypy. Type-locality: not known by Felder, 1875, ibidemt 2 (Abt.2) Erklärung to pls 75-107: 9, [SOUTH AFRICA]: [Cape province].

TECMESSA Burmeister, 1878, Descr. phys. Répub. Argent. 5: 504.

Type-species: *Thosea annulipes* Berg, 1878, *An. Soc. cient. argent.* 5: 186, by monotypy. Argentina: Buenos Aires. See also *Eunaduna* Dognin, 1901.

TEGULATA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 110.

Type-species: *Tegulata tumida* Walker, 1862, *ibidem* 6: 110, by monotypy. Borneo: Sarawak.

TEIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 4: 778 (key), 803.

Type-species: *Teia anartoides* Walker, 1855, *ibidem* 4: 804, by monotypy. Australia: Sydney and [Tasmania] Van Dieman's Land.

TEINOCLADIA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.96, fig.9.

GEOMETRIDAE

Type-species: *Teinocladia cuculloides* Felder, 1874, *ibidem* 2 (Abt.2): pl.96, fig.9, by monotypy. [Australia]: Tasmania, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 9.

Teinocladia when established was not placed in a family; it was placed in the Notodontidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 930; and it was transferred to the Geometridae by Turner, 1903, Proc. Linn. Soc. N.S.W. 28: 75.

TEINOMASTYX Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxviii, 799.

Type-species: *Chamaita griseobasis* Rothschild, 1913, *Novit. zool.* **20**: 219, by original designation. New Guinea: [Papua New Guinea], Mambare R., Biagi.

TEINOPHALERA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 59.

Type-species: *Phalera elongata* Rothschild, 1917, *Novit. zool.* **24**: 253, pl.7, fig.17, by original designation. [India]: Assam, Khasia Hills.

TELECLITA Turner, 1903, *Proc. Linn. Soc. N.S.W.* **28**: 45 (key), 53.

Type-species: *Teleclita cydista* Turner, 1903, *ibidem* **28**: 53, by monotypy. Australia: Queensland, Townsville.

In the BMNH there is a short series of the type-species of *Teleclita* from Queensland, reared by F.P. Dodd; two of the specimens are labelled '*Notodonta dryinopa* Lower' in Dodd's

107: 9.

handwriting, one is labelled 'Hoplitis cydista Turn.' in Turner's handwriting. Teleclita cydista was placed in the genus Hoplitis Hübner, [1819], by Turner in 1922, Proc. Linn. Soc. N.S.W. **47**: 380.

No reference has been found to the publication by Lower of the name Notodonta dryinopa, but a larva and a pupa were described under that name by Dodd, 1902, Entomologist 35: 42, 193. Dodd's description of the larva of Notodonta dryinopa and Turner's description of the similar larva of Teleclita cydista in 1903, Proc. Linn. Soc. N.S.W. 28: 54, together with the evidence of the labelled moths, suggest strongly that the two are synomynous. Teleclita cydista Turner, 1903, is therefore here treated as a junior subjective synonym of Notodonta dryinopa Dodd (Lower MS), 1902, syn. n.

TELIONEURA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.31. CTEN Type-species: Telioneura glaucopis Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.31, by monotypy. [Brazil]: Amazon R., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-

TEUCER Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 165.

An unnecessary objective replacement name for Telioneura Felder, 1874, which is not preoccupied by *Telejoneura* Rondani, 1864, – Insecta, Diptera.

TENSHA Matsumura, 1925, Zool. Mag. Tokyo 37: 392.

Type-species: Tensha striatella Matsumura, 1925, ibidem 37: 392, pl.7, fig.8, by original designation. Talwan.

TENUINACLIA Griveaud, 1964, Faune Madagascar 17: 62.

Type-species: Micronaclia oberthueri Rothschild, 1911, Novit. zool. 18: 155, by original designation. MADAGASCAR: Antananariyo.

TEPILIA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1158 (key), 1176.

Type-species: Tepilia biluna Walker, 1855, ibidem 5: 1176, by monotypy. Brazil.

Tepilia was established in the Drepanulidae, now Drepanidae; it was placed in the Lymantriidae by Schaus, 1927, in Seitz, Gross-Schmett. Erde 6: 556.

See also Phecada Walker, 1856.

TERACOTONA Butler, 1878, Proc. zool. Soc. Lond. 1878: 382.

Type-species: Aloa rhodophaea Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 302, by original designation. [Botswana]: Lake Ngami.

MACRONYX Felder, 1874, in Feler & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.100, fig.22.

Type-species: Macronyx debilis Felder, 1874, ibidem 2 (Abt.2): pl.100, fig.22, by monotypy. [GUYANA]: Demerara, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 6.

A junior homonym of Macronyx Swainson, 1827, Zool. J. 3: 344, - Aves. There is no objective replacement name but the type-species of Macronyx is currently considered to be congeneric with the type-species of Teracotona Butler, 1878.

M. debilis is a junior subjective synonym of Virbia submacula Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 472.

See also Microgiton Felder, 1874.

TERAGRA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 1064. Type-species: Teragra conspersa Walker, 1855, ibidem 5: 1064, by monotypy. [South Africa]: [Natal], [Durban] Port

Teragra was established in the Notodontidae; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1, placed it in the Notodontidae on page 614 and in the Zeuzeridae, now Cossidae, on page 865; it was placed in the Arbelidae, now Metarbelidae, by Hampson, 1910, Ann. Mag. nat. Hist. (8) 6: 117, 125.

‡TERASIA; misspelling. See Stauropus Germar, 1812. NOTO

TERASION Hübner, [1819]. See Stauropus Germar, 1812.

TERATOPORA Meyrick, 1889, Trans. ent. Soc. Lond. 1889: 459.

Type-species: Teratopora haplodes Meyrick, 1889, ibidem 1889: 459, by monotypy. New Guinea: [Papua New Guinea], Port Moresby.

TERMESSA Newman, 1856 April 21, Trans. ent. Soc. Lond. (2) 3 (8): 285 (but attributed to Walker).

Type-species: Termessa shepherdi Newman, 1856 April 21, ibidem (2) 3 (8): 285, fig., by monotypy. AUSTRALIA: Victoria.

The name Termessa originated from Walker but was used and made nomenclaturally available by Newman prior to its proposal as a name for a new genus by Walker, 1856 [May 10], List Specimens lepid. Insects Colln Br. Mus. 7: 1689.

TERMESSA Walker, 1856 [May 10], List Specimens lepid. Insects Colln Br. Mus. 7: 1689.

Type-species: Termessa laeta Walker, 1856, ibidem 7: 1689,

by monotypy. [AUSTRALIA]: Tasmania; Sydney.

A junior homonym of Termessa Newman, 1856 April 21. There is no objective replacement name but the type-species of these two nominal genera are currently considered to be congeneric so that Termessa Newman, 1856 April 21 is available for use as a subjective replacement name.

TERMESSA Walker, 1856 [May 10]. See Termessa Newman, 1856 April 21.

TERNA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 267.

Type-species: Terna ampla Walker, [1865] 1864, ibidem 31: 267, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 399. [BRAZIL]: [Amazon R.], [Tefé] Ega.

TERPHOTHRIX Holland, 1893, Psyche, Camb. 6: 474. LYMA

Type-species: Terphothrix lanaria Holland, 1893, ibidem 6: 474, 568, pl.18, fig.19, by original designation. [GABON]: [Ogooué] Ogové R. valley.

‡TERPHOTRIX; Bryk, 1934, in Strand, Lepid. Cat. 62: 312. An incorrect subsequent spelling of Terphothrix Holland, 1893.

‡TERPHOTRIX; misspelling. See Terphothrix Holland, 1893.

TESMA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 48.

Type-species: Phryganopsis nigrapex Strand, 1912, Arch. Naturgesch. 78 (A) 7: 173, by original designation. CAMERoun: northern, Johann-Albrechts-Höhe.

TESSELLA Breyer, 1957, Revta Soc. ent. argent. 19: 14.

Type-species: Halysidota sertata Berg, 1882, An. Soc. cient. argent. 13: 214, by original designation. ARGENTINA: Buenos Aires province.

TESSELLARCTIA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 6, 90.

Type-species: Halysidota semivaria Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1707 (but cited as ‡Halesidota, an incorrect subsequent spelling), by original designation (but cited as \$\frac{1}{2}Halesidota). Brazil.

TESSELLOTA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: xv, 7, 430.

Type-species: Halysidota cancellata Burmeister, Descr. phys. Répub. Argent. 5 (1): 445, (but cited as ‡Halesidota, an incorrect subsequent spelling), by original designation (but cited as \$\frac{1}{2}Halesidota). Argentina: Buenos Aires.

TESSMANNIA Bryk, 1915, Arch. Naturgesch. 81 (A) 4: 1. LYMA

Type-species: Tessmannia braueri Bryk, 1915, ibidem 81 (A) 4: 2, fig.4, by original designation. [EQUATORIAL GUINEA]: [Rio Muni] Spanisch-Guinea, Uelleburg.

TEUCER Kirby, 1892. See Telioneura Felder, 1874.

TEULISNA Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 109. LITH

Type-species: Teulisna plagiata Walker, 1862, ibidem (Zool.) 6: 109, by original designation. Borneo: Sarawak.

THACONA Walker, [1863] 1864, J. Proc. Linn. Soc. (Zool.) 7: 169.

Type-species: Thacona costivitta Walker, [1863], ibidem

(Zool.) 7: 169, by monotypy. Borneo: Sarawak.

Thacona was established in the Hypogrammidae, now Noctuidae; it was placed in the Notodontidae by Swinhoe, 1900, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 2: 300.

THAGONA Möschler, 1883, Verh. zool.-bot. Ges. Wien 32 (Abh.): 337.

Type-species: Thagona uniformis Möschler, 1883, ibidem 32 (Abh.): 338, by monotypy. Surinam: Paramaribo.

THAILA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 414.

Type-species: Thaila cinerascens Kiriakoff, 1974, ibidem 17: 415, text-fig.28, pl.2, fig.5, by original designation. THAI-LAND: Khzo Yzi.

THALESA Schaus, 1896, Jl N.Y. ent. Soc. 4: 136. Type-species: Charidea seruba Herrich-Schäffer, [1855] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.51, fig.280 (but attributed to Boisduval, an incorrect authorship), by original designation.

C. seruba is a junior subjective synonym of Phalaena citrina Sepp, [1852], Surinaamsche Vlinders 2: [113].

THALLARCHA Meyrick, 1886, Proc. Linn. Soc. N.S.W. (2) 1: 692, 736.

Type-species: Thallarcha phalarota Meyrick, 1886, ibidem (2) 1: 736, by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 499. AUSTRALIA: New South Wales, Sydney; Victoria, Beechworth.

THAMBETA Collenette, 1953, Ann. Mag. nat. Hist. (12) 6:

Type-species: Thambeta haigi Collenette, 1953, ibidem (12) 6: 577, pl.13, fig.8, by original designation. NIGERIA: S., Obudu.

THAMNOCERA Holland, 1893, Psyche, Camb. 6: 454, 469.

Type-species: Thamnocera albilinea Holland, 1893, ibidem

6: 469, 568, pl.18, fig.7, by original designation. [GABON]: [Ogooué] Ogové R. valley.

THANATARCTIA Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 395.

Type-species: Thanatarctia infernalis Butler, 1877, ibidem (4) 20: 395, by original designation. JAPAN: Hakodate.

THAUMATOPOEA Agassiz, 1847. See Thaumetopoea Hübner, [1820].

THAUMETOPOEA Hübner, [1820] 1816, Verz. bekannter Schmett.: 185.

Type-species: Phalaena processionea Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 500, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 591. TYPE-LOCAL-ITY: not stated, [EUROPE].

CNETHOCAMPA Stephens, 1828, Illust. Br. Ent. (Haustellata) 2: 46.

Type-species: Phalaena processionea Linnaeus, 1758, by subsequent designation by Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 64.

A junior objective synonym of Thaumetopoea Hübner, [1820].

Cnethocampa was again proposed by Stephens, 1829 [June], Nom. Br. Insects: 39, and 1829 [July], Syst. Cat. Br. Insects

THAUMATOPOEA Agassiz, 1847, Nomencl. zool. (Index univl.): 367.

An unjustified emendation of *Thaumetopoea* Hübner, [1820]. The 'Index universalis' formed fascicle 12 of the Nomencl. zool.; Thaumatopoea is dated from the wrapper of the fascicle, not from its title-page which is dated 1846.

‡CNETOCAMPA; Rambur, 1858, Cat. syst. Lépid. Andalousie (1): pl.4, figs 5, 6.

An incorrect subsequent spelling of *Cnethocampa* Stephens, 1828.

THEAGES Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 3: 721.

Type-species: Theages leucophaea Walker, 1855, ibidem 3: 722, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 202. [BRAZIL]: Rio [de] Janeiro.

THEBRONE Boisduval, 1870, Considérations Lépid. Guatemala: 85.

Type-species: Phalaena aglaura Cramer, 1780, Uitlandsche Kapellen (Papillons exot.) 3: 126, [173], pl.263, fig.F, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 182. Surinam.

P. aglaura is dated from the wrapper of the part and not from the title-page of the volume, 1782.

P. aglaura is a junior subjective synonym of Phalaena tricolora Sulzer, 1776, Geschichte Insecten: 160, pl.22, fig.5.

See also Aphisaon Boisduval, 1870.

THELDE Walker, 1862, J. Proc. Linn. Soc. (Zool.) 6: 139.

Type-species: Thelde patula Walker, 1862, ibidem (Zool.) 6: 140, by monotypy. [Borneo]: Sarawak.

T. patula is a subspecies of Repena invaria Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1724.

Thelde was established in the Notodontidae; it was placed in the Lymantriidae by Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 217.

THEMACA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 394. LYMA

Type-species: *Themaca comparata* Walker, 1865, *ibidem* 32: 395, by monotypy. [INDIA]: North Hindostan.

THEMERASTIS Turner, 1903, Proc. Linn. Soc. N.S.W. 28: 46 (key), 63.

Type-species: *Themerastis celaena* Turner, 1903, *ibidem* **28**: 64, by monotypy. AUSTRALIA: Victoria, Melbourne.

T. celaena is a junior subjective synonym of Discophlebia lucasii Rosenstock, 1885, Ann. Mag. nat. Hist. (5) 16: 421, pl.11, fig.4.

Themerastis was established in the Notodontidae; it was placed in the Thaumetopoeidae by Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 2.

THERIOPHILA Hering, 1932, Revue Zool. Bot. afr. 22: 103.

Type-species: *Theriophila miara* Hering, 1932, *ibidem* **22**: 103, pl.2, fig.27, by original designation. [CAMEROUN] Kamerun: Bipindi.

THERMEOLA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: viii, 82, 96.

Type-species: *Thermeola tasmanica* Hampson, 1900, *ibidem* 2: 96, fig. 30, by original designation. [Australia]: Tasmania, Hobart.

THERMIDARCTIA Talbot, 1929, Bull. Hill Mus. Witley 3: 133.

Type-species: *Thermidarctia thermidoides* Talbot, 1929, *ibidem 3*: 133, pl.3, fig.7, by original designation. VENEZUELA: Mérida.

THERMOGRAPTA Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xxii, 568.

Type-species: *Thermograpta rufizonata* Hampson, 1914, *ibidem* (Suppl.) 1: 568, fig.159, by original designation. NEW GUINEA: [Irian Jaya], Snow Mts, upper Setekwa R.

THEROA Schaus, 1901, Trans. ent. Soc. Lond. 1901: 258 (key), 270.

Type-species: *Dasylophia zethus* Druce, 1898, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 2: 454, pl.90, fig.11, by original designation. MEXICO: Jalapa.

THIACIDAS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1027.

NOCTUIDAE

Type-species: *Thiacidas postica* Walker, 1855, *ibidem* 5: 1028, by monotypy. NEPAL.

Thiacidas was established in the Notodontidae; it was placed in the Lymantriidae by Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 445; and was transferred to the Noctuidae by Dyar, 1897, Can. Ent. 29: 12.

THIRMIDA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 466.

Type-species: *Thirmida dimidiata* Walker, 1854, *ibidem* **2**: 466, by monotypy. Colombia.

THOLOCOLEUS Robinson, 1975, Macrolepid. Fiji and Rotuma: 238. NOCTUIDAE

Type-species: Lophocoleus astrifer Butler, 1886, Trans. ent. Soc. Lond. 1886: 417, by original designation. Holotype \mathfrak{P} , Fiji: Viti Islands.

Tholocoleus in the Noctuidae, Hypeninae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

THOLODA Gistl, 1848. See Clostera Samouelle, 1819.

THOSEA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1068.

Type-species: *Thosea unifascia* Walker, 1855, *ibidem* 5: 1068, by monotypy. [Indonesia] East Indies.

Thosea was established in the Notodontidae; it was placed in the Limacodidae by Moore, 1877, Ann. Mag. nat. Hist. (4) 20: 348.

THRENOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: viii, 82, 96.

Type-species: *Tigrioides heminephes* Meyrick, 1886, *Proc. Linn. Soc. N.S.W.* (2) 1: 697, by original designation. AUSTRALIA: New South Wales, Bathurst.

THRINACIA Butler, 1876, J. Linn. Soc. (Zool.) 12: 384.

Type-species: *Glaucopis afflicta* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 1: 144, by original designation. [Brazil]: Para.

G. afflicta is a junior subjective synonym of Sphinx temenus Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 148, pl.367, fig.D.

‡THUMATA; misspelling. See Thumatha Walker, 1866.

THUMATHA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1900.

Type-species: *Thumatha fuscescens* Walker, 1866, *ibidem* 35: 1901, by monotypy. AUSTRALIA: Moreton Bay.

See also Pelobrochis Lucas, 1892.

‡THUMATA; Draudt, 1914, in Seitz, Gross-Schmett. Erde 10: 159.

An incorrect subsequent spelling of *Thumatha* Walker, 1866, in both the German and the English editions.

THUMATHOIDES Holloway, 1979, Series Ent. 15: 383.

Type-species: *Thumathoides fumosa* Holloway, 1979, *ibidem* 15: 384, text-fig.81, pl.71, fig.7, by original designation. New CALEDONIA: Grand Lac.

THYANA Dognin, 1923, Hétérocères nouv. Am. Sud 21: 13.

Type-species: *Thyana sumptuosa* Dognin, 1923, *ibidem* 21: 13, by original designation. BOLIVIA: Rio Songo, [BRAZIL]: Amazon, Sao Paulo de Olivença.

‡THYGORINA; misspelling. See Thyrgorina Walker, [1865].

THYLACIGYNA Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 283.

Type-species: *Orgyia ericae* Germar, [1824], *in* Ahrens, *Fauna Insect. Eur.* 8: pls 17, 18, by subsequent designation by Kirby, 1870, *Zool. Rec.* 6: 389. [GERMANY (B.R.D.)]: Lüneburg.

Thylacigyna is dated from the wrapper of livraison 2, and not from the title-page of the work, 1858.

THYLACOPTERA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: [ix], 22, 52.

Type-species: *Thylacoptera albipuncta* Hampson, 1898, *ibidem* 1: 53, fig.24, by original designation. [Philippines]: Palawan.

‡TALYCOPTERA; Seitz, 1912, in Seitz, Macrolepid. World 10: 66.

An incorrect subsequent spelling of *Thylacoptera* Hampson, 1898, in the English edition.

‡TYLACOPTERA; Seitz, 1912, in Seitz, Gross-Schmett. Erde

An incorrect subsequent spelling of *Thylacoptera* Hampson, 1898, in the German edition.

THYONAEA Dyar, 1910, Zoologica, N.Y. 1: 127. LITH Type-species: Thyone perbella Schaus, 1905, Proc. U.S. natn.

Mus. 29: 198, by original designation. FRENCH GUIANA: St Jean.

THYONE Walker, 1854. See Euthyone Watson nom. n.

THYRARCTIA Hampson, 1901. See Parathyris Hübner, [1819].

THYRETARCTIA Strand, 1912, Arch. Naturgesch. 78 (A) 6: THYRET

Type-species: Metarctia haematica Holland, 1893, Psyche, Camb. 6: 396, by monotypy. [GABON]: [Ogooué] Ogové R.

THYRETES Boisduval, 1847, in Delegorgue, Voyage Afr. australe 2: 596.

Type-species: Sphinx hippotes Cramer, 1780, Uitlandsche Kapellen (Papillons exot.) 3: 166, 175, pl.286, fig.A, but included by Boisduval as 'Thyretes Hiphothes Cramer' an incorrect subsequent spelling, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 103. [INDIA]: [Bengal coast] Côte de Bengale.

S. hippotes is dated from the wrapper of the part and not

from the title-page of the volume, 1782.

THYRGIS Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 316.

Type-species: Phalaena militta Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 159, 250, pl.370, fig.1, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 400. SURINAM.

P. militta is dated from the wrapper of the part, and not

from the title-page of the volume, 1782.

THYRGORINA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 317.

Type-species: Thyrgorina spilosomata Walker, [1865] 1864, ibidem 31: 318, by monotypy. [INDIA]: [S.], South Hindostan. See also Echlida Walker, 1865.

‡THYGORINA; Dyar, 1897, Can. Ent. 29: 212, 215.

An incorrect subsequent spelling of Thyrgorina Walker,

THYROGONIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xi, 23, 139.

Type-species: Syntomis efulensis Holland, 1898, Ent. News 1898: 12, by original designation. [CAMEROUN] Cameroons: Bulé Country, Efulen.

‡TYROGONIA; Pagenstecher, 1909, Geogr. Verbreitung Schmett: 420.

An incorrect subsequent spelling of Thyrogonia Hampson, 1898.

THYROMOLIS Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 2, 68.

Type-species: Idalus pythia Druce, 1900, Ann. Mag. nat. Hist. (7) 6: 67, by original designation. Bolivia: Songo [R.].

THYROSTICTA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xi, 23, 129.

Type-species: Naclia quadrimacula Mabille, 1879, Bull. Soc. zool. Fr. 3: 85, by original designation. MADAGASCAR.

THYSANOPLUSIA Ichinosé, 1973, Kontyu 41: 137.

Type-species: Phytometra intermixta Warren, 1913, in Seitz, Gross-Schmett. Erde 3: 357, pl.64 row g, by original designation. Lectotype by PRESENT DESIGNATION & CHINA: W., Pu-tsu-fu, 8-10000ft, (BMNH).

Thysanoplusia in the Noctuidae, Plusiinae, is cited here as it was not included in Nye, 1975, Generic Names Moths World

THYSANOPRYMNA Butler, 1875, Ann. Mag. nat. Hist. (4)

Type-species: Eucereon pyrrhopyga Walker, 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 105 (but cited as ‡Eucerea, an incorrect subsequent spelling by Walker), by original designation. BRAZIL.

THYSANOPTYX Hampson, 1894, Fauna Br. India (Moths) 2: vii, 40, 74.

Type-species: Lithosia tetragona Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 510, by original designation. [BANGLADESH]: [Sylhet] Silhet.

TICILIA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 394. DREPANIDAE

Type-species: Ticilia argentilinea Walker, 1865, ibidem 32: 394, by monotypy. SINGAPORE.

T. argentilinea is a junior subjective synonym of Damna

gelidata Walker, [1863] 1862, ibidem 26: 1570.

Ticilia was established in the Liparidae, now Lymantriidae; its type-species was placed in the 'Drepanulidae' by Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 339.

TIFAMA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 981 (key), 1077.

Type-species: Phalaena chera Drury, 1773, Illust. nat. Hist. exot. Insects 2: 36, index, pl.20, fig.4 (but included by Walker as Phalaena simois Stoll, 1780, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 39, 252, pl.308, fig.F, a misidentification; see below), by monotypy. SURINAM.

Walker's diagnosis of Tifama was based on two female specimens of Phalaena chera Drury, 1773, which Walker misidentified as Phalaena simois Stoll, 1780. The two specimens which Walker studied are in the BMNH; they have been compared with Drury's figure of P. chera and match it well. The use by Walker of the name simois was possibly a lapsus, for Phalaena chera Drury, 1773, was figured by Stoll, 1780, in Cramer, ibidem 4: 39, pl.308, fig.E, next to P. simois Stoll, 1780, on the same plate.

Under the Code, Article 70(a), the case of a misidentified

type-species should be referred to the Commission.

P. simois is dated from the wrapper of part 26, not from the title-page of volume 4 which is dated 1782.

CHAETOGNATHA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.98, fig.6; 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 3.

Type-species: Phalaena altrix Stoll, 1780, by monotypy. A junior objective synonym of Tifama Walker, 1855.

TIGRIDANIA Kaye, 1918, Ann. Mag. nat. Hist. (9) 2: 225.

Type-species: Tigridania quadricincta Kaye, 1918, ibidem (9) 2: 226, by original designation. [Peru]: upper Amazon. Rio Ucayali.

TIGRIOIDES Butler, 1877, Trans. ent. Soc. Lond. 1877: 359.

Type-species: Setina alterna Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 520, by original designation. [Australia] New Holland: Sydney.

TIMALUS Watson nom. n. for Pterygopterus Butler, 1876.

Type-species: Pterygopterus clavipennis Butler, 1876, J. Linn.

Soc. (Zool.) 12: 411, by original designation (for Pterygopterus Butler). [Brazil.]: [Espirito Santo] Espiritu Santo.

PTERYGOPTERUS Butler, 1876, J. Linn. Soc. (Zool.) 12: 411.

Type-species: *Pterygopterus clavipennis* Butler, 1876, *ibidem* 12: 411, by original designation.

A junior homonym of *Pterygopterus* Kner, 1867, *Sber. Akad. Wiss. Wien* 55 (1): 722, – Pisces. The objective replacement name is *Timalus* Watson nom. n.

TIMORACA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 212.

Type-species: *Chadisra meeki* Rothschild, 1917, *Novit. zool.* **24**: 251, pl.5, fig.3, by original designation. New Guinea: [Irian Jaya] Mt Goliath.

TINEOPSIS Felder, 1861, Sber. Akad. Wiss. Wien 43 (Abt.1): 37.

Type-species: *Tineopsis saturata* Felder, 1861, *ibidem* 43 (Abt.1): 37, by monotypy. [Indonesia]: [Moluccas], [Ambon I.] Amboinensium.

TINOLIODES Wileman, 1915, Entomologist 48: 110 (but attributed to Hampson, an incorrect authorship).

Type-species: *Tinoliodes benguetensis* Wileman, 1915, *ibidem* 48: 110, by original designation. Philippines: Luzon, Benguet subprovince, Palali.

TIPULODES Boisduval, 1832, in d'Urville, Voyage de Découvertes de l'Astrolabe (Faune ent.) 1: 195.

Type-species: *Tipulodes ima* Boisduval, 1832, *ibidem* 1: 196, pl.3, fig.7, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 113. BRAZIL.

TITHRAUSTES Druce, 1885, Biologia cent.-am. (Zool.) Lepid.-Heterocera 1: 158.

Type-species: *Tithraustes haemon* Druce, 1885, *ibidem* 1: 158, pl.14, fig.20, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 413. PANAMA: Volcan de Chiriqui.

TMETOPTERA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.106, fig.29. LITH

Type-species: *Tmetoptera phryganeoides* Felder, 1874, *ibidem* 2 (Abt.2): pl.106, fig.29, by monotypy. BRAZIL: cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 9.

TMETOPTERYX Kiriakoff, 1964, in Wytsman, Genera Insect. 217 (A): 175.

Type-species: Scalmicauda bisecta Rothschild, 1917, Novit. zool. 24: 260, pl.5, fig.12, by original designation. SIERRA LEONE.

TODDIA Kiriakoff, 1967. See Toddiana Kiriakoff, 1973.

TODDIANA Kiriakoff, 1973, Bull. Annls Soc. r. ent. Belg. 109: 42.

Type-species: Fentonia eingana Schaus, 1928, Proc. U.S. natn. Mus. 73 (19): 80, by original designation (for Toddia Kiriakoff, 1967). CHINA: Szechwan, Mt Omei.

Toddiana was established as the objective replacement name for Toddia Kiriakoff, 1967.

TODDIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 111.

Type-species: Fentonia eingana Schaus, 1928, by original designation.

A junior homonym of *Toddia* França, 1911, *Archos Inst. bact. Camara Pestana* 3 (2): 232, – Protozoa. The objective replacement name is *Toddiana* Kiriakoff, 1973.

TOGARISHACHIA Matsumura, 1925, Zool. Mag. Tokyo 37: 399.

Type-species: *Togarishachia argentopicta* Matsumura, 1925, *ibidem* 37: 399, pl.6, fig.3, by original designation. TAIWAN.

TOGARITENSHA Matsumura, 1929, Insecta matsum. 4: 43. NOTO

Type-species: *Togaritensha acuta* Matsumura, 1929, *ibidem* 4: 43, pl.1, fig.3, by original designation. TAIWAN: Horisha.

T. acuta is a junior subjective synonym of Norraca curvilinea Wileman, 1911, Entomologist 44: 174.

TOGEPTERYX Matsumura, 1920, Zool. Mag. Tokyo 32: 149.

Type-species: *Drymonia velutina* Oberthür, 1880, *Etudes d'Entomologie* 5: 64, pl.8, fig.2, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

Togepteryx was again proposed by Matsumura, 1921, Thousand Insects Japan (Additam.) 4: 811.

TOLIMICOLA Prout, 1918, Novit. zool. 25: 396 (key), 403.

Type-species: *Tolimicola fassli* Prout, 1918, *ibidem* **25**: 404, by original designation. Colombia: Monte Tolima.

TOLMIANA Kiriakoff, 1974, Veröff. zool. StSamml. Münch. 17: 375.

Type-species: *Tolmiana cyanosticta* Kiriakoff, 1974, *ibidem* 17: 376, text-fig.2, pl.1, fig.2, by original designation. SUMATRA: S.W. Siantar.

TOPOMESA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1921.

Type-species: *Topomesa subinanis* Walker, 1866, *ibidem* 35: 1921, by monotypy. JAVA.

TOPOMESOIDES Strand, 1910, in Seitz, Gross-Schmett. Erde 2: 133.

Type-species: Aroa jonasii Butler, 1877, Ann. Mag. nat. Hist. (4) 20: 402, by original designation. JAPAN: Yokohama.

The date of publication of the part of Seitz containing page 133, was cited on page 129 as 22.xii.1910 in both the German and the English editions.

TORIGEA Matsumura, 1934, Insecta matsum. 8: 180. NOTO Type-species: Bireta plumosa Leech, 1889, Proc. zool. Soc. Lond. 1888: 620, pl.31, fig.4, by original designation (but cited as plumosa Butler, an incorrect authorship). [JAPAN]: [Ohoyama] Oyama.

TORONA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 467.

Type-species: *Torona ferrifera* Walker, 1865, *ibidem* 32: 468, by monotypy. [INDIA]: Hindostan.

TORTRICOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xiv, 84, 291.

Type-species: *Tortricosia excisa* Hampson, 1900, *ibidem* 2: 291, fig. 208, by original designation. Burma: Mergui, [Sumatra]: Perak, Padang Rengas.

TORYCUS Herrich-Schäffer, 1866, CorrespBl. zool.-min. Ver. Regensburg 20: 119.

Type-species: *Torycus tricolor* Herrich-Schäffer, 1866, *ibidem* **20**: 119, by monotypy. CUBA.

TOSPITIS Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 426.

Type-species: Tospitis nulliferana Walker, 1863, ibidem 28: 427, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 365. BORNEO: Sarawak.

TOULGOETINACLIA Griveaud, 1964, Faune Madagascar

Type-species: Thyrosticta obliquipuncta Rothschild, 1924, Ann. Mag. nat. Hist. (9) 14: 306, by original designation. MADAGASCAR: Diego Suarez.

TRABANTA Kiriakoff, 1962, Annls Mus. r. Afr. cent. (8vo Sci. zool.) 114: 21.

Type-species: Turnaca nigripuncta Rothschild, 1917, Novit. zool. 24: 258, pl.3, fig.25, by original designation. [GHANA] Gold Coast: Takwa.

TRAUMATOCAMPA Wallengren, 1871, Skand. Heterocer-Fjärilar 2 (2): 158.

Type-species: Gastropacha pinivora Treitschke, 1834, in Ochsenheimer, Schmett. Eur. 10 (1): 194, by subsequent designation by Agenjo, 1941, Eos. Madr. 17: 75. GERMANY: northern.

Traumatocampa was established as a subgenus of Cnethocampa Stephens, 1828.

TRIANEURA Butler, 1876, J. Linn. Soc. (Zool.) 12: 353.

Type-species: Glaucopis subaurata Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 149, by original designation. [SRI LANKA] Ceylon.

TRICHAEA Herrich-Schäffer, 1866, CorrespBl. zool.-min. Ver. Regensburg 20: 108 (key), 115. **PYRALIDAE**

Type-species: Trichaea pilicornis Herrich-Schäffer, 1866, ibidem 20: 115, by subsequent designation by Alayo & Arteaga, 1976, Serie biol. Acad. Cienc. Cuba 65: 5. CUBA.

Trichaea was established in the 'Syntomina', now Ctenuchidae; it was transferred to the Pyralidae by Gundlach, 1881, Contr. Ent. Cubana 1: 229.

TRICHAETA Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 52. **CTEN**

Type-species: Syntomis teneiformis Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1595, by monotypy. [INDIA] Hindostan.

‡TRICHALIS; misspelling. See Trischalis Hampson, 1894.

TRICHAREVA Dognin, 1914 February, Hétérocères nouv. Am.

Type-species: Trichareva fulvilaniata Dognin, 1914 Febru-

ary, ibidem 7: 11, by monotypy. Panama: Lino.

The name Trichareva probably originated from Hampson but was used and made nomenclaturally available by Dognin prior to its proposal by Hampson, 1914 [November] as the name for a new genus.

TRICHAREVA Hampson, 1914 [November], Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: xx, 518.

Type-species: Trichareva fulvilaniata Dognin, 1914 February, Hétérocères nouv. Am. Sud 7: 11, by original designation.

A junior homonym of Trichareva Dognin, 1914 February. The objective replacement name is Trichareva Dognin, 1914 February.

TRICHAREVA Hampson, 1914 [November]. See Trichareva Dognin, 1914 February.

TRICHELA Herrich-Schäffer, [1853] 1850-1858, Sanıml. neuer oder wenig bekannter aussereur. Schmett. 1 (1): wrapper, pl.13, fig.53.

Type-species: Trichela tolumnensis Herrich-Schäffer, [1853] 1850-1858, ibidem 1 (1): wrapper, pl.13, fig.53 (but attributed to Boisduval, an incorrect authorship), by monotypy. [Col-OMBIA]: Bogota.

See also Enope Walker, 1854.

‡TRICHETRA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling probably of Trichela Herrich-Schäffer, [1853].

TRICHETRA Westwood, 1840. See Trichiocercus Stephens, 1835.

‡TRICHETRA White, 1841. See Trichiocercus Stephens, 1835.

‡TRICHETRA; misspelling. See Trichela Herrich-Schäffer, [1853].

TRICHIA Nietner, 1861, Observations on the Enemies of the Coffee Tree in Ceylon: 20.

Type-species: Trichia exigua Nietner, 1861, ibidem: 20, by monotypy. [SRI LANKA] Ceylon.

TRICHIOCERCUS Stephens, 1835, Illust. Br. Ent. (Haustellata) 4: 386.

Type-species: Arcturus sparshalli Curtis, 1830, Br. Ent. 7: 336, by original designation (for Arcturus Curtis, 1830). Type-LOCALITY: stated to have been found near Horning [England, Norfolk], but the type was evidently an imported specimen, for

the species occurs in AUSTRALIA.

Trichiocercus was established as the objective replacement name for Arcturus Curtis, 1830. Both these names and Trichetra were established in the Arctiidae; they were placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 440; Trichetra was transferred to the Eupterotidae by Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 497; A. sparshalli, the type-species, was placed in the Cnethocampinae, now Thaumetopoeidae, by Turner, 1922, Proc. Linn. Soc. N.S.W. 47: 363.

ARCTURUS Curtis, 1830, Br. Ent. 7: 336.

Type-species: Arcturus sparshalli Curtis, 1830, by original designation.

A junior homonym of Arcturus Berthold, 1827, in Latreille, Nat. Fam. Thierreichs: 283, - Crustacea. The objective replacement name is Trichiocercus Stephens, 1835.

TRICHETRA Westwood, 1840, Introd. mod. Classif. Insects 2 (Synopsis Genera Br. Insects): 92.

Type-species: Arcturus sparshalli Curtis, 1830, by original designation (for Arcturus Curtis, 1830).

Proposed unnecessarily as an objective replacement name for Trichiocercus Stephens, 1835, which Westwood misspelt as ‡Trichocercus, and mistakenly believed to be preoccupied by Trichocerca Lamarck, 1801.

‡TRICHOCERCUS; Westwood, 1840, Introd. mod. Classif. Insects 2 (Synopsis Genera Br. Insects): 92.

An incorrect subsequent spelling of Trichiocercus Stephens, 1835.

‡TRICHETRA; White, 1841, in Grey, Jls two Exped. Disc. N.W. and W. Australia . . . 1837, 38, and 39 2: 479, 482.

White was not proposing a new genus, as suggested by Neave, 1940, Nomencl. zool. 4: 537, but was using Trichetra Westwood, 1840, for which he cited the original reference on page 482.

ARCTOPROCTUS Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 92.

Type-species: Arcturus sparshalli Curtis, 1830, by PRES-ENT DESIGNATION.

A junior objective synonym of Trichiocercus Stephens, 1835. Arctoproctus was included by Herrich-Schäffer in his 'Synopsis familiarum Lepidopterorum' which was published first in Syst. Bearbeitung Schmett. Eur. 6: 82-132. The title-page of volume 6 is dated 1843-1856 but Herrich-Schäffer, 1869, CorrespBl. zool.-min. Ver. Regensburg 23: 214, has stated that Heft 68 (which included the Synopsis) was published in 1855;

the Heft wrapper is so dated.

The same Synopsis was included in Samml. neuer. oder wenig bekannter aussereur. Schmett. 1: 2-52. The title-page of this work is dated 1850-1858 but Gerstaecker, 1857, Arch. Naturgesch. 23 (2): 431, has stated that pages 1-52 were published in 1856. Further information is given by Fletcher, 1979, in Nye, Generic Names Moths World 3 under Dates of some important Publications.

‡TRICHIOSOMA; misspelling. See Somatrichia Kirby, 1892.

‡TRICHOCERCUS; misspelling. See Trichiocercus Stephens, 1835.

TRICHOCEROSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 91, 563.

Type-species: *Trichocerosia atrifulva* Hampson, 1900, *ibidem* 2: 563, pl.35, fig.11, by original designation. New Guinea: [Papua New Guinea], Milne Bay.

TRICHODESMA Hampson, 1898. See Epidesma Hübner, [1819].

TRICHOFELTIA McDunnough, [1929 April] 1928, Bull. Dep. Mines, Can. 55: 61.

Type-species: Agrotis circumdata Grote, 1883, Ann. Mag. nat. Hist. (5) 11: 53, by original designation.

Further information in Nye, 1975, Generic Names Moths World 1: 489.

TRICHOFELTIA Barnes & Benjamin, 1929 September, Bull. Brooklyn ent. Soc. 24: 177.

Type-species: Agrotis circumdata Grote, 1883, by original designation

Trichofeltia Barnes & Benjamin is a junior objective synonym and also a junior homonym of Trichofeltia McDunnough, [1929]

April] 1928.

Trichofeltia Barnes & Benjamin, in the Noctuidae, Hadeninae is cited here as it was not included in Nye, 1975, Generic

Names Moths World 1.

TRICHOFELTIA Barnes & Benjamin, 1929 September. See Trichofeltia McDunnough, [1929 April]. NOCTUIDAE

TRICHOGYIA Hampson, 1894, Fauna Br. India (Moths) 2: 103.

Type-species: *Trichogyia semifascia* Hampson, 1894, *ibidem* 2: 103, fig.68, by original designation. [Burma]: Tenasserim.

Trichogyia was established in the Arctiidae; it was transferred to the Limacodidae by Snellen, 1895, Dt. ent. Z. Iris 8: 136.

TRICHOLEPIS Hampson, 1891. See Chiretolpis Watson nom. n.

TRICHOLOBA Hampson, 1910, Ann. Mag. nat. Hist. (8) 5:

Type-species: Antheua carteri Druce, 1888, Proc. zool. Soc. Lond. 1887: 685, by original designation. Gambia.

In the original combination of the type-species, the generic name was published as ‡Antheura, an incorrect subsequent spelling.

OVERLAETIA Kiriakoff, 1954, Bull. Inst. r. Sci. nat. Belg. 30 (14): 3.

Type-species: Overlaetia grisescens Kiriakoff, 1954, ibidem 30 (14): 4, fig.3, by original designation. [ZAIRE] Congo Belge: Lupweji.

A junior homonym of *Overlaetia* Schouteden, 1932, *Annls Mus. r. Congo Belge* (C Zool.) (3 Arthropoda, Section 2) 1: 191, – Insecta, Hemiptera. There is no objective replacement

name but *O. grisescens* is congeneric with *Antheua carteri* Druce, 1888, the type-species of *Tricholoba* Hampson, 1910; the latter is thus available for use as a subjective replacement name.

TRICHOMAPLATA Wing, [1850], *Proc. zool. Soc. Lond.* **1849**: 105.

Type-species: *Trichomaplata vittata* Wing, [1850], *ibidem* **1849**: 105, Annulosa pl.14, fig.1, by original designation. BRAZII.

In the original publication the generic name was spelt twice on page 105 and once on page 179 as *Trichomaplata*; on the legend to Annulosa plate 14 it was spelt as ‡*Trichomoplata*, an incorrect (of a multiple) original spelling.

‡TRICHOMOPLATA; Wing, [1850], Proc. zool. Soc. Lond. 1849: Annulosa pl.14, legend.

An incorrect (of a multiple) original spelling of *Trichoma-plata* Wing, [1850].

ICHTHYOSOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.17.

Type-species: *Ichthyosoma tigniferum* Felder, 1874, *ibidem* 2 (Abt.2): pl.97, fig.17, by monotypy. [Brazil]: River Amazon, cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 5 (as ‡*Jchthyosoma*, an incorrect subsequent spelling).

A junior homonym of *Ichthyosoma* Montrouzier, 1855, *Annls Soc. Agric. Lyon* (2) 7: 58, – Insecta, Coleoptera. There is no objective replacement name, but *I. tigniferum* is a junior subjective synonym of *Trichomaplata vittata* Wing, [1850], the type-species of *Trichomaplata* Wing, [1850]; the latter is thus available for use as a subjective replacement name.

Felder, 1874, in his script legends to his plates, did not distinguish between the capital letters I and J; the spelling *Ichthyosoma* has generally been adopted by authors.

‡JCHTHYOSOMA; Felder, 1875, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2) Erklärung to pls 75-107: 5.

An incorrect subsequent spelling of *Ichthyosoma* Felder, 1874.

TRICHOMELIA Schaus, 1892 August, Proc. zool. Soc. Lond. 1892: 283.

Type-species: *Trichomelia celenna* Schaus, 1892 August, *ibidem* 1892: 283, by monotypy. BRAZIL: Rio de Janeiro.

T. celenna was redescribed by Schaus, 1892 November, Am. Lepid.: 13, pl.2, fig.5

‡TRICHOMOPLATA; misspelling. See Trichomaplata Wing, [1850].

‡TRICHONOCHA; misspelling. See Tritonaclia Hampson, 1898.

TRICHOSOMA Rambur, 1832. See Somatrichia Kirby, 1892.

‡TRICHOSOMUM; misspelling. See Somatrichia Kirby, 1892.

TRICHOTA Rambur, 1866. See Nudaria Haworth, [1809].

TRICHOTIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.97, fig.19.

Typc-species: *Trichotis picta* Felder, 1874, *ibidem* 2 (Abt.2): pl.97, fig.19, by monotypy. Type-locality: cited by Felder, 1875, *ibidem* 2 (Abt.2) Erklärung to pls 75-107: 9, as 'fl. Amazonas', but this was probably an error, as the species is currently recorded from the states of Texas, Georgia and Florida in the U.S.A.

T. picta is a form of Heterocampa astarte Doubleday, 1841, Entomologist 1: 57.

See also *Heterocampa* Doubleday, 1841, and *Balia* Doubleday, 1869.

TRICHROA Fairmaire, 1894, Annls Soc. ent. Belg. 38: 395.

Established as a replacement name for *Trichromia* Fairmaire, 1892, *Annls Soc. ent. Fr.* 61 (Bull.): xxi, – Insecta, Coleoptera.

The name *Trichroa* Fairmaire has appeared in the literature of Lepidoptera (e.g. Strand, 1919, in Wagner, Lepid. Cat. 22: 88 – as a junior synonym of Neritos Walker) probably because of the misleading citation by Fairmaire, 1894, ibidem 38: 395 – 'Au lieu de . . . Trichromia (Hübn., Lépid.) =TRICHROA' whereas Fairmaire was in fact replacing Trichromia Fairmaire, 1892, Annls Soc. ent. Fr. 61: xxi, – Insecta, Coleoptera [a junior homonym of Trichromia Hübner, [1819], – Lepid., Arctiidae] with Trichroa Fairmaire, 1894.

TRICHROMIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 164.

Type-species: *Phalaena onytes* Cramer, 1777, *Uitlandsche Kapellen (Papillons exot.)* 2: 107, 150, pl.165, fig.E, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 290. SURINAM.

P. onytes is dated from the wrapper of the part and not from

the title-page of the volume, 1779.

Trichromia was established in the 'Lithosiae'; it was retained in the 'Lithosiidae' by Kirby, 1892; and was placed in the 'Syntomidae' by Hampson, 1905, Ann. Mag. nat. Hist. (7) 15: 429, who incorrectly cited Sphinx capys Cramer, 1775, as type-species. Trichromia is here transferred to the Arctiidae.

TRICHURA Hübner, [1819] 1816, Verz. bekannter Schmett.: 126.

Type-species: *Trichura druryi* Hübner, [1819] 1816, *ibidem*: 126, by monotypy. [GUYANA]: Demerara and Berbice.

T. druryi was established for the concept misidentified by Cramer, 1775, Uitlandsche Kapellen (Papillons exot.) 1: 7, pl.4, figs F, G as Sphinx coarctata Drury, 1773, Illust. nat. Hist. exot. Insects 2: index, 46, fig. The syntypes of T. druryi Hübner are therefore the specimens represented by Cramer's 1775 figures F and G.

TRICYPHA Möschler, 1878, Verh. zool.-bot. Ges. Wien 27 (Abh.): 654.

Type-species: *Tricypha furcata* Möschler, 1878, *ibidem* 27 (Abh.): 655, by monotypy. Surinam: interior of.

See also Maenoleneura Butler, 1878.

TRINCOMALA Kiriakoff, 1968, in Wytsman, Genera Insect. 217 (C): 213.

Type-species: Chadisra semiferrea Hampson, 1910, J. Bombay nat. Hist. Soc. 20: 92, pl.F, fig.35, by original designation (but cited as 'Chadisra bipars var. semiferrea Hampson', an incorrect original combination). [SRI LANKA] Ceylon: Trincomalee.

‡TRIPHEROMERA; misspelling. See Trypheromera Butler; 1881.

TRIPURA Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 298. LITH

Type-species: *Tripura prasena* Moore, [1860] 1858-9, *in* Horsfield & Moore, *ibidem* 2: 299, pl.7a, fig.6, by monotypy. INDIA: northern.

TRISCHALIS Hampson, 1894, Fauna Br. India (Moths) 2: vii, 42, 101.

Type-species: Hemonia flava Hampson, 1893, Illust. typical

Specimens Lepid. Heterocera Colln Br. Mus. 9: 14, 83, fig., by original designation. [SRI LANKA] Ceylon: Nawala-pittia.

H. flava is a junior subjective synonym of Tospitis absconditana Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 28: 432.

‡TRICHALIS; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 422.

An incorrect subsequent spelling of *Trischalis* Hampson, 1894.

‡TRISCHALLIS; Seitz, 1914, in Seitz, Gross-Schmett. Erde 10: 123.

An incorrect subsequent spelling of *Trischalis* Hampson, 1894, in both the German and the English editions.

‡TRISCHALLIS; misspelling. See Trischalis Hampson, 1894.

TRISSOBROCHA Turner, 1915, Proc. Linn. Soc. N.S.W. 39: 548.

Type-species: *Trissobrocha eugraphica* Turner, 1915, *ibidem* 39: 548, by monotypy. Australia: New South Wales, Ebor scrub.

TRISTANIA Kirby, 1892. See Nyctemera Hübner, [1820].

TRISULA Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 420.

NOCTUIDAE

Type-species: *Trisula variegata* Moore, [1860] 1858-9, *in* Horsfield & Moore, *ibidem* 2: 420, pl.12a, fig.1, by monotypy. INDIA: northern and Madras.

Trisula was established in the Bombyces; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 458; and was placed in the Noctuidae by Hampson, 1894, Fauna Br. India (Moths) 2: 437.

TRISULOIDES Butler, 1881, Ann. Mag. nat. Hist. (5) 7: 36.
NOCTUIDAE

Type-species: *Trisuloides sericea* Butler, 1881, *ibidem* (5) 7: 36, by monotypy. [INDIA]: [Darjeeling] Darjiling, and Assam, Shillong.

Trisuloides was established in the Liparidae, now Lymantriidae; it was placed in the Noctuidae, by Hampson, 1894, Fauna Br. India (Moths) 2: 436.

TRITONACLIA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xi, 23, 135.

Type-species: *Hydrusa kefersteinii* Butler, 1882, *Cistula ent.* 3: 2, by original designation. MADAGASCAR: Ankafana.

‡TRICHONOCHA; Pagenstecher, 1909, Geogr. Verbreitung Schmett.: 420.

An incorrect subsequent spelling of *Tritonaclia* Hampson, 1898.

TRITOPHIA Kiriakoff, 1967, in Wytsman, Genera Insect. 217 (B): 141.

Type-species: *Phalaena phoebe* Siebert, 1790, *in* Scriba, *Beitr*. *Insekten-Geschichte* 1: 18, pl.2, by original designation. Type-LOCALITY: not stated, [GERMANY].

P. phoebe is a junior subjective synonym of Bombyx tritophus [Denis & Schiffermüller], 1775, Ankündung syst. Werkes Schmett. Wienergegend: 63.

TROCHUDA Schaus, 1904, Trans. Am. ent. Soc. 30: 140.

Type-species: *Trochuda bilinea* Schaus, 1904, *ibidem* 30: 140, by monotypy. [GUYANA] British Guiana: Omai.

TROCODIMA Watson nom. n. for Microdota Dognin, 1906.

Type-species: *Microdota lenistriata* Dognin, 1906, *Annls Soc. ent. Belg.* 50: 183, by original designation (for *Microdota* Dognin, 1906). [PARAGUAY, [ARGENTINA]: Tucuman.

MICRODOTA Dognin, 1906, Annls Soc. ent. Belg. 50: 183.

Type-species: Microdota lenistriata Dognin, 1906, ibidem

50: 183, by original designation.

A junior homonym of *Microdota* Mulsant & Rey, 1873, *Hist.* nat. Coléoptères France (Brévipennes): 295, – Insecta, Coleoptera. The objective replacement name is *Trocodima* Watson nom. n.

‡MICRODATA; Strand, 1919, in Wagner, Lepid. Cat. 22: 54. An incorrect subsequent spelling of Microdota Dognin, 1906.

TROPACME Hampson, 1894, Fauna Br. India (Moths) 2: viii,

Type-species: *Tropacme cupreimargo* Hampson, 1894, *ibidem* 2: 101, fig.65, by original designation. [Burma]: Tenasserim.

TROTONOTUS Butler, 1898, Proc. zool. Soc. Lond. 1898: 431.

Type-species: Trotonotus bettoni Butler, 1898, ibidem 1898: 431, pl.33, fig.2, by original designation. [Kenya]: Mgana.

Trotonotus was established in the Eupterotidae; it was placed in the Notodontidae by Bethune-Baker, 1911, Ann. Mag. nat. Hist. (8) 7: 558.

See also Prionocentrum Janse, 1920.

TRUMANDA Dognin, 1911, Hétérocères nouv. Am. Sud 3: 19.

Type-species: *Trumanda fifiana* Dognin, 1911, *ibidem* 3: 19, by original designation. FRENCH GUIANA.

TRYPHEROMERA Butler, 1881, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 5: ix, 45. ARCT

Type-species: *Nyctemera plagifera* Walker, 1854, *List Specimens lepid. Insects Colln Br. Mus.* 2: 400, by original designation. [Bangladesh]: [Sylhet] Silhet, Hong Kong.

N. plagifera is a junior subjective synonym of Phalaena adversata Schaller, 1788, Naturforscher Berlin 23: 52.

‡TRIPHEROMERA; Swinhoe, 1903, Trans. ent. Soc. Lond. 1903: 55, 67.

An incorrect subsequent spelling of Trypheromera Butler, 1881.

TSARAFIDYNIA Griveaud, 1964, Faune Madagascar 17: 56.

Type-species: Aglaope perpusilla Mabille, 1880, Annls Soc. ent. Fr. (5) 9: 348, by original designation. MADAGASCAR.

TSIRANANACLIA Griveaud, 1964, Faune Madagascar 17: 118.

Type-species: *Tsirananaclia tripunctata* Griveaud, 1964, *ibidem* 17: 119, text-figs 284-286, pl.1, fig.82, by original designation. MADAGASCAR: central, Ambohimahasoa, Tsarafidy forest.

TUERTELLA Kiriakoff, 1976, Tierreich 98: 114.

Type-species: *Tuerta rema* Druce, 1910, *Ann. Mag. nat. Hist.* (8) 6: 169, by original designation. Type(s) 3, [TANZANIA] German East Africa, (BMNH).

Tuertella in the Agaristidae is cited here as it was not included in Nye, 1975, Generic Names Moths World 1.

TUINA Butler, 1877, Trans. ent. Soc. Lond. 1877: 326. LITH Type-species: Ituna cingulata Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 481, by monotypy (of Ituna Walker, 1854). HONDURAS.

Tuina Butler was established as the objective replacement name for Ituna Walker, 1854.

ITUNA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 480.

Type-species: *Ituna cingulata* Walker, 1854, *ibidem* 2: 481, by monotypy. HONDURAS.

A junior homonym of *Ituna* Doubleday, 1847, *in* Doubleday & Westwood, *Genera diurnal Lepid*. 1: 113, – Lepid., Nymphalidae. The objective replacement name is *Tuina* Butler, 1877.

TUMICLA Wallengren, 1863, Wien. ent. Monatschr. 7: 146, 147.

Type-species: Setina sagenaria Wallengren, 1860, ibidem 4: 46, by original designation (but cited as ‡faginaria, an incorrect subsequent spelling). [SOUTH AFRICA]: Caffraria, eastern.

Tumicla was redescribed by Wallengren as new in 1865, K. svenska Vetensk Akad. Handl. 5 (4): 45.

‡TUMIDA; Kirby, 1892, Synomynic Cat. Lepid. Heterocera 1: 306.

An incorrect subsequent spelling of *Tumicla* Wallengren, 1863.

‡TUMIDA; misspelling. See Tumicla Wallengren, 1863.

‡TURENNA; misspelling. See Turuenna Walker, 1865.

TURLINA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 226.

Type-species: *Turlina punctata* Griveaud, [1976] 1975, *ibidem* 80: 226, by original designation. MADAGASCAR: E., région de Moramanga, route de Beparasy, Km 17.

TURLINIA Toulgoët, 1976, Bull. Soc. ent. Fr. 81: 191. LITH Type-species: Turlinia phalaenoides Toulgoët, 1976, ibidem 81: 191, fig.14, by original designation. MADAGASCAR: W., East of Sakaraha, réserve spéciale du Zombitsy.

TURNACA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 454.

Type-species: *Turnaca acuta* Walker, 1865, *ibidem* 32: 454, by monotypy. [INDIA]: [Kanara] Canara.

TURNACOIDES Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 427.

Type-species: *Turnacoides basipuncta* Gaede, 1928, *ibidem* **14**: 427, pl.71 row a, by monotypy. [ZIMBABWE-]RHODESIA.

The date of publication of the part of Seitz containing page 427 was cited on page 425 as 30.iii.1928 in both the German and the English editions.

TURRIGA Walker, 1869, Characters undescr. Lepid. Heterocera: 15.

Type-species: *Turriga invasa* Walker, 1869, *ibidem*: 15, by monotypy. [Australia]: [Queensland], Moreton Bay.

T. invasa is a junior subjective synonym of Olene mendosa Hübner, 1823, Zuträge Samml. exot. Schmett. 2: 19, figs 293, 294.

See also Olene Hübner, 1823.

TURUENNA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 545.

Type-species: *Turuenna dirphioides* Walker, 1865, *ibidem* 32: 545, by monotypy. Type-locality: not known, [South America]: [northern].

Turuenna was established in the Saturniidae; it was placed in the Lasiocampidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 791, as ‡Turumena, an incorrect subsequent spelling; and was placed in the Lymantriidae by Dyar, 1912, Proc. ent. Soc. Wash. 14: 57, as ‡Turenna, an incorrect subsequent spelling.

See also Isoctenia Felder, 1874.

‡TURUMENA; Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 791.

An incorrect subsequent spelling of Turuenna Walker, 1865.

‡TURENNA; Dyar, 1912, Proc. ent. Soc. Wash. 14: 57. An incorrect subsequent spelling of Turuenna Walker, 1865.

‡TURUMENA; misspelling. See Turuenna Walker, 1865.

TURUPTIANA Walker, 1869, Characters undescr. Lepid. Heterocera: 19.

Type-species: Turuptiana obliqua Walker, 1869, ibidem: 19, by monotypy. Type-locality: not stated, [AMERICA]: [tropical].

See also Sallaea Felder, 1874.

‡TYLACOPTERA; misspelling. See Thylacoptera Hampson, 1898.

TYLANTHES Meyrick, 1889, Trans. ent. Soc. Lond. 1889:

Type-species: Tylanthes ptochias Meyrick, 1889, ibidem 1889: 460, by montypy. NEW GUINEA: [Papua New Guinea], Port Moresby.

TYMPANOPHORA Laboulbène, 1865. See Cymbalophora Rambur, 1866.

TYRIA Hübner, [1819] 1816, Verz. bekannter Schmett.: 166.

Type-species: Phalaena jacobaeae Linnaeus, 1758, Syst. Nat. (Edn 10) 1: 511, by subsequent designation by Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: 185. TYPE-LOCALITY: not stated, [EUROPE].

‡HIPOCRITA Hübner, [1806], Tentamen determinationis

digestionis . . .: [1].

Included in a work rejected for nomenclatural purposes by the International Commission on Zoological Nomenclature, 1926, Smithson. misc. Collns 73 (4) Opinion 97: 19, also idem, 1954, Opin. Decl. Int. Commn Zool. Nom. 6 Opinion 278: 140. Only included species Phalaena jacobaeae Linnaeus, 1758.

EUCHELIA Boisduval, 1828, Eur. Lepid. Index meth.: 39.

Type-species: Phalaena jacobaeae Linnaeus, 1758, by subsequent designation by Blanchard, 1845, Hist. Insectes 2: 377 (but cited as ‡jacobeae, an incorrect subsequent spelling).

A junior objective synonym of Tyria Hübner, [1819].

‡TYROGONIA; misspelling. See Thyrogonia Hampson, 1898. CTEN

ULINA Kiriakoff, 1967, Biologia gabon. 3: 353. NOTO Type-species: Ulina oculata Kiriakoff, 1967, ibidem 3: 354, fig.39, by original designation. GABON: Makokou.

ULINELLA Kiriakoff, 1954, Revue Zool. Bot. afr. 49: 312.

Type-species: Ulinella cotytto Kiriakoff, 1954, ibidem 49: 312, pl.1, fig.10, pl.2, figs 10, 11, by original designation. [ZAIRE] Congo Belge: Katanga, [Lubumbashi] Elisabethville.

ULINODES Kiriakoff, 1968, Bull. Annls Soc. r. ent. Belg. 104: 139.

Type-species: Ulinodes costalis Kiriakoff, 1968, ibidem 104: 140, fig.5, by original designation. MOZAMBIQUE: Chiluva Hills.

ULIOLEPIS Warren, 1897, Novit. zool. 4: 385. LYMA Type-species: Uliolepis pilosa Warren, 1897, ibidem 4: 386, by original designation. Afghanistan: Sefir-Kuh.

U. pilosa is a junior subjective synonym of Ocneria signatoria Christoph, 1893, Dt. ent. Z. Iris 6: 88.

Uliolepis was established in the Geometridae; it was transferred to the Lymantriidae by Prout, 1938, in Seitz, Gross-Schmett. Erde 4 (Suppl.): 216.

PSEUDOLABIS Hartig, 1963, in Hartig & Prola, Mitt. münch. ent. Ges. 53: 66.

Type-species: Ocneria signatoria Christoph, 1893, Dt. ent. Z. Iris 6: 88, by monotypy. [U.S.S.R.]: Transcaucasi, Eldar;

A junior homonym of Pseudolabis Matthew, 1904, Bull. Am. Mus. nat. Hist. 20: 211, - Mammalia. There is no objective replacement name but O. signatoria is the senior subjective synonym of Uliolepis pilosa Warren, 1897, the type-species of Uliolepis Warren, 1897. The latter is therefore available for use as a subjective replacement name.

UMNCUMBATA Kiriakoff, 1975, Arnoldia, (Rhodesia) 7 (15):

Type-species: Umncumbata nyikae Kiriakoff, 1975, ibidem 7 (15): 5, fig.4, by original designation. Zambia/ Malawi: border, Nyika Plateau.

URAGA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 464.

Type-species: Uraga hoemorrhoa Walker, 1854, ibidem 2: 465, by monotypy. Jamaica.

URANOPHORA Hübner, [1831] 1825, Zuträge Samml. exot. Schmett. 3: 14.

Type-species: Uranophora chalybea Hübner, [1831] 1825, ibidem 3: 14, pl.[76], figs 439, 440, by monotypy. CUBA.

URGEDRA Dyar, 1908, Proc. ent. Soc. Wash. 9: 64. Type-species: Heterocampa striata Druce, 1906, Ann. Mag. nat. Hist. (7) 17: 410, by original designation. Peru: southeastern, Oconeque.

‡URGERDA; Druce, 1911, Ann. Mag. nat. Hist. (8) 8: 145. An incorrect subsequent spelling of *Urgedra* Dyar, 1908.

‡URGERDA; misspelling. See Urgedra Dyar, 1908. NOTO

UROCAMPA Staudinger, 1892, in Romanoff, Mém. Lépid. 6: 343.

Type-species: Harpyia ocypete Bremer, 1861, Bull. Acad. imp. Sci. St. Pétersb. 3: 481, by monotypy. [U.S.S.R.]: Between Ussuri estuary and [River] Noor.

Urocampa was established as a subgenus of Uropus Rambur, 1832.

See also Fentonia Butler, 1881, and Neoshachia Matsumura, 1925.

UROCOMA Herrich-Schäffer, [1858] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 82. LYMA

Type-species: Porthesia limbalis Herrich-Schäffer, [1855] 1850-1858, *ibidem* 1(1): wrapper, pl.68, fig.389; ibidem 1: 82, by monotypy. [Australia] Nov. Holl.

Urocoma was first published in the synonymy of Porthesia limbalis as 'Urocoma boeckeae Bd. in litt.' but Urocoma Herrich-Schäffer has been treated as a valid generic name by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 468, and so is an available generic name under the Code, Article 11(d). ‡U. boeckeae has not been treated as a valid specific name, and is therefore an unavailable name and under the Code, Article 68(c), is not eligible for fixation as type-species.

ELA Walker, 1862, Trans. ent. Soc. Lond. (3) 1: 273.

Type-species: Ela leucophaea Walker, 1862, ibidem (3) 1: 273, by monotypy. AUSTRALIA.

A junior homonym of Ela Giebel, 1856, Fauna Vorwelt 2 (1): 256, - Neuroptera. There is no objective replacement name but *E. leucophaea* is here placed as a junior subjective synonym of *Porthesia limbalis* Herrich-Schäffer, [1855], the type-species of *Urocoma* Herrich-Schäffer, [1858]; the latter is therefore available for use as a subjective replacement name.

Ela Walker, 1862, was established in the Notodontidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 924, and since then has not been cited; it is here placed as a junior subjective synonym of Urocoma Herrich-Schäffer, [1858].

URODONTA Staudinger, 1887, in Romanoff, Mém. Lépid. 3: 217.

Type-species: *Notodonta albimacula* Staudinger, 1887, *ibidem* 3: 217, pl.12, figs 7a, b, by monotypy. [U.S.S.R.]: Vladivostok, Ussuri.

Urodonta was established as a subgenus of Notodonta Ochsenheimer, 1810; it was treated as a genus by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 598.

URODONTOIDES Matsumura, 1929, Insecta matsum. 4: 48.

Type-species: *Uropus branickii* Oberthür, 1880, *Etudes d'Entomologie* 5: 60, pl.6, fig.6, by original designation. [U.S.S.R.]: [S.E. of Vladivostok], Askold Island.

URODONTOPSIS Matsumura, 1929, Insecta matsum. 4: 48.

Type-species: *Urodonta arcuata* Alphéraky, 1897, *in* Romanoff, *Mém. Lépid.* **9**: 154, pl.11, fig.9, by original designation. [U.S.S.R.]: [Sedimi 43°05′N 131°24′E] Sidemi.

UROLASIA Hampson, 1898 [December], Cat. Lepid. Phalaenae Br. Mus. 1: xvii, 27, 370.

Type-species: *Syntrichura brodea* Schaus, 1896, *Jl N.Y. ent. Soc.* 4: 132, by original designation. TRINIDAD.

See Urolosia Druce, 1898 May.

UROLOSIA Druce, 1898 May, Ann. Mag. nat. Hist. (7) 1: 404.

Type-species: *Urolosia opalocincta* Druce, 1898, *ibidem* (7) 1: 404, by monotypy. [FRENCH GUIANA]: Cayenne.

The name *Urolosia* probably originated as a misspelling of *Urolasia* Hampson, 1898 [December], but was used and made nomenclaturally available by Druce prior to the establishment of *Urolasia* by Hampson.

UROPUS Rambur, 1832. See Dicranura R.L., 1817. NOTO

UROPYIA Staudinger, 1892, in Romanoff, Mém. Lépid. 6: 344.

Type-species: *Notodonta meticulodina* Oberthür, 1884, *Etudes d'Entomologie* **10**: 16, pl.1, fig.3, by monotypy. [U.S.S.R.]: [S.E. of Vladivostok], Sidemi.

UROZANA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 89, 378.

Type-species: *Odozana cordatula* Druce, 1885, *Biologia cent.-am.* (Zool.) Lepid.-Heterocera 1: 125, pl.12, fig.20, by original designation. Guatemala: El Tumbador and Las Mercedes.

URSIA Barnes & McDunnough, 1911, Jl N.Y. ent. Soc. 19: 160.

Type-species: *Ursia noctuiformis* Barnes & McDunnough, 1911, *ibidem* 19: 160, fig.1, by original designation. U.S.A.: Arizona, Yavapai County.

USIMBARA Collenette, 1955, Trans. R. ent. Soc. Lond. 107: 189.

Type-species: *Aroa lata* Holland, 1893, *Psyche, Camb.* **6**: 432, 567, pl.10, fig.13, by original designation. [GABON]: [Ogooué] Ogové R. valley.

UTETHEISA Hübner, [1819] 1816, Verz. bekannter Schmett.: 168.

Type-species: *Phalaena ornatrix* Linnaeus, 1758, *Syst. Nat.* (Edn 10) 1: 511, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 345. AMERICA: [northern].

‡UTHETHEISA; Hübner, [1824] 1806, Samml. exot. Schmett. 2: pl.[181].

An incorrect subsequent spelling of *Utetheisa* Hübner, [1819].

‡UTETHESIA; Moore, [1860] 1858-9, in Horsfield & Moore, Cat. lepid. Insects Mus. nat. Hist. East-India House 2: 306.

An incorrect subsequent spelling of *Utetheisa* Hübner, [1819].

‡UTETHESIA; misspelling. See Utetheisa Hübner, [1819].

‡UTHETHEISA; misspelling. See Utetheisa Hübner, [1819].

UTIDAVA Walker, [1863] 1862, List Specimens lepid. Insects Colln Br. Mus. 26: 1689.

Type-species: *Utidava incomptaria* Walker, [1863] 1862, *ibidem* **26**: 1689, by monotypy. [SRI LANKA] Ceylon.

Utidava was established in the Geometridae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 456.

UTIDAVIANA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 436.

Type-species: *Utidava citana* Schaus, 1893, *in* Schaus & Clements, *On a Colln of Sierra Leone Lepid*.: 26, pl.1, fig.10, by monotypy. SIERRA LEONE.

The date of publication of the part of Seitz containing page 436 was cited on page 433 as 30.v.1928 in the German edition; and as 26.xi.1928 in the English edition.

‡UTRICULIFERA; misspelling. See Utriculofera Hampson, 1893.

UTRICULOFERA Hampson, 1893, Illust. typical Specimens Lepid. Heterocera Colln Br. Mus. 9: 41, 81.

Type-species: *Utriculofera fuscapex* Hampson, 1893, *ibidem* 9: 13, 82, pl.158, figs 7, 17, by monotypy. [Sri Lanka] Ceylon: Nawala-pittia.

‡UTRICULIFERA; Draudt, 1914, in Seitz, Gross-Schmett. Erde 10: 176.

An incorrect subsequent spelling of *Utriculofera* Hampson, 1893, in both the German and the English editions.

UXELA Walker, 1866. See Gangamela Walker, [1865].
CTEN

UXIA Walker, 1866, List Specimens lepid. Insects Colln Br. Mus. 35: 1897.

Type-species: *Uxia albida* Walker, 1866, *ibidem* 35: 1897, by monotypy. North America.

U. albida is a junior subjective synonym of Clemensia albata Packard, 1864, Proc. ent. Soc. Philad. 3: 101.

See also Clemensia Packard, 1864, and Repa Walker, 1866.

‡UZELA; misspelling. See Gangamela Walker, [1865].

VADONACLIA Griveaud, 1964, Faune Madagascar 17: 81.

Type-species: *Vadonaclia marginepuncta* Griveaud, 1964, *ibidem* 17: 81, text-figs 189-191, pl.1, fig.52, by original designation. MADAGASCAR: Maroantsetra, Ambodivoangy.

VAMUNA Moore, 1878, Proc. zool. Soc. Lond. 1878: 10.

Type-species: *Lithosia remelana* Moore, 1866, *ibidem* **1865**: 798, by original designation. INDIA: Darjeeling.

VANEECKEIA Kiriakoff, 1967, Tijdschr. Ent. 110: 49. NOTO Type-species: Stauropus ovalis Eecke, 1929, Zool. Meded. Leiden 12: 166, pl.14, fig.7, by original designation. SUMATRA: Fort de Kock.

S. ovalis is currently treated as a subspecies of Stauropus pallidifascia Hampson, [1893] 1892, Fauna Br. India (Moths) 1: 151.

VANESSODES Grote & Robinson, 1870, Trans. Am. ent. Soc. 3: 176.

Type-species: Vanessodes clarus Grote & Robinson, 1870,

ibidem 3: 176, by monotypy. U.S.A.: Texas.

V. clarus is a junior subjective synonym of Pagara simplex Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1679.

See also Pagara Walker, 1856, and Comacla Walker, [1865].

VARATRA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 231.

Type-species: *Dasychira acosmeta* Collenette, 1939, *Ento-mologist* 72: 144, pl.2, fig.2, by original designation. MADA-GASCAR: Central, Ambalarondra.

VARMINA Moore, 1888, Proc. zool. Soc. Lond. 1888: 405.

Type-species: Gluphisia indica Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1039, by monotypy. EAST INDIES.

Varmina was established in the Lasiocampidae; it was placed in the Liparidae, now Lymantriidae, by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 467.

VENEDICTOFFIA Toulgoët, 1977, Nouv. Revue Ent. 7: 335, 337.

Type-species: Venedictoffia flavicollis Toulgoët, 1977, ibidem 7: 335, 337, fig.8, by original designation. Ecuador: Prov. Napo, Laguna de Papallacta.

VIANANIA Orfila, 1954, Physis, B. Aires 20: 479.

Type-species: Cisthene argentinensis Rothschild, 1912, Novit. zool. 19: 230, by original designation. ARGENTINA: Salta and Tucuman.

VIETTEELLA Kiriakoff, 1958, Mém. Inst. scient. Madagascar (E) 9: 223.

Type-species: Desmeocraera madagascariensis Draeseke, 1937, Dt. ent. Z. Iris 51: 190, by original designation. MADAGASCAR: Ambalarondra.

VIETTEMA Griveaud, 1977, Faune Madagascar 43: 216.

Type-species: Euproctis ratovosoni Viette, 1967, Bull. mens. Soc. linn. Lyon 36: 56, fig.1, by original designation. MADAGASCAR: W., au Nord de Morondava, foret d'Andranomena.

VIETTERIA Griveaud, [1976] 1975, *Bull. Soc. ent. Fr.* **80**: 229.

Type-species: Dasychira ampliata Butler, 1878, Ann. Mag. nat. Hist. (5) 2: 460, by original designation. MADAGASCAR: Ellongo.

VIETTESIA Toulgoët, 1959, Annls Soc. ent. Fr. 128: 122.

Type-species: Coracia plumicornis Butler, 1882, Cistula ent. 3: 4, by original designation. MADAGASCAR: Ankafana.

VIRBIA Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 2: 470.

Type-species: *Virbia mentiens* Walker, 1854, *ibidem* 2: 471, by subsequent designation by Kirby, 1892, *Synonymic Cat. Lepid. Heterocera* 1: 400. VENEZUELA.

VIRIDIFENTONIA Nakamura, 1974, Trans. lepid. Soc. Japan 25: 118.

Type-species: *Heterocampa plagiviridis* Moore, 1879, in Hewitson & Moore, *Descr. new Indian lepid. Insects Colln late Mr W.S. Atkinson* (1): 61, by original designation. INDIA: Darjeeling.

VITRONACLIA Griveaud, 1964, Faune Madagascar 17: 17.

Type-species: Naclia veronica Oberthür, 1893, Etudes d'Entomologie 18: 8, pl.1, fig.15, by original designation. MADAGASCAR: Alahakato.

VIVIENNEA Watson, 1975, Bull. Br. Mus. nat. Hist. (Ent.)
Suppl. 25: 11.

Type-species: Automolis moma Schaus, 1905, Proc. U.S. natn. Mus. 29: 218, by original designation. GUYANA: Omai.

VOHITRA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 231.

Type-species: *Dasychira melissograpta* Collenette, 1936, *Novit. zool.* 40: 163, by original designation. MADAGASCAR: E., Périnet (149km E. of Tananarivo).

VOLANA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 230.

Type-species: *Dasychira cyclota* Collenette, 1959, *Annls Soc. ent. Fr.* **128**: 110, pl.2, fig.15, by original designation. MADAGASCAR: Centre, Ankaratra, Ambahona.

VOLGARCTIA Alphéraky, 1908. See Acymba Rambur, 1866.

VULMARA Schaus, 1924, Proc. U.S. natn. Mus. 65 (7): 24.

Type-species: Vulmara drostana Schaus, 1924, ibidem 65 (7): 25, by original designation. Guatemala: Chejel.

VULSINIA Schaus, 1928, Proc. ent. Soc. Wash. 30: 54. CTEN Type-species: Vulsinia socorra Schaus, 1928, ibidem 30: 54, by original designation. PHILIPPINES: Socorro.

VUNGA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 453.

Type-species: *Vunga delineata* Walker, 1865, *ibidem* 32: 453, by monotypy. [AUSTRALIA]: [Queensland], Moreton Bay.

Vunga was established in the Notodontidae; it was placed in the Geometridae by Meyrick, 1892, Proc. Linn. Soc. N.S.W. (2) 6: 667.

WANDERBILTIA Rego Barros, 1958, Bolm Mus. nac. Rio de J. (Nov. Ser.) Zool. 177: 1.

Type-species: Wanderbiltia wanderbilti Rego Barros, 1958, ibidem 177: 2, fig., by original designation. Brazil: Itatiaia, Agulhas Negras.

WILEMANUS Nagano, 1916, Bull. Nawa ent. Lab. 1: 2.

NOTO

Type-species: Stauropus bidentatus Wileman, 1911, Trans. ent. Soc. Lond. 1911: 287, pl.30, fig.9, by original designation. JAPAN: Yoshino, Yamato.

See also Chadisroides Matsumura, 1924.

WOERDENIA Snellen, 1872, Tijdschr. Ent. 15: 8, 40. LYMA Type-species: Woerdenia weyenberghi Snellen, 1872, ibidem 15: 8, 40, pl.3, figs 9-12, by monotypy. [CAMEROUN/ GABON/ CONGO/ ANGOLA] Neder-Guinea.

W. weyenberghi is a junior subjective synonym of Hysibada varipes Walker, 1865, the type-species of Hysibada Walker, 1865.

See also Hysibada Walker, 1865.

‡XANTESTHES; misspelling. See Xanthesthes Rambur, 1866.

XANTHARETE Draudt, 1915, in Seitz, Gross-Schmett. Erde 6: 148.

Type-species: *Xantharete stellans* Draudt, 1915, *ibidem* 6: 148, pl.22, row b, by monotypy. Colombia: eastern, Rio Negro.

The date of publication of the part of Seitz containing page 148 was cited on page 145 as 30.iv.1915 in both the German and English editions.

‡XANTHESTES; misspelling. See Xanthesthes Rambur, 1866.

XANTHESTHES Rambur, 1866, Cat. syst. Lépid. Andalousie (2): 228.

Type-species: Xanthesthes albocincta Rambur, 1866, ibidem (2): 229, by monotypy. MADAGASCAR.

‡XANTESTHES; Bryk, 1937, Lepid. Cat. 82: 26.

An incorrect subsequent spelling of Xanthesthes Rambur, 1866.

‡XANTHESTES; Bryk, 1937, Lepid. Cat. 82: 93.

An incorrect subsequent spelling of Xanthesthes Rambur, 1866.

XANTHETIS Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvi, 89, 414.

Type-species: Cyme luzonica Felder & Rogenhofer, 1875, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.139, fig.53, by original designation. [Philippines]: Manilla.

XANTHOARCTIA Travassos, 1951, Revta bras. Biol. 11: 393.

ARCT

Type-species: *Automolis pseudameoides* Rothschild, 1910, *Novit. zool.* 16: 43, by original designation. PERU: Carabaya, R. Huacamayo.

XANTHOCRASPEDA Hampson, 1894, Fauna Br. India (Moths) 2: viii, 121.

Type-species: Nudaria marginata Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 274, by original designation. [INDIA]: Hindostan, southern.

XANTHODONTA Gaede, 1928, in Seitz, Gross-Schmett. Erde 14: 435.

Type-species: *Xanthodonta debilis* Gaede, 1928, *ibidem* 14: 435, pl.72 row a, by monotypy. [Tanzania] Tanganyika: western.

The date of publication of the part of Seitz containing page 435 was cited on page 433 as 30.v.1928 in the German edition; and as 26.xi.1928 in the English edition.

XANTHODULE Butler, 1886, Trans. ent. Soc. Lond. 1886: 384.

Type-species: *Xanthodule semiochrea* Butler, 1886, *ibidem* **1886**: 384, pl.9, fig.1, by monotypy. Australia: [Queensland], near Clermont, Peak Downs.

XANTHODURA Butler, 1880, Ann. Mag. nat. Hist. (5) 5: 384.

Type-species: Xanthodura trucidata Butler, 1880, ibidem (5) 5: 385, by original designation.

Xanthodura was established in the Liparidae, now Lymantriidae; and was transferred to the Geometridae by Swinhoe, 1904, Trans. ent. Soc. Lond. 1904: 546.

XANTHOLOPHA Schaus, 1899, Jl N.Y. ent. Soc. 7: 216.

Type-species: Xantholopha purpurascens Schaus, 1899, ibidem 7: 216, by monotypy. [BRAZIL]: Parana, Castro.

The name Xantholopha probably originated from Hampson but was used and made nomenclaturally available by Schaus prior to its proposal by Hampson, 1900, as the name for a new genus.

XANTHOLOPHA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xv, 89, 356.

Type-species: Xantholopha purpurascens Schaus, 1899, by original designation.

A junior homonym of *Xantholopha* Schaus, 1899. The objective replacement name is *Xantholopha* Schaus, 1899.

XANTHOLOPHA Hampson, 1900. See *Xantholopha* Schaus, 1899.

XANTHOMAENAS Roepke, 1940, Ent. Z., Frankf.a.M. 54: 25.

Type-species: Xanthomaenas singularis Roepke, 1940, ibidem 54: 26, by monotypy. JAVA: south coast, Wijnkoopsbaai.

XANTHOMIS Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xv, 24, 309.

Type-species: Automolis grandis Druce, 1884, Biologia cent.am. (Zool.) Lepid.-Heterocera 1: 75, pl.8, fig.25, by original designation. Costa Rica: Irazu.

XANTHONYGMIA Toxopeus, 1948, Treubia 19: 478. LYMA Type-species: Xanthonygmia oculata Toxopeus, 1948, ibidem 19: 478, by original designation. Borneo: [Kalimantan], Tanah Grogot, Amuntai.

XANTHOPHAEINA Hampson, 1901, Cat. Lepid. Phalaenae Br. Mus. 3: viii, 5, 82.

Type-species: *Phaegoptera levis* Druce, 1899, *Ann. Mag. nat. Hist.* (7) 3: 467, by original designation (but cited as ‡*Phegoptera*, an incorrect subsequent spelling). BRAZIL: southern.

XANTHOPLEURA Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: xviii, 27, 426.

Type-species: Calonotos trotschi Druce, 1884, Biologia cent.am. (Zool.) Lepid.-Heterocera 1: 44, pl.8, fig.8, by original designation. Panama: Chiriqui Volcano.

XANTHYDA Hampson, 1920. See Hyda Walker, 1854.

XENIARA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 978 (key), 1017. NOTO

Type-species: *Xeniara sobria* Walker, 1855, *ibidem* 5: 1018, by monotypy. [Brazil]: Rio de Janeiro.

XENOCLOSTERA Kiriakoff, 1970, Tijdschr. Ent. 113: 123.

Type-species: *Xenoclostera argyrocraspeda* Kiriakoff, 1970, *ibidem* 113: 123, fig.2, by original designation. [SRI LANKA] Ceylon: Maskelyia.

XENOMIGIA Warren, 1906, Proc. U.S. natn. Mus. 30: 413.

DIOP

Type-species: Xenomigia veninotata Warren, 1906, ibidem

30: 414, by original designation. COLOMBIA.

Xenomigia was established in the Cyllopodinae, now Geometridae, Sterrhinae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 406.

XENORMA Prout, 1918, Novit. zool. 25: 397 (key), 404.

Type-species: Phaeochlaena cytheris Druce, 1891, Biologia cent.-am. (Zool.) Lepid.-Heterocera 2: 2, pl.41, fig.11, by original designation. MEXICO: Orizaba, Cuesta de Misantla, COSTA RICA: Caché.

XENORMICOLA Hering, 1928, Dt. ent. Z. Iris 42: 271.

Type-species: Xenormicola prouti Hering, 1928, ibidem 42: 271, by original designation. BOLIVIA: Yungas, San Antonio.

XENOSOMA Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.99, figs 9, 10. PERI

Type-species: Xenosoma nigricosta Felder, 1874, ibidem 2 (Abt.2): pl.99, fig.9, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 431, but cited as Eloria flaviceps Walker, 1865, a species not originally included in Xenosoma. Kirby, however, at the same time placed flaviceps as the senior synonym of nigricosta, a species originally included in Xenosoma. Under the Code, Article 69(a)(iv), this designation constitutes the fixation of the originally included species as the type-species. Guatemala: cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 10.

X. nigricosta is a junior subective synonym of Eloria flaviceps Walker, 1865, List Specimens lepid. Insects Colln Br. Mus.

32: 339.

XEROPTERYX Kiriakoff, 1964. See Spatalina Bryk, 1950.

XULORA Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 979 (key), 1071.

Type-species: Xulora lignosa Walker, 1855, ibidem 5: 1072, by monotypy. [Brazil]: Rio de Janeiro.

XYLECATA Swinhoe, 1904, Trans. ent. Soc. Lond. 1904: 148.

Type-species: Xylecata druna Swinhoe, 1904, ibidem 1904: 148, by PRESENT DESIGNATION. GABON.

XYLINODES Packard, 1864, Proc. ent. Soc. Philad. 3: 366.

Type-species: Xylinodes virgata Packard, 1864, ibidem 3: 367, by monotypy. [U.S.A.]: Cambridge, Michigan, Lansing, Rhode Island, Seekonk.

X. virgata is a junior subjective synonym of Ianassa lignicolor Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5:

See also *Ianassa* Walker, 1855.

YAMATOA Kiriakoff, 1967, in Wytsman, Genera Insect. 217

Type-species: Rosama cinnamomea Leech, 1889, Proc. zool. Soc. Lond. 1888 (4): 637, pl.31, fig.11, by original designation. [JAPAN]: Nagasaki, Ohoyama.

YAZAWAIA Marumo, 1920, J. Coll. Agric. imp. Univ. Tokyo 6: 283 (key), 305.

Type-species: Yazawaia japonica Marumo, 1920, ibidem 6: 306, text-fig.14, pl.23, fig.14, pl.25, figs 4, 8, pl.29, fig.2, pl.35, fig.4, by original designation. [JAPAN]: Shinano, Mt Shirouma.

Y. japonica is a junior subjective synonym of Epinotodonta fumosa Matsumura, 1919, Zool. Mag. Tokyo 31: 76. See also Epinotodonta Matsumura, 1919.

YELVA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1:82.

Type-species: Yelva obscura Birket-Smith, 1965, ibidem 1: 82, fig., by monotypy. NIGERIA: southern, Owenna.

YPIRANGA Rego Barros, 1959. See Lepypiranga Rego Barros, 1966.

ZADADRA Moore, 1878, Proc. zool. Soc. Lond. 1878: 25.

Type-species: Zadadra distorta Moore, 1878, ibidem 1878: 25, by monotypy. INDIA: Darjeeling.

ZADADRINA Kiriakoff, 1954, Revue Zool. Bot. afr. 50: 174.

Type-species: Zadadrina metallica Kiriakoff, 1954, ibidem 50: 175, pl.1, fig.4, pl.2, figs 8, 9, by original designation. ZAIRE: [Lubumbashi] Elisabethville.

ZAEVIUS Dyar, 1910, Zoologica, N.Y. 1: 126. ARCT Type-species: Zaevius calocore Dyar, 1910, ibidem 1: 126, fig.41, no.3, by monotypy. GUYANA: Hoorie.

ZAGARIS Walker, 1855, List Specimens lepid. Insects Colln Br. Mus. 5: 1096.

Type-species: Zagaris crassa Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.99, fig.16 (but included by Walker as 'Zagaris Meteus Cram.' a misidentification; see below), by monotypy. [South AFRICA]: [Cape Province], Knysna, cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls 75-107: 10.

Walker, 1855, cited one originally included nominal species Sphinx meteus Stoll, 1781, in Cramer, Uitlandsche Kapellen (Papillons exot.) 4: 109, pl.347, fig.B. Hampson, 1898, Cat. Lepid. Phalaenae Br. Mus. 1: 149, indicated that Walker had misidentified specimens as *meteus* which in reality belonged to a taxon later named as Zagaris crassa Felder.

Zagaris is a case of a genus that has a misidentified typespecies and so should be referred to the Commission under Article 70(a) of the Code.

ZAMA Walker, 1865. See Zamana Fletcher nom. n. NOTO

ZAMANA Fletcher nom. n. for Zama Walker, 1865. NOTO Type-species: Zama ursipes Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 469, by monotypy (of Zama Walker, 1865). WEST AFRICA.

ZAMA Walker, 1865, List Specimens lepid. Insects Colln Br. Mus. 32: 469.

Type-species: Zama ursipes Walker, 1865, by monotypy. A junior homonym of Zama Herrich-Schäffer, 1855, Syst. Bearbeitung Schmett. Eur. 6: 87, - Lepid., Zygaenidae. The objective replacement name is Zamana Fletcher nom. n.

ZAMOLIS Dyar, 1910, Proc. U.S. natn. Mus. 38: 235. ARCT Type-species: Zamolis noctella Dyar, 1910, ibidem 38: 235,

by original designation. MEXICO: Cuernavaca.

Z. noctella is a junior subjective synonym of Ctenucha dolens Druce, 1897, Biologia cent.-am. (Zool.) Lepid.-Heterocera 2: 360, pl.73, fig.4.

See also Stenucha Hampson, 1901, and Ardonipsa Dyar, 1912.

ZANA Walker, 1856, List Specimens lepid. Insects Colln Br. Mus. 7: 1700.

Type-species: Zana lignosa Walker, 1856, ibidem 7: 1700, by monotypy. INDIA: northern.

‡ZANGRA; misspelling. See Zaranga Moore, 1884. NOTO

‡ZANODA; misspelling. See Zonoda Schaus, 1896. LITH

ZARANGA Moore, 1884, Trans. ent. Soc. Lond. 1884: 357.

Type-species: Zaranga pannosa Moore, 1884, ibidem 1884: 357, by monotypy. INDIA: Punjab, Ambala district.

‡ZANGRA; Matsumura, 1909, Thousand Insects Japan (Suppl.) 1: 65.

An incorrect subsequent spelling of Zaranga Moore, 1884.

ZARFA Walker, 1869, in Chapman, Proc. nat. Hist. Soc. Glasgow 1: 326, 338.

Type-species: Zarfa lunifera Walker, 1869, ibidem 1: 326,

339, by monotypy. Congo.

Z. lunifera is a junior subjective synonym of Naroma signifera Walker, 1856, the type-species of Naroma Walker, 1856.

Zarfa was established in the Liparidae, now Lymantriidae; it was placed in the Bombycidae by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 716; and returned to the Lyman-triidae by Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 133. See also Naroma Walker, 1856.

ZASTONIA Herrich-Schäffer, [1858] 1850-1858, Samml. neuer oder wenig bekannter aussereur. Schmett. 1: 67 (but attributed to Boisduval, an incorrect authorship).

Type-species: Bombyx panda Boisduval, 1847, in Delegorgue, Voyage Afr. australe 2: 600, by monotypy. [South Africa]: [Natal], [Durban] Port Natal.

HENOSIS Wallengren, 1865, K. svenska Vetensk Akad. Handl. (N.F.) 5 (4): 51.

Type-species: Bombyx panda Boisduval, 1847, by monotypy. A junior objective synonym of Zastonia Herrich-Schäffer, [1858].

‡ZATHREPHA; misspelling. See Zatrephes Hübner, [1819]. ARCT

ZATREPHES Hübner, [1819] 1816, Verz. bekannter Schmett.: 171.

Type-species: Phalaena nitida Cramer, 1780, Uitlandsche Kapellen (Papillons exot.) 3: 147, 176, pl.274, fig.F, by subsequent designation by Kirby, 1892, Synonymic Cat. Lepid. Heterocera 1: 196. SURINAM.

P. nitida is dated from the wrapper of the part and not from the title-page of the volume, 1782.

‡ZATHREPHA; Hübner, [1826] 1816, Verz. bekannter Schmett. (Anzeiger): 29.

An incorrect subsequent spelling of Zatrephes Hübner, [1819].

ZAVANA Griveaud, [1976] 1975, Bull. Soc. ent. Fr. 80: 227.

Type-species: Dasychira acroleuca Hering, 1926, in Seitz, Gross-Schmett. Erde 14: 176, pl.27, row f, by original designation. MADAGASCAR: Mohanoro.

ZELICA Hübner, [1825] 1816, Verz. bekannter Schmett.: 396.

Type-species: Phalaena zelica Stoll, 1790, in Cramer, Uitlandsche Kapellen (Papillons exot.) (Aanhangsel): 73, pl.16, figs 2, 2C, 2D (but included by Hübner as ‡zelia, an incorrect subsequent spelling), by monotypy. SURINAM.

P. zelica is dated from the wrapper of the part, not from the title-page of the volume which is dated 1791.

See also Rosema Walker, 1855.

ZELLATILLA Dyar, 1914, Insecutor Inscit. menstr. 2: 111.

Type-species: Zellatilla columbia Dyar, 1914, ibidem 2: 111. by monotypy. Cuba: [Isla de Pinos] Isle of Pines, Columbia; and unstated locality.

ZELOMERA Butler, 1882, Cistula ent. 3: 25. NOTO Type-species: Zelomera imitans Butler, 1882, ibidem 3: 26. by monotypy. MADAGASCAR.

ZERAFIA Strand, 1915, Arch. Naturgesch. 80 (A) 10: 102.

Type-species: Zerafia drymonides Strand, 1915, ibidem 80 (A) 10: 102, by original designation. [SUDAN]: Bahr el Abiad, Meschra Zeraf.

ZIGIRA Walker, [1865] 1864, List Specimens lepid. Insects Colln Br. Mus. 31: 152. Type-species: Zigira quadrata Walker, [1865] 1864, ibidem 31: 153, by monotypy. [COLOMBIA]: Bogota.

ZOBIDA Birket-Smith, 1965, Pap. Fac. Sci. Haile Sellassie I Univ. (C) 1: 31.

Type-species: Eilema trinitas Strand, 1912, Arch. Naturgesch. 78 (A) 7: 181, by original designation. [CAMEROUN] Kamerun, Togo: 'Bismarckburg'.

ZONODA Schaus, 1896, Jl N.Y. ent. Soc. 4: 150. Type-species: Zonoda fasciata Schaus, 1896, ibidem 4: 150 (but cited as ‡Zanoda, an incorrect (of a multiple) original spelling), by subsequent designation by Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: 363. BRAZIL: São Paulo.

As first reviser, Strand, 1922, in Wagner, Lepid. Cat. 26: 714, chose the spelling Zonoda from the multiple original spellings Zonoda and ‡Zanoda.

‡ZANODA; Schaus, 1896, Jl N.Y. ent. Soc. 4: 150.

An incorrect (of a multiple) original spelling of Zonoda Schaus, 1896.

ZONOSOMA Butler, 1881. See Nyctemera Hübner, [1820]. ARCT

ZOOTE Hübner, [1820]. See Arctia Schrank, 1802. ARCT

ZUNACETHA Walker, 1863, List Specimens lepid. Insects Colln Br. Mus. 27: 133.

Type-species: Zunacetha bipartita Walker, 1863, ibidem 27: 134, by monotypy. HONDURAS.

Zunacetha was established in the Phycidae, now Pyralidae; it was placed in the Dioptidae by Prout, 1918, Novit. zool. 25: 406.

ZYGAENOPSIS Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.19.

Type-species: Zygaenopsis squamicornis Felder, 1874, ibidem 2 (Abt.2): pl.102, fig.19, by monotypy. [Brazil]: Amazon R., cited by Felder, 1875, ibidem 2 (Abt.2) Erklärung to pls

Z. squamicornis is a junior subjective synonym of Euchromia contenta Walker, 1854, List Specimens lepid. Insects Colln Br. Mus. 1: 258.

‡ZYGAENOPSIS Swinhoe, 1892. See Zygaenosia Hampson, 1900.

ZYGAENOSIA Hampson, 1900, Cat. Lepid. Phalaenae Br. Mus. 2: xvii, 86, 423.

Type-species: Zygaenopsis fuscimarginalis Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 61, by original designation (for Zygaenopsis Swinhoe, 1892). [New Guinea]: [Irian Jaya], [north-western], [Dore] Dorey.

Zygaenosia Hampson was established as the objective replacement name for Zygaenopsis Swinhoe, 1892.

ZYGAENOPSIS Swinhoe, 1892, Cat. east. and Aust. Lepid. Heterocera Colln Oxf. Univ. Mus. 1: 61.

Type-species: Zygaenopsis fuscimarginalis Swinhoe, 189?, by original designation.

A junior homonym of Zygaenopsis Felder, 1874, in Felder & Rogenhofer, Reise öst. Fregatte Novara (Zool.) 2 (Abt.2): pl.102, fig.19, – Lepid., Arctiidae. The objective replacement name is Zygaenosia Hampson, 1900.

Index to Species-group Names

Species-group names are listed with author and original genus. The names following a semicolon are the genus-group names, listed in the catalogue in bold type, under which the species-group names will be found.

abbreviata Kenrick, Dasychira; Fanala abdominalis Grote, Pygarctia; Pygarctia abnormis Hampson, Sphecosoma; Abnormipterus abrupta Hübner, Hypocrita; Ephialtias absconditana Walker, Tospitis; Trischalis acasia Schaus, Somera; Acasina accipiter Schaus, Hapigia; Rhapigia accurata van Eecke, Neosiccia; Neosiccia acharista West, Spatalia; Coscodaca achatina Smith, Phalaena; Parorgyia acosmeta Collenette, Dasychira; Varatra acrea Drury, Phalaena; Estigmene acrobela Turner, Themerastis; Antithemerastis acroleuca Hering, Dasychira; Zavana acroxantha Perty, Glaucopis; Mochloptera acuminata Matsumura, Neofentonia; Neofentonia acuta Hampson, Halysidota; Lepidonewtonia acuta Matsumura, Togaritensha; Togaritensha acuta Walker, Turnaca; Turnaca admirabilis Cramer, Phalaena; Idalus admirabilis Staudinger, Lophopteryx; Hagapteryx adscitina Dognin, Ardonissa; Ardonissa adusta Felder, Dipteromorpha; Dipteromorpha adversata Schaller, Phalaena; Trypheromera aedumena Turner, Heliosa; Aclytophanes aegrotum Swinhoe, Leptosoma; Exitelica aenea Kiriakoff, Oreoceryx; Oreoceryx aenescens Hampson, Metareva; Metareva aenone Butler, Halysidota; Syntarctia aeola Bryk, Harpandrya; Harpandrya aequalis Walker, Alpenus; Alpenus aequalis Walker, Eugoa; Eugoa aequata Walker, Satara; Satara aequinoctialis Walker, Gnophaela; Gnophaela aequipars Walker, Nystalea; Proelymiotis aeschra Hampson, Sminthopses; Sminthopses aetheria Strand, Ruanda; Ruanda aethiopica Laporte, Dorstiana; Dorstiana aethiopica Laporte, Rougeotia; Rougeotia affine Butler, Callicereon; Callicereon affinis Hering, Euproctillopsis; Euproctillopsis affinis Rothschild, Pseudapiconoma; Balacrella afflicta Walker, Glaucopis; Thrinacia afghanus Ebert, Bifurcifer; Bifurcifer africana Holland, Lepasta; Neomardara africana Holland, Maguza; Pseudoscrancia afroalpina Laporte, Michelliana; Michelliana aganais Felder, Homoeognatha; Homoeognatha aganice Fawcett, Bithra; Galtara agaristoides Boisduval, Phaegorista; Phaegorista aglaura Cramer, Phalaena; Thebrone aglaurus Kiriakoff, Prosphoroplitis; Prosphoroplitis aglone Herrich-Schäffer, Lobeza; Lobeza agramma Hampson, Phyllaliodes; Phyllaliodes agyllia Hampson, Glaucosia; Glaucosia alba Druce, Symmerista; Kalkoma alba Moore, Caltura; Caltura alba Moore, Dondera; Dondera alba Moore, Dura; Dura albata Packard, Clemensia; Clemensia albertae Dyar, Dodia; Dodia albescens Rothschild, Pasteosia; Prosiccia albicancellata Burmeister, Arctia; Caridarctia albicans Walker, Carales; Carales albicans Walker, Rethona; Pseudorethona albicauda Bryk, Phalera; Periphalera

albicincta Schaus, Eupyra; Mydropastea albicollis Felder, Eucyrta; Eucyrta albicollis Pagenstecher, Hypocrita; Caprimima albicosta Hampson, Diduga; Androstigma albicosta Hübner, Noctua; Symmerista albicosta Matsumura, Mimodonta; Mimodonta albicostata Hampson, Demolis; Demolis albida Hampson, Anaemosia; Anaemosia albida Walker, Nudaria; Phaneropseustis albida Walker, Uxia; Uxia albidator Walker, Halysidota; Psychophasma albidilinea Gaede, Eurystauropsis; Eurystauropsis albidilinea Gaede, Eutrotonotus; Eutrotonotus albifinis Walker, Sidyma; Sidyma albifrons Smith, Phalaena; Edema albifrons Walker, Phusiana; Phusiana albifusa Wileman, Pydna; Loxobireta albilinea Holland, Thamnocera; Thamnocera albilunula Collenette, Dasychira; Rahona albilunulata Karsch, Bathmochtha; Bathmochtha albimacula Hampson, Gargetta; Stictogargetta albimacula Staudinger, Notodonta; Urodonta albimacula Wallengren, Palasea; Palasea albimacula Warren, Macroneurodes; Macroneurodes albimaculata Okano, Peridea; Pulia albiplaga Walker, Euchromia; Metriophyla albiplaga Warren, Momonipta; Momonipta albiplagiata Gaede, Peratodonta; Bernardita albipuncta [Denis & Schiffermüller], Noctua; Hyphilare albipuncta Gaede, Desmeocraera; Epimetula albipuncta Hampson, Thylacoptera; Thylacoptera albipuncta Schaus, Hyperthaema; Hyperthaema albipunctella Kiriakoff, Synete; Synete albisigna Walker, Glaucopis; Pheia albistriga Moore, Niganda; Poncetia albiventus Druce, Neacerea; Neacerea albocalccolata Bryk, Phaleromimus; Phaleromimus albocincta Rambur, Xanthesthes; Xanthesthes albocinerea Moore, Ghoria; Ghoria alboflavida Bryk, Pydna; Palessa albolineata Strand, Scrancia; Gargettoscrancia albomaculata Kiriakoff, Pachyceryx; Pachyceryx albonotata Walker, Allata; Brykia albosigma Druce, Lochmaeus; Tachuda albosparsatus Kenrick, Lerodes; Paralerodes albovittata Hampson, Phycidopsis; Phycidopsis albulata Herrich-Schäffer, Mieza; Paramulona alciphron Cramer, Phalaena; Euimata aliaria Druce, Trichura; Riccia aliena Walker, Euchromia; Pompiliodes aliena Wallengren, Creagra; Creagra allocota Bryk, Argyrothyris; Argyrothyris almon Cramer, Sphinx; Autochloris alpina Bellier, Clostera; Rhegmatophila alpina Quensel, Bombyx; Acerbia alpium Osbeck, Phalaena; Diphteramoma alterna Walker, Setina; Tigrioides alternata Romieux, Gaedea; Gaedeina altrix Stoll, Phalaena; Tifama amabilis Hering, Pyrrhopteryx; Pyrrhopteryx amamiana Nakamura, Pseudofentonia; Nipponotensha amanda Boisduval, Euchelia; Alytarchia amanda Staudinger, Ocnerogyia; Ocnerogyia amasis Cramer, Phalaena; Rhodogastria amata Druce, Opharus; Pseudopharus amatame Dyar, Ribaldia; Ribaldia

amazonum Felder, Microrgyia; Microrgyia amazoula Boisduval, Thyretes; Epitoxis ambages Strand, Schalidomitra; Schalidomitra amblyiodes Turner, Scythrophanes; Hobartina amica Stoll, Phalaena; Calodesma amicus White, Agagles; Agagles ammoscia Lower, Microstola; Microstola amnaea Swinhoe, Aemene; Hyposiccia amoureli Dognin, Areva; Paragylla ampla Walker, Enome; Enome ampla Walker, Terna; Terna ampliata Butler, Dasychira; Vietteria amplificata Gaede, Phalera; Phusiana amygdaloides Walker, Melora; Melora analalava Griveaud, Erika; Erika anapheles Nye, Sideridis; Sideridis anartoides Walker, Mosoda; Mosoda anartoides Walker, Teia; Teia anastasia Oberthür, Naclia; Stictonaclia anastomosis Linnaeus, Phalaena; Ichthyura anceps Goeze, Phalaena; Peridea anceps Walker, Cerasana; Cerasana ancilla Linnaeus, Phalaena; Dysauxes angasii Angas, Agarista; Cremnophora angasii Walker, Apina; Cremnophora anguina Smith, Phalaena; Dasylophia angulata Gaede, Lophopteryx; Afropteryx angulata Gaede, Pseudobarobata; Pseudobarobata angulifer Schaus, Eugonosia; Eugonosia angulinea Schaus, Dunama; Dunama anguliscripta Lucas, Chiriphe; Goniosema angulosa Walker, Lyces; Lyces angustata Möschler, Sphecosoma; Pleurosoma angustipennis Joannis, Stenopterosia; Stenopterosia angustipennis Kiriakoff, Microchadisra; Microchadisra angustipennis Matsumura, Betashachia; Betashachia angustipennis Rothschild, Anaphleps; Anaphleps ania Hering, Dasychira; Jabaina annulata Gaede, Stauroplitis; Stauroplitis annulata Schaus, Cresera; Cresera annulatum Boisduval, Leptosoma; Nyctemera annulipes Berg, Thosea; Tecmessa anomala Benjamin, Acsala; Acsala anomala Bowring, Epipyrops; Epipyrops anomala Butler, Lechriolepis; Lechriolepis anomala Butler, Maenoleneura; Maenoleneura anomala Moore, Sorocaba; Sorocaba anomala Walker, Peroma; Peroma anomala Westwood, Epipyrops; Epipyrops ansorgei Kirby, Neuroxena; Neuroxena antennata Walker, Laora; Laora anthracina Walker, Euchromia; Anycles antica Walker, Arestha; Arestha antica Walker, Dediama; Dediama antica Walker, Lacida; Lacida antica Walker, Lithosia; Brunia antica Walker, Tanada; Tanada antipolo Semper, Syntpmoides; Symphlebomis antiqua Linnaeus, Phalaena; Orgyia antongilensis Viette, Desmeocraera; Exantongila apatela Kiriakoff, Metriaeschra; Metriaeschra apateloides Holland, Dasychira; Deinarchia apicalis Herrich-Schäffer, Charidea; Metriophyla apicalis Kiriakoff, Allodonta; Allodontina apicalis Moore, Stauropus; Grangula apicalis Rothschild, Teucer; Mesocerea apicalis Walker, Arna; Arna apicalis Walker, Bembina; Bembina apicalis Walker, Mosara; Mosara apicalis Walker, Nyctemera; Podomachla apicalis Walker, Pida; Pida apicalis Walker, Sphetta; Sphetta

apicalis Zeller, Calligenia; Chionosia apiculatus Rothschild, Stauropus; Epistauropus apostata Schaus, Tessellota; Castronia appendiculatus Herrich-Schäffer, Creatonotus; Hyperandra apsara Moore, Dasychira; Gazalina apulana Schaus, Procolax; Procolax apulus Cramer, Phalaena; Colax arauna Wallengren, Aristodaema; Aristodaema arborifera Butler, Lacydes; Paralacydes archias Stoll, Sphinx; Eucereon archonta Kiriakoff, Odontoperas; Odontoperas arctata Walker, Nyctemera; Arctata arctata Walker, Pseudosphex; Sphecops arctioides Butler, Spilobotys; Spilobotys arctipennis Butler, Lygrotes; Lygrotes arctipennis Holland, Norraca; Galanthella arcuata Alphéraky, Urodonta; Urodontopsis arcuata Kiriakoff, Hypostaurella; Hypostaurella arcuata Kiriakoff, Somerina; Somerina arculifera Felder, Eucreagra; Eucreagra areolata Walker, Dioptis; Hyrmina arge Drury, Phalaena; Mimarctia argentata Butler, Dasychira; Calliteara argentata Moore, Lophopteryx; Spatalina argentata Packard, Halysidota; Euschausia argentata Walker, Teara; Marane argentea Fleming, Pseudomacroptila; Pseudomacroptila argentea Hampson, Procanthia; Procanthia argentea Janse, Bicelluphora; Bicelluphora argentea Walker, Redoa; Himala argentea Walker, Salapola; Salapola argenteodivisa Kiriakoff, Bireta; Struba argenteogutta Strand, Neritonaclia; Neritonaclia argenteomaculata Aurivillius, Notodonta; Graphidura argenteopunctata Möschler, Euxoga; Euxoga argentifascia Hampson, Eutornopera; Eutornopera argentifera Moore, Heterocampa; Pseudofentonia argentifera Walker, Allata; Allata argentilinea Druce, Didugua; Didugua argentina [Denis & Schiffermüller], Bombyx; Spatalia argentinensis Rothschild, Cisthene; Vianania argentopicta Matsumura, Togarishachia; Togarishachia arginalis Hampson, Areas; Callindra argus Walker, Hypercompe; Sebastia argynnis Fabricius, Zygaena; Marmax argyraspis Holland, Ptilura; Ptilura argyria Butler, Tiauspa; Ctianopha argyrochlora Carcasson, Tuerta; Pseudotuerta argyrocraspeda Kiriakoff, Xenoclostera; Xenoclostera argyrophora Hampson, Spatalia; Baliopteryx arida Walker, Clathe; Clathe arnaui Orfila, Lepidojulia; Lepidojulia arne Cramer, Phalaena; Eudmoe aroa Bethune-Baker, Lasioceros; Lasioceros aroa Bethune-Baker, Paraceryx; Paraceryx aroata Schaus, Betola; Betola arontes Stoll, Sphinx; Pelochyta arsilonchoides Möschler, Talmenia; Talmenia artaxidia Butler, Miltochrista; Nudina arthemis Schaus, Hemipecteros; Hemipecteros asbolina Turner, Epicoma; Epicomana asiatica Krulikovski, Sarrothripus; Dufayella aspasta Hering, Hoppiana; Hoppiana aspersa Grote, Alexicles; Alexicles aspersa Walker, Lithosia; Euconosia aspersum Felder, Tearosoma; Tearosoma assimilis Hering, Myonia; Isionda astarte Doubleday, Heterocampa; Heterocampa astreus Drury, Sphinx; Amerila astrifer Butler, Lophocoleus; Tholocoleus astur Cramer, Phalaena; Senia asurella Strand, Agylloides; Agylloides

atalopa Turner, Destolmia; Commonia ataracta Hampson, Ionthas; Ionthas athola Turner, Oreopola; Oreopola atima Collenette, Rhypotoses; Rhypotoses atomosa Walker, Halysidota; Euhalisidota atra Gaede, Pheosina; Pheosina atricollis Kiriakoff, Bilulua; Bilulua atricornis Wallengren, Asinusca; Asinusca atrifasciata Hampson, Psalisodes; Psalisodes atrifrons Walker, Gaugamela; Gaugamela atrifulva Hampson, Trichocerosia; Trichocerosia atrinotata Butler, Antiphella; Pirgula atripennis Grote, Dahana; Dahana atriplaga Staudinger, Notodonta; Lophocosma atrizona Hampson, Pseudlepista; Pseudlepista atroguttata Aurivillius, Leucoperina; Leucoperina atrosquama Walker, Gogana; Choerotricha atrovittatus Bremer, Asteroscopus; Shaka atuntseica Daniel, Asuridoides; Asuridoides auge Linnaeus, Sphinx; Cosmosoma aulaea Geyer, Arachnis; Arachnis aulica Linnaeus, Phalaena; Hyphoraia aurantiaca Hampson, Stenilema; Stenilema aurantiaca Hübner, Eubaphe; Holomelina aurantiaca Rothschild, Lambula; Lambulosia auranticeps Hampson, Chrysoscota; Chrysoscota aurantiistriga Kiriakoff, Bireta; Subniganda aurata Moore, Niganda; Midasia aureobrunnea Bryk, Mimesisomera; Mimesisomera aureogrisea Rothschild, Automolis; Regobarrosia aureola Hübner, Bombyx; Eilema aureopsis Hering, Porthesaroa; Porthesaroa aureorubra Hampson, Chrysallactis; Chrysallactis aurifera Klages, Pseudosphex; Sphecomimax aurifera Rothschild, Oeonistis; Oeonosia auriflamma Geyer, Josia; Scea auriginea Franclemont, Synorthodes; Synorthodes auripennis Matsumura, Eushachia; Eushachia auripes Butler, Leucoma; Ivela auriplaga Kiriakoff, Graphidurina; Graphidurina aurivilliusi Kiriakoff, Macronadata; Archinadata aurora Kiriakoff, Bireta; Oraura aurora Smith, Phalaena; Hyparpax aurorina Herrich-Schäffer, Halurgia; Halurgia aurosigna Hampson, Plusiogramma; Plusiogramma austera [Denis & Schiffermüller], Bombyx; Ochrostigma azurea Walker, Dioptis; Euagra

baibarana Matsumura, Damatoides; Damatoides bakeri Schaus, Siopastea; Siopastea bangeni Laporte, Dargeochaeta; Dargeochaeta banksiae Lewin, Bombyx; Danima barbicostata Hampson, Chrysomesia; Chrysomesia barnesi Kiriakoff, Diopeithes; Diopeithes basalis Moore, Stauropus; Neostauropus basalis Walker, Argila; Argila basalis Walker, Euchromia; Pseudosphenoptera basalis Walker, Perina; Perina basalis Walker, Phastia; Phastia basifera Walker, Rhuda; Rhuda basifusca Walker, Nemeta; Nemeta basigera Walker, Cossa; Cossa basigutta Walker, Noleca; Noleca basilinea Hering, Scoturopsis; Scoturopsis basilinea Hulstaert, Dichrostoptera; Dichrostoptera basilinea Wileman, Notodonta; Fusadonta basilutea Walker, Glaucopis; Hyda basimaculatum Matsumura, Medama; Medama basinota Moore, Lithosia; Simareea

basinotata Wileman, Notodonta; Norracoides basiplaga Felder, Nyctochroa; Nyctochroa basipuncta Gaede, Turnacoides; Turnacoides basipuncta Schaus, Anita; Anita basistriata Kiriakoff, Egonociades; Egonociades basitriens Walker, Notodonta; Chatfieldia basivacua Walker, Dediama; Calapana batesii Felder, Josiodes; Josiodes batialis Walker, Illice; Illice baucis Dalman, Charidea; Lepidolutzia beanii Neumoegen, Antarctia; Neoarctia bela Swinhoe, Pydna; Rosiora belae Grote, Didasys; Didasys bella Bethune-Baker, Omestia; Omestia bella Bethune-Baker, Peratodonta; Bostrychogyna bella Guérin-Méneville, Glaucopis; Cyanopepla bellatrix Dalman, Callimorpha; Amphicallia bellatrix Stoll, Phalaena; Crinodes belli Field, Parvicincia; Parvicincia benga Holland, Scalmicauda; Scalmicauda bengueta Schaus, Midara; Midara benguetensis Wileman, Tinoliodes; Tinoliodes bermejona Schaus, Pagrasana; Pagrasana bettoni Butler, Trotonotus; Trotonotus bicellulata Kaye, Crinophora; Crinophora bicolor Distant, Chadisra; Sarimarais bicolor Hulstaert, Diadesmola; Diadesmola bicolor Kiriakoff, Graptophalera; Graptophalera bicolor Möschler, Campylona; Campylona bicolor Rothschild, Asura; Phlogomera bicolor Rothschild, Scaptesyle; Pseudoscaptia bicolor Viette, Pseudohypsoides; Pseudohypsoides bicolor Walker, Euchromia; Hydrusa bicolor Walker, Rhagonis; Rhagonis bicolor Walker, Sarapus; Asparus bicolor Walker, Sorama; Sorama bicolora Bethune-Baker, Acco; Acco bicolora Boisduval, Lithosia; Scoliacma bicolora Sulzer, Phalaena; Esthema bicolora Walker, Decimia; Decimia bicolorana Bethune-Baker, Garudinodes; Garudinodes bicoloria [Denis & Schiffermüller], Bombyx; Leucodonta bicosta Walker, Lithosia; Palaeosia bidens Oberthür, Semidonta; Sinodonta bidensis Kaye, Acutia; Atucia bidentata Walker, Nerice; Nerice bidentatus Wileman, Stauropus; Wilemanus bieti Oberthür, Chelonia; Euleechia bifacies Walker, Glissa; Glissa bifacies Walker, Nemeta; Nemeta bifacies Walker, Sidana; Sidana bifasciata Köhler, Neobourquinia; Neobourquinia bigutta Walker, Lysceia; Lysceia bilinea Schaus, Trochuda; Trochuda bilinea Walker, Setina; Grammarctia bilineata Hampson, Graphosia; Graphosia bilineata Hampson, Phalera; Antiphalera bilineata Packard, Cecrita; Seirodonta bilineatus Hampson, Siculifer; Siculifer biloba Oberthür, Drymonia; Semidonta biluna Walker, Tepilia; Tepilia bilunata Toulgoët, Cristulosia; Cristulosia bimaculata Moore, Challa; Challa binotata Butler, Leptocneria; Leptocneria binotata Kiriakoff, Chalepa; Chalepa bipars Butler, Hypsoides; bipars Walker, Chadisra; Chadisra bipars Walker, Stonia; Stonia bipartita Butler, Nerice; Nericoides bipartita Matsumura, Stenoshachia; Stenoshachia bipartita Walker, Zunacetha; Zunacetha biplagata Bethune-Baker, Parascaptia; Parascaptia

biplagella Butler, Teulisna; Macotasa bipuncta Hampson, Pachycerosia; Pachycerosia bipuncta Hampson, Schistophleps; Schistophleps bipuncta Hering, Suarezia; Salvatgea bipunctata Kenrick, Gonepteronia; Gonepteronia bipunctata Okano, Pseudofentonia; Pantherinus bipunctata Walker, Adrallia; Adrallia bipunctata Walker, Andraca; Andraca bipunctigera Wallengren, Lithosia; Lexis bipunctoides Toulgoët, Philenora; Parexilisia birmalina Bryk, Spatalina; Spatalina bisecta Rothschild, Scalmicauda; Tmetopteryx bisignata Kiriakoff, Antiscranciola; Antiscranciola bisignata Kiriakoff, Bugandita; Bugandita bistigmalis Strand, Anthelymantria; Anthelymantria bitia Stoll, Phalaena; Biotes bivittata Clemens, Ectypia; Ectypia bivittata Kiriakoff, Gluphisiodes; Gluphisiodes bivittata Kiriakoff, Schoutenia; Schoutenia bivittata Strand, Anotodonta; Anotodonta blanda van Eecke, Mellona; Mellona boeticum Rambur, Trichosoma; Pachylischia boisduvalii Newman, Oenosandra; Oenosandra boisduvallii van der Hoeven, Glaucopis; Ambryllis bombycina Oberthür, Notodonta; Phalerodonta bombycoides Walker, Bagisara; Caroia bostrychonota Tams, Seudyra; Sarbissa bourgognei Kiriakoff, Apisa; Parapisa bracata Draudt, Osmocneme; Osmocneme braco Herrich-Schäffer, Haematerion; Haematerion braconoides Walker, Glaucopis; Ilipa bractea Felder, Nystalea; Lepasta bracteola Geyer, Phaeochlaena; Brachyglene branickii Oberthür, Uropus; Urodontoides braueri Bryk, Tessmannia; Tessmannia braziliensis Schaus, Mintopola; Mintopola briachisia Schaus, Stauropus; Briachisia brinoni Holloway, Colnettia; Colnettia brodea Schaus, Syntrichura; Urolasia broidricci Dyar, Scevesia; Scevesia brontonepha Collenette, Leucoma; Homaroa brunnea Aurivillius, Peratodonta; Peratodonta brunnea Janse, Eurystaura; Eurystaura brunnea Schaus, Pseudhapigia; Pseudhapigia brunneipennis Hering, Metarctia; Pterophaea brunneoviridans Dognin, Synalamis; Synalamis brunnescens Bartel, Ischnarctia; Ischnarctia brunnescens Rothschild, Scoliacma; Scoliosia bucephala Linnaeus, Phalaena; Phalera buceros Kiriakoff, Rheneina; Rheneina bufo Guenée, Barydia; Barydia butleri Field, Paracincia; Paracincia butleri Mabille, Syntomis; Dubianaclia

cadaverosa Grote, Pareuchaetes; Pareuchaetes cadburyi Franclemont, Comachara; Comachara cadmia Guenée, Hemiceras; Hemiceras caecina Druce, Lithosia; Nephelosia caelicolor Walker, Seripha; Seripha caerulea Druce, Holophaea; Holophaea caeruleifascia Walker, Balacra; Balacra caeruleocephala Linnaeus, Phalaena; Diloba caerulescens Butler, Scaptesyle; Atopomima caerulescens Hampson, Syntomimorpha; Syntomimorpha caesarea Goeze, Phalaena; Epatolmis caffra Walker, Siccia; Siccia caja Linnaeus, Phalaena; Arctia calcarata Walker, Euchromia; Orcynia caledonica Holloway, Pseudocoarica; Pseudocoarica

californiae Walker, Nemeophila; Leptarctia californica Packard, Phryganidia; Phryganidia caliginosa Walker, Blavia; Blavia calligramma Walker, Mardara; Ramadra callipyga Wallengren, Munychia; Munychia callistia Hampson, Chrysochlorosia; Chrysochlorosia callisto Walker, Apina; Apina calocore Dyar, Zaevius; Zaevius calpe Felder, Platyodonta; Platyodonta camelina Linnaeus, Phalaena; Ptilodon camelinerdes Walker, Naprepa; Naprepa camerunica Kiriakoff, Belisaria; Belisaria cana Walker, Repa; Repa cancellata Burmeister, Halysidota; Tessellota candida Hering, Eopirga; Eopirga canescens Walker, Apisa; Apisa caniola Hübner, Bombyx; Eilema caniplaga Walker, Cymatophora; Ellida canosparsa Walker, Epicoria; Epicoria cantiana Schaus, Fentonia; Pellewia capillata Wallengren, Simyra; Polienus capucina Linnaeus, Phalaena; Ptilodon capys Cramer, Sphinx; Patreliura carama Druce, Eustema; Eustemides carbonaria [Denis & Schiffermüller], Geometra; Parascotia caricae Fabricius, Noctua; Psephea carmelita Esper, Phalaena; Odontosia carmelitaria Guenée, Ceratonyx; Sorocaba carmelitoides Guenée, Canodia; Canodia carnea Weymer, Antiloba; Antiloba carneola Moore, Barhona; Barhona carnipicta Butler, Ammatho; Asuridia carteri Druce, Antheua; Tricholoba caryae Harris, Lophocampa; Lophocampa cassandra Linnaeus, Sphinx; Saurita castanea Rothschild, Ichthyura; Erythroclostera castaneola Hampson, Brachiosia; Brachiosia castor Kiriakoff, Ptilurodes; Ptilurodes catilina Cramer, Phalaena; Chetone catocalina Felder, Discophlebia; Discophlebia catocalina Walker, Clisobara; Clisobara catocaloides Herrich-Schäffer, Catochria; Catochria catoxantha Felder, Inopsis; Inopsis caudata Collenette, Leptepilepta; Leptepilepta caudata Kiriakoff, Rodneya; Rodneya caudata Schaus, Bryantia; Bryantia caudatissima Strand, Scranciana; Scranciana caudatula Felder, Anaphlebia; Anaphlebia caudatula Kiriakoff, Amata; Miracidion caurensis Klages, Pseudargyroeides; Pseudargyroeides cedonulli Stoll, Phalaena; Parathyris celadon Cramer, Phalaena; Eucyane celaena Turner, Themerastis; Themerastis celaenocera Collenette, Sevastopuloa; Sevastopuloa celebensis Kiriakoff, Loda; Loda celebensis Roepke, Miltasura; Miltasura celenna Schaus, Trichomelia; Trichomelia celina Boisduval, Calepidos; Calepidos celsicola Toulgoët, Seydelia; Seydelia centralis Herrich-Schäffer, Notodonta; Dicentria centristicta Hampson, Pheosia; Pseudoteleclita ceramboides Herrich-Schäffer, Correbia; Correbia cerurata Dognin, Eunaduna; Eunaduna ceruroides Walker, Blera; Sericochroa cervicalis Walker, Asura; Asura cervina Lucas, Scoliacma; Meteura cervina Moore, Baryaza; Baryaza cervina Walker, Lebadia; Lebadia cervina Walker, Repena; Repena cessogae Schaus, Pseudaethria; Pseudaethria ceylanica Felder, Migoplastis; Migoplastis chadisroides Kiriakoff, Parascrancia; Parascrancia

chalciope Hübner, Isanthrene; Nyridela chalcocraspedon Dyar, Chalcopasta; Cuahtemoca chalybea Hübner, Uranophora; Uranophora chalybsa Hampson, Pseudmelisa; Pseudmelisa chiguinda Druce, Bepara; Heliactinidia chiloides Walker, Teulisna; Corcura chionora Meyrick, Lithosia; Hesychopa chloana Holland, Somera; Enomotarcha chloauchena Holland, Somera; Pittheides chloe Holland, Somera; Stauropussa chloropasta Turner, Syntypistis; Syntypistis chloropyga Walker, Hypsa; Agape choerotricha Felder, Micromorphe; Micromorphe cholerota Meyrick, Chlorogenia; Chlorogenia chordigera Felder, Oligoclona; Oligoclona chrysargyria Hampson, Caripodia; Caripodia chrysis Hübner, Poecilosoma; Poecilosoma chrysogastra Perty, Glaucopis; Corematura chrysogastrides Draudt, Paradinia; Paradinia chrysogramma Hering, Dasychirinula; Dasychirinula chrysolopha Kollar, Liparis; Oligoclona chrysorrhoea Linnaeus, Phalaena; Euproctis chrysotis Hampson, Metallosia; Metallosia ciliosa Felder, Physetocneme; Physetocneme cincticollis Felder, Diabaena; Diabaena cinderella Kiriakoff, Dasychirella; Dasychirella cinderella Schaus, Parapalosia; Parapalosia cineracea Kiriakoff, Archistilbia; Archistilbia cineraria Kiriakoff, Pseudofentonia; Antifentonia cinerascens Kiriakoff, Thaila; Thaila cinerascens Walker, Halysidota; Lepidozikania cinerascens Walker, Tadana; Tadana cinerea Butler, Peridea; Suzukiana cinerea Janse, Taeniopteryx; Metopteryx cinerea Schaus, Afilia; Afilia cingalesa Walker, Panassa; Panassa cingulata Walker, Ituna; Tuina cinnamomea Leech, Rosama; Yamatoa cinnamomea Wallengren, Hexaneura; Hexaneura circulosa Gaede, Desmeocraera; Crestonica circumcincta Saalmüller, Notodonta; Rhenea circumdata Grote, Agrotis; Trichofeltia circumdata Walker, Etobema; Etobema circumscripta Butler, Drymonia; Microhoplitis citana Schaus, Utidava; Utidaviana citrago Toulgoët, Metexilisia; Metexilisia citricostata Hampson, Proschaliphora; Proschaliphora citrina Sepp, Phalaena; Thalesa citrina Wileman, Gangarides; Gangaridopsis cladara Turner, Nesiotica; Nesiotica clara Stoll, Phalaena; Notoplusia clarus Grote & Robinson, Vanessodes; Vanessodes clathrata Toulgoët, Paulianosia; Paulianosia clavata Walker, Glaucopis; Mallodeta clavipennis Butler, Pterygopterus; Timalus cleta Turner, Baeomorpha; Hobapromea clio Packard, Seirarctia; Euverna clite Walker, Josia; Getta clusia Druce, Laemacharis; Hypocharis clymene Brown, Phalaena; Haploa coavestis Hampson, Nudaria; Hemipsilia coccinea Schaus, Nyctosia; Nyctosia cocho Schaus, Sansaptera; Sansaptera coenosa Hübner, Bombyx; Laelia coeruleipuncta Gaede, Alandria; Alandria cognata Walker, Pseudosphex; Sphecosoma coleta Stoll, Phalaena; Coleta collaris Drury, Phalaena; Leucopsumis collaris Herrich-Schäffer, Paraplesis; Paraplesis collaris Herrich-Schäffer, Phaegoptera; Amastus collaris Jones, Castronia; Castronia collaris Möschler, Macronadata; Macronadata

collartorum Kiriakoff, Collartisa; Collartisa collateralis Hampson, Neoscaptia; Neoscaptia collenettei Kiriakoff, Asurgylla; Asurgylla collucens Lucas, Porthesia; Euzora colombina Fabricius, Zygaena; Eunomia colorata Walker, Darna; Crocomela columbia Dyar, Zellatilla; Zellatilla combusta Herrich-Schäffer, Anodonta; Arhacia combusta Walker, Anticyra; Dinara comita Schaus, Euharpyia; Euharpyia comma Strand, Stenostauridia; Stenostauridia comorana Jordan, Aegocera: Nesaegocera comparata Walker, Themaca; Themaca compsa Jordan, Pseudapiconoma; Compsochromia concentrica Oberthür, Fentonia; Homocentridia conchyfera Guenée, Nystalea; Nystalea concinna Smith, Phalaena; Oedemasia concinna Walker, Cottobara; Cottobara concocta Dognin, Odoxia; Odoxia concolor Walker, Nadiasa; Nadiasa conferta Walker, Mazaeras; Mazaeras conferta Walker, Pitane; Nepita confinis Herrich-Schäffer, Charidea; Nelphe confinis Walker, Glaucopis; Eressa confinis Walker, Phecada; Phecada conformis Hampson, Apothosia; Apothosia confusa Wileman, Stauropus; Taiwa confusum Roepke, Calyptronotum; Calyptronotum congerens Felder, Cymella; Myclela congrua Dyar, Congruia; Congruia conia Collenette, Coniortodes: Melgona conspersa Aurivillius, Lobilema; Lobilema conspersa Walker, Cincia; Cincia conspersa Walker, Teragra; Teragra conspersus Walker, Elysius; Elysius contenta Walker, Euchromia; Zygaenopsis contigua Walker, Charadra; Charadra contingata Möschler, Antiora; Pseudantiora contraria Walker, Paracles; Paracles contraria Walker, Teara; Teara contristis Hübner, Epicoma; Epicoma convexa Hering, Liparodonta; Liparodonta cora Hübner, Cerma; Cerma coracina Boisduval, Doracis; Doracis cordatula Druce, Odozana; Urozana coreana Matsumura, Coreodonta; Coreodonta cornea Herrich-Schäffer, Phaegoptera; Hemihyalea cornifrons Roepke, Allotoma; Allotoma cornuta Kiriakoff, Pygaeridia; Pygaeridia correbioides Felder, Acreagris; Acreagris correbioides Felder, Dycladia; Dycladia correcta Walker, Hypsa; Pachyphilona corsicum Rambur, Trichosoma; Somatrichia corticalis Butler, Hupodonta; Hupodonta corticina Aurivillius, Macrolaelia; Macrolaelia corydon Druce, Eupyra; Euchlorostola cosmeta Lower, Spilosoma; Cheliosea costalis Kiriakoff, Ulinodes; Ulinodes costalis Moore, Caragola; Caragolina costalis Walker, Abella; Clathe costalis Walker, Antipha; Antipha costalis Walker, Melia; Ilema costalis Walker, Moresa; Moresa costata Möschler, Pryteria; Pryteria costata Moore, Diduga; Diduga costigera Walker, Gargetta; Gargetta costimacula Mabille, Lithosia; Archithosia costivitta Walker, Thacona; Thacona cota Kiriakoff, Quista; Quista cotytto Kiriakoff, Ulinella; Ulinella courregesi Dognin, Purius; Metacrisia crassa Felder, Homoeocera; Homoeocera

crassa Felder, Zagaris; Zagaris crassicornis Walker, Chamaita; Caulocera craushayi Hampson, Micrilema; Micrilema credula Fabricius, Bombyx; Composia crenata Esper, Bombyx; Gluphisia crenulata Bethune-Baker, Rhodesana; Rhodesana crenulata Hampson, Lophopteryx; Ptilodontosia cresus Cramer, Phalaena; Chliara cretata Hampson, Euproctosia; Euproctosia cribraria Clerck, Phalaena; Argina cribraria Linnaeus, Phalaena; Coscinia cribrum Linnaeus, Phalaena; Coscinia crinada Dognin, Macroptila; Macroptila cristata Butler, Trabala; Rabtala critobulus Fawcett, Stauropus: Simesia crocea Schaus, Aptilosia; Aptilosia crocicollis Herrich-Schäffer, Liparis; Antiphella crocipes Boisduval, Cypra; Collenettema crocodora Meyrick, Licnoptera; Licnoptera croesus Hampson, Chrysozana; Chrysozana crossoea Druce, Heterocampa; Boriza crotalistria Bryk, Gylla; Gylla crucifera Perty, Chelonia; Dysschema cruenta Clemens, Cyanopepla; Cyanopepla cruenta Herrich-Schäffer, Gundlachia; Burtia cruenta Hübner, Pericopis; Phaloe cryptica Turner, Antimima; Antimima cryptocephala Bryk, Angustiala; Angustiala cryptoleuca Walker, Corydalla; Agyrta ctenuchoides Walker, Dotha; Dotha cubana Herrich-Schäffer, Pericopis; Sphaeromachia cucadma Druce, Cosmosoma; Leucopleura cucullatella Linnaeus, Phalaena; Nola cucullina [Denis & Schiffermüller], Bombyx; Ptilodontella cucullodes Walker, Cimbina; Cimbina cuculus Staudinger, Odontosia; Lophontosia culta Hübner, Sommeria; Sommeria cunea Drury, Phalaena; Hyphantria cunegunda Schaus, Jorgensenia; Geriojennsa cuneifera Dognin, Xenomigia; Ceraeotricha cuneiplena Walker, Darantasia; Darantasia cuneonotatus Walker, Ammatho; Ammatho cupra Dognin, Lafajana; Lafajana cuprea Schaus, Odozana; Metalobosia cupreimargo Hampson, Tropacme; Tropacme cuprina Moore, Apamea; Acanthopolia cuprizona Hampson, Padenodes; Padenodes curtula Linnaeus, Phalaena; Clostera curvatula Rothschild, Hoplitis; Scaeopteryx curvifera Walker, Pegella; Pegella curvilinea Swinhoe, Hyperaeschra; Acrasiella curvilinea Wileman, Norraca; Togaritensha curvinervis Felder, Eurylomia; Eurylomia cyanea Leech, Somera; Egonocia cyanogramma Hampson, Anaphosia; Anaphosia cyanomela Neumoegen, Daphne; Didaphne cyanomelas Felder, Phanoptis; Phanoptis cyanopyga Felder, Agape; Agape cyanosticta Kiriakoff, Tolmiana; Tolmiana cyclopea Kiriakoff, Archigargetta; Archigargetta cyclota Collenette, Dasychira; Volana cycnoptera Lower, Notodonta; Pheressaces cydista Turner, Teleclita; Teleclita cyma Hübner, Dioptis; Dioptis cymatophoroides Hampson, Dohertya; Dohertya cytheraea Druce, Lithosia; Neoplynes cytheris Druce, Phaeochlaena; Xenorma cyttarrosticta Hampson, Stenadonta; Stenadonta czekanowskii Grum-Grshimailo, Hyperborea; Hyperborea

daedala Collenette, Soreutoneura; Soreutoneura daisetsuzana Matsumura, Konokareha; Konokareha dalmeidai d'Andretta &c Travassos, Romualdisca; Romualdisca dalsa Schaus, Gartha; Gartha dama Druce, Eupyra; Cyanarctia danbyi Neumoegen, Melia: Eumelia danieli Kiriakoff, Rhipidarctia; Hemirhipidia daphaena Hampson, Pseudapiconoma; Daphaenisca dara Druce, Stilpnotia; Eustema dardania Druce, Tifama; Drastoma dasychirina Gaede, Hoplitis; Afroplitis dasychiroides Butler, Stauropus; Simesia dasychiroides Janse, Breyeria; Eubreyeria dasychiroides Strand, Pseudodura; Pseudodura dasypyga Felder, Diastrophia; Diastrophia davidsoni Schaus, Ginaldia; Ginaldia debilis Felder, Macronyx; Teracotona debilis Gaede, Xanthodonta; Xanthodonta decaryi Griveaud, Fletcherinia; Fletcherinia decempunctata Tessman, Pirgula; Pirgula deceptans Dyar, Parillice; Parillice decia Boisduval, Lithosia; Leptarctia decipiens Kiriakoff, Codita; Codita decipiens Kiriakoff, Hepatophalera; Hepatophalera decora Linnaeus, Phalaena; Pais decora Walker, Cissura; Cissura decora Walker, Sacada; Sacada decorata Kiriakoff, Antistaura; Antistaura decorata Walker, Anaxita; Anaxita decreta Butler, Lithosia; Aedoea decurrens Dyar, Postanita; Postanita defecta Walker, Barsine; Barsine defoliata Walker, Alpis; Alpis dehanna Pagenstecher, Arctia; Niasana deldon Druce, Polypoetes; Polypoetes delecta Butler, Hysia; Hypatia deletus Rebel, Arctiites; Arctiites delia Leech, Drymonia; Neodrymonia delicatula Boisduval, Cypra; Cypra delineata Walker, Rolepa; Rolepa delineata Walker, Vunga; Vunga demas Kiriakoff, Afrogluphisia; Afrogluphisia dentata Hampson, Hyperaeschra; Polyaeschra dentata Kiriakoff, Antichadisra; Antichadisra dentata Schaus, Symmerista; Tagela dentata Walker, Cetola; Cetola dentatus Gaede, Melagona; Melagona denticulata Schaus, Psorocampa; Psorocampa denudata Walker, Sitvia; Sitvia deornata Walker, Salamboria; Salamboria depressa Esper, Noctua; Katha derelicta Prout, Crambometra; Crambometra deserta Felder, Pyralidia; Pyralidia desiccata Kiriakoff, Xeropteryx; Spatalina detecta Walker, Bazisa; Baziza; detrita Esper, Phalaena; Parocneria detrita Walker, Cyana; Cyana devestita Walker, Odagra; Odagra deyrolii Walker, Glaucopis; Erruca dharma Moore, Gangarides; Gangarides dialampa Staudinger, Epimydia; Epimydia diana Druce, Phaenareta; Phaenarete diaphana Eversmann, Lithosia; Hyalocoa diaphana Kollar, Syntomis; Caeneressa diaphanella Hampson, Palaeopsis; Palaeopsis dichroa Hübner, Hypocrita; Hypocrita dictaea Linnaeus, Phalaena; Pheosia didugana Draudt, Dasippia; Dasippia diehli Kiriakoff, Pantanopsis; Pantanopsis diffusum Felder, Orthosoma; Rhuda dilatata Walker, Josia; Scedros

dilucida Warren, Tanaostyla; Tanaostyla diluta Felder, Cratoplastis; Cratoplastis dimidiata Walker, Thirmida; Thirmida diminuta Walker, Dioptis; Micragyrta dinawa Bethune-Baker, Pseudilema; Pseudilema dioptis Felder, Chamesthema; Chamesthema diores Cramer, Phalaena; Phiditia diplaga Hampson, Euproctis; Medama diplisticta Bethune-Baker, Eilema; Nanna dipotamica Tams, Sumeria; Sumeria diptera Walker, Euchromia; Melisa dirphioides Walker, Turuenna; Turuenna discalis Brandt, Sidemia; Brandtaxia discalis Moore, Challa; Challa discalis Walker, Aroa; Aroa discalis Walker, Baritius; Baritius discalis Walker, Eloria; Axuenna discalis Walker, Halesidota; Azatrephes discalis Walker, Lithosia; Stenaulis discalis Walker, Lusura; Lusura discata Schaus, Cyclosticta; Cyclosticta discivitta Moore, Heracula; Heracula discocellularis Kiriakoff, Atornoptera; Atornoptera discocellularis Strand, Norracoides; Norracoides discoidalis Gaede, Pydna; Oglana discoidalis Matsumura, Hyperaeschra; Allodontoides disjuncta Dognin, Nerice; Pronerice disjuncta Walker, Brycea; Brycea dispar Kiriakoff, Loudonta; Loudonta dispar Linnaeus, Phalaena; Porthetria dispar Rothschild, Aegocera; Epischausia dispar Walker, Pantana; Pantana disrupta Moore, Celeia; Metaschalis dissimilis Hampson, Oxacme; Oxacme dissodectes Kiriakoff, Cerura; Oreocerura dissozona Meyrick, Hectogama; Hectogama distigmata Hampson, Ilema; Cragia distincta Rothschild, Tessellarctia; Pydnaodes distinguenda Walker, Aethalida; Aethalida distorta Moore, Pangora; Pangora distorta Moore, Zadadra; Zadadra ditissimus Walker, Scaptius; Scaptius divaricata Gaede, Phalera; Snellenita divergens Kiriakoff, Desmeocraera; Liriochroa diversa Bethune-Baker, Pseudogargetta; Pseudogargetta diversa Walker, Avela; Avela diversipes Walker, Biturix; Biturix dives Oberthür, Spatalia; Spataloides divisa Boisduval, Pyralopsis; Pyralopsis divisa Gaede, Parastaura; Parastaura divisa Moore, Hesudra; Hesudra divisa Walker, Apela; Apela divisa Walker, Arctia; Catarctia divisa Walker, Eloria; Marbla divisa Walker, Lamnis; Lamnis divisus Herrich-Schäffer, Creatonotos; Echeta dochmoschema Turner, Porphyrochrysa; Porphyrochrysa docis Hübner, Automolis; Himerarctia dodonaea [Denis & Schiffermüller], Bombyx; Drymonia dohertyi Rothschild, Pseudoceryx; Pseudoceryx dolens Druce, Ctenucha; Stenucha dominia Cramer, Phalaena; Neochera dominula Linnaeus, Phalaena; Callimorpha dora Neumoegen & Dyar, Doa; Doa doriae Oberthür, Pseudocallimorpha; Pseudocallimorpha dorsalis Kiriakoff, Closteroides; Closterellus dorsalis Walker, Ichthyura; Gaugamela dorsalis Walker, Rosema; Rosema dorsipuncta Hampson, Neonerita; Neonerita dorsisigna Walker, Balbura; Balbura dorsoflavida Kiriakoff, Epicosmolopha; Epicosmolopha dotata Walker, Byrsia; Byrsia

doubledayi Newman, Castulo; Cluaca dracoena Kiriakoff, Metarctia; Collocaliodes dromedarius Linnaeus, Phalaena; Notodonta drostana Schaus, Vulmara; Vulmara druna Swinhoe, Xylecata; Xylecata druryi Hübner, Trichura; Trichura dryas Cramer, Phalaena; Sychesia dryinopa Dodd, Notodonta; Teleclita drymonides Strand, Zerafia; Zerafia dubia Warren, Cyphopora; Cyphopora dubiosa Hering, Dasychoproctis; Dasychoproctis dubiosa Schaus, Sauritinia; Sauritinia dudgeoni Elwes, Nudaria; Ovipennis dufranei Kiriakoff, Apisa; Lempkeella dulcicula Swinhoe, Hemonia; Cyclosiella duplex Gaede, Onophalera; Onophalera duplicata Walker, Gallaba; Gallaba durvillii Boisduval, Cocytia; Cocytia duvaucelii Boisduval, Psichotoe; Psichotoe

eagrus Cramer, Sphinx; Dinia ebalea Stoll, Phalaena; Nystalea eborella Boisduval, Lithosia; Lamprosiella eborina [Denis & Schiffermüller], Noctua; Cybosia eburneana Walker, Tospitis; Garudinistis echo Smith, Phalaena; Seirarctia ecnomoda Swinhoe, Leucoma; Carriola ectophaea Hampson, Asura; Habrochroma ectozona Hampson, Neoduma; Neoduma ectrocta Hampson, Aethosia; Aethosia editha Walker, Caralisa; Caralisa edwardsi Collenette, Orithrepta; Orithrepta effecta Schaus, Dottia; Dottia efulensis Holland, Syntomis; Thyrogonia egaca Walker, Bepara; Bepara egle Drury, Phalaena; Euchaetes eingana Schaus, Fentonia; Toddiana eion Druce, Pseudocharidea; Pseudocharidea electraula Meyrick, Amalodeta; Amalodeta elegans Boisduval, Damias; Damias elegans Janse, Leucophalera; Leucophalera elegans Kiriakoff, Schedostauropus; Schedostauropus elegans Moore, Grotea; Sebastia elegantissima Herrich-Schäffer, Phaegoptera; Echemis eleuthera Stoll, Phalaena; Pseudeucereon eliera Schaus, Dugonia; Dugonia elissa Schaus, Automolis; Selenarctia elongata Hering, Scoturopsis; Euscoturopsis elongata Matsumura, Ginshachia; Ginshachia elongata Rothschild, Phalera; Teinophalera elongata Schaus, Malupa; Malupa empusaria Guenée, Leucula; Leucula endophaea Dognin, Paragylla; Paragylla enervis Schaus, Idalus; Hypidalia entella Cramer, Phalaena; Oeonistis eone Hübner, Agerocha; Agerocha epelytes Collenette, Lymantica; Lymantica ephippiata Wallengren, Sirenopyga; Sirenopyga epigena Stoll, Phalaena; Rhogalia epinephela Bethune-Baker, Metopolophota; Metopolophota ericae Germar, Orgyia; Thylacigyna erichrysa Meyrick, Metacrias; Metacrias eriopis Herrich-Schäffer, Agarista; Pristoceraea eriphia Fabricius, Zygaena; Antichloris erlangeri Rothschild, Metarctia; Oenarctia erminea Esper, Phalaena; Apocerura eroki Bethune-Baker, Epidonta; Epidonta erosa Herrich-Schäffer, Halysidota; Psychophasma erratica Schaus, Mantruda; Mantruda

erratipennis Walker, Maschane; Maschane erubescens Butler, Cosmosoma; Lepidoneiva erubescens Hampson, Tricholepis; Chiretolpis erycinoides Felder, Dioptis; Isostyla erycinoides Felder, Marsypophora; Marsypophora erythrarchos Walker, Glaucopis; Rezia erythrias Meyrick, Haplopseustis; Acnissa erythrina Holland, Lomadonta; Lomadonta erythrophora Felder, Ptychoglene; Ptychoglene erythropygum Felder, Camptoloma; Camptoloma erythrotelus Walker, Glaucopis; Hyaleucerea erythrozonata Hampson, Lasiosceles; Scelilasia eryx Fabricius, Zygaena; Belemnia eschara Dyar, Epicrisias; Epicrisias esmeralda Hampson, Cerura; Hampsonita esperanza Schaus, Talara; Abrochocis esquamata Bethune-Baker, Paraxena; Paraxena etiennei Viette, Matarum; Matarum eucalypti Doubleday, Hylaeora; Hylaeora euchromoides Walker, Coreura; Coreura euchrysa Meyrick, Heterallactis; Heterallactis eucyaniformis Dognin, Seileria; Seileria eugenia Staudinger, Pteroma; Pterotes eugoana van Eecke, Malesia; Malesia eugraphica Turner, Trissobrocha; Trissobrocha eugraphica Walker, Spilosoma; Leucaloa eumenes Herrich-Schäffer, Eumenogaster; Eumenogaster eumenides Newman, Myrmecopsis; Myrmecopsis eumeta Druce, Lophopteryx; Herbertina eupatagia Hampson, Pydna; Innisca euprepiaeformis Herrich-Schäffer, Arctiomorpha;

Arctiomorpha eurocilia Cramer, Phalaena; Pericopis euryzona Lower, Euproctis; Habrophylla evadnes Stoll, Sphinx; Chrostosoma evergista Stoll, Phalaena; Deilemera evidens Hübner, Noctua; Sideridis evippe Walker, Josia; Erbessa evippus Druce, Idalus; Hypomolis excellens Strand, Pheositis; Pheositis excellens Strand, Shisa; Shisa excisa Hampson, Adoxosia; Adoxosia excisa Hampson, Tortricosia; Tortricosia excisa Rothschild, Garudinia; Macadumosia excurvata Hampson, Pheosia; Poecilopheosia exigua Nietner, Trichia; Trichia exilicosta de Joannis, Hanoisiella; Hanoisiella exposita Butler, Stenelopsis; Stenelopsis extensa Walker, Lemyra; Lemyra extravagans Warren, Episcea; Episcea extrema Walker, Contrebia; Contrebia exusta Butler, Gelastocera; Gelastocera

fadella Mabille, Hylemera; Microhyle
fagi Linnaeus, Phalaena; Stauropus
falsimonia Schaus, Dolichesia; Dolichesia
fascelina Linnaeus, Phalaena; Dicallomera
fasciata Bethune-Baker, Hirsutopalpis; Hirsutopalpis
fasciata Moore, Gampola; Gampola
fasciata Moore, Laelioides; Laelioides
fasciata Moore, Pheosia; Neopheosia
fasciata Moore, Pydna; Honveda
fasciata Schaus, Zonoda; Zonoda
fasciata Walker, Curoba; Curoba
fasciata Walker, Dulichia; Knappetra
fasciculata Walker, Agylla; Agylla
fasciculosa Felder, Polyptychia; Polyptychia
fasciolata Rothschild, Caulocera; Schalodeta
fasciolatum Butler, Sphecosoma; Sphecosoma

fassli Draudt, Horamella; Horamella fassli Prout, Tolimicola; Tolimicola fastosa Turner, Heterotropa; Heterotropa fenestra Linnaeus, Phalaena; Hyalurga fenestra Walker, Dioptis; Nepe fenestrata Walker, Coborisa; Coborisa fenestrata Walker, Gyara; Gyara fernandezi Watson, Emurena; Emurena ferox Walker, Euchromia; Hippola ferraria Walker, Acherdoa; Acherdoa ferrifera Walker, Torona; Torona ferruginea Aurivillius, Brachychira; Brachychira ferruginea Butler, Bettonia; Bettonia ferruginea Packard, Lophodonta; Lophodonta ferruginea Walker, Pithea; Pithea ferruginosa Walker, Euchromia; Dipaenae fervens Walker, Pitane; Pitane fervidus Walker, Ammalo; Ammalo festiva Hufnagel, Phalaena; Eucharia ficus Fabricius, Noctua; Lacides fifiana Dognin, Trumanda; Trumanda figulina Butler, Gangamela; Scearctia fimbria Möschler, Sychesia; Sychesia fissura Saalmüller, Acroctena; Acroctena flabrifera Moore, Nishada; Nishada flammans Walker, Euchromia; Phauda flammeola Distant, Heteronygmia; Hemerophanes flammula Geyer, Calosoma; Osmacola flava Hampson, Hemonia; Trischalis flava Hering, Parabitecta; Parabitecta flava Wallengren, Saenura; Saenura flavens Mabille, Coenostegia; Coenostegia flaveola Rothschild, Bitecta; Eurynora flavescens Kiriakoff, Rheneades; Rheneades flavibasis Hering, Anexotamos; Anexotamos flavicapilla Hübner, Atolmis; Scotura flavicapilla Wallengren, Leucoma; Homalomeria flaviceps Druce, Cyptonychia; Cyptonychia flaviceps Walker, Eloria; Xenosoma flavicollis Druce, Atolmis; Gnamptonychia flavicollis Leech, Crinola; Halseyella flavicollis Moore, Mahavira; Mahavira flavicosta Wallengren, Lithosia; Pusiola flavicostata Hampson, Cyclosodes; Cyclosodes flavicostata Snellen, Pitane; Diduga flavida Hampson, Pydna; Antheuella flavilimbata Herrich-Schäffer, Olenoptera; Olenoptera flavimarginata Hampson, Aglossosia; Aglossosia flavimargo Hering, Laelaroa; Laelaroa flavinotata Butler, Aclonophlebia; Aclonophlebia flavipalpata Staudinger, Ocneria; Casama flavipes Hampson, Cispia; Aruta flaviplaga Heylaerts, Gnophrioides; Gnophrioides flavipunctata Bethune-Baker, Eressades; Eressades flavissima Walker, Lyces; Erilyces flavivena Hampson, Metarctia; Notharctia flavogrisea Leech, Hypeugoa; Hypeugoa flavopunctata Bethune-Baker, Halone; Chrysasura flexilisana Walker, Conchylis; Stictosia flexuosa Walker, Certila; Certila flocciferus Möschler, Strophocerus; Strophocerus floccosa Walker, Odozana; Odozana focula Stoll, Phalaena; Rhuda fontainei Collenette, Sankurua; Kunusara forbesi Druce, Cypra; Rajacoa forficula Draudt, Proctocopis; Proctocopis formosae Strand, Cleapa; Marushachia formosae Strand, Creatonotus; Amsactoides formosana Matsumura, Shachihoka; Shachihoka formosicola Strand, Pydna; Achepydna formula Grote, Robinsonia; Robinsonia

fractilinea Snellen, Pitane; Stictane

fractivittarum Dyar, Nudur; Nudur franki Kiriakoff, Galos; Galos friedae Kiriakoff, Blacodes; Neoblacodes friga Druce, Thyrarctia; Gonotrephes frigida Walker, Devara; Euchontha frugalis Leech, Pydna; Curuzza fucosa Hübner, Hypoprepia; Hypoprepia fulgurifera Walker, Crino; Megashachia fulgurifera Walker, Gozarta; Bardaxima fuliginea Butler, Pyramocera; Pyramocera fuliginosa Linnaeus, Phalaena; Phragmatobia fuliginosa Saalmüller, Calliteara; Abakabaka fulleri Druce, Nyctemera; Eohemera fulva Stretch, Kodiosoma; Kodiosoma fulva Walker, Lagela; Lagela fulvescens Walker, Syntomis; Ceryx fulvia Linnaeus, Phalaena; Clerckia fulviceps Felder, Cymopsis; Cymopsis fulviceps Walker, Icta; Icta fulvicollis Hübner, Glaucopis; Cisseps fulvicolora Hampson, Leptaroa; Leptaroa fulvilaniata Dognin, Trichareva; Trichareva fulvinigra Hampson, Eurozonosia; Eurozonosia fulvinota Hampson, Opsaroa; Opsaroa fulvipicta Hampson, Nudaria; Gymnochroma fulvipuncta Hampson, Machairophora; Machairophora fulvohirta Walker, Alphaea; Alphaea fulvohirta Walker, Ardices; Ardices fumeola Walker, Lithosia; Macrosia fumipennis Walker, Euchromia; Ceramidia fumosa Holloway, Thumathoides: Thumathoides fumosa Matsumura, Dudusa; Dudusopsis fumosa Matsumura, Epinotodonta; Epinotodonta furcata Möschler, Tricypha; Tricypha furcifasciata Butler, Gnophria; Carcinopodia furcifera Walker, Setina; Phaeophlebosia furcilla Packard, Platycerura; Platycerura furcula Clerck, Phalaena; Furcula furva Leech, Ocneria; Maimaia fusca Hampson, Microtane; Microtane fusca Walker, Acyphas; Acyphas fusca Walker, Massicyta; Massicyta fuscapex Hampson, Utriculofera; Utriculofera fuscata Talbot, Eurystaura; Diceratostaura fuscescens Walker, Thumatha; Thumatha fuscicollis Gaede, Pseudogargetta; Roepkeella fuscifusa Hampson, Nudosia; Nudosia fuscimarginalis Swinhoe, Zygaenopsis; Zygaenosia fuscipennana Walker, Tospitis; Omiosia fuscivena Swinhoe, Stracena; Stracena fusiformis Walker, Nioda; Nioda

gabonensis Kiriakoff, Antitrotonotus; Antitrotonotus gabunica Holland, Euproctidion; Euproctidion gallans Karsch, Nunua; Nunua gangis Linnaeus, Phalaena; Creatonotos garleppi Druce, Zatrephes; Gorgonidia gavisa Walker, Agyrta; Micragyrta gelida Grote, Ellida; Ellida gelida Walker, Caprima; Caprima gelida Walker, Doliche; Doliche gelidata Walker, Damna; Ticilia geminata Herrich-Schäffer, Mystrocneme; Mystrocneme gentilis Felder, Sthenognatha; Sthenognatha geographica Meyrick, Choromeles; Choromeles geometrica Grote, Eupsychoma; Eupsychoma gerardi Holloway, Fabresema; Fabresema germana Rothschild, Odozana; Neozana gibbosa Smith, Phalaena; Nadata

gigantea Druce, Josia; Josiomorphoides gigantea Elwes, Notodonta; Oreodonta gigantea Röber, Callindra; Callindra gigas Distant, Hoplitis; Melebaeas gigas Walker, Lithosia; Macrobrochis gilveola Ochsenheimer, Lithosia; Manulea gladstonei Janse, Campyloctys; Campyloctys glandulosa Felder, Choerotricha; Choerotricha glauca Turner, Parameces; Parameces glauca Walker, Osica; Osica glaucaspis Walker, Dioptis; Phelloe glaucoalba Walker, Phreata: Phreata glaucopis Felder, Telioneura; Telioneura glaucoviridis Rothschild, Stauropus; Kakostauropus gnatula Boisduval, Naclia; Cacosoma goodii Holland, Oecura; Oecura gortynoides Grote, Carathis; Carathis gortynoides Walker, Bellura; Bellura graminosa Walker, Sizalisca; Sizalisca grammica Linnaeus, Phalaena; Spiris grammodes Hering, Dasychira; Orana grammophora Felder, Autoceras; Autoceras grandis Butler, Numenoides; Numenoides grandis Druce, Automolis; Xanthomis grata Fabricius, Bombyx; Cyphocampa grisea Bethune-Baker, Parakanchia; Parakanchia grisea Butler, Microphalera; Microphalera grisea Gaede, Dasyarctia; Dasyarctia grisea Kiriakoff, Epicerurina; Epicerurina grisea Schaus, Nezula; Nezula griseobasis Rothschild, Chamaita; Teinomastyx griseodivisa Bryk, Pydna; Blennema griseola Hübner, Bombyx; Collita griseola Kiriakoff, Stauropida; Stauropida griseoviridis Laporte, Nocthadena; Nocthadena grisescens Bethune-Baker, Parapelosia; Parapelosia grisescens Daniel, Pareugoa; Apogurea grisescens Dufrane, Metarctia; Dufraneella grisescens Kiriakoff, Overlaetia; Tricholoba grisescens Roepke, Stauropodopsis; Stauropodopsis grisescens Staudinger, Cnethodonta; Cnethodonta griseus Hampson, Stauropus; Poliostauropus griveaudi Kiriakoff, Italaviana; Italaviana grotei Packard, Anatolmis; Anatolmis grotesca Bryk, Pontalamia; Pontalamia guacolda Poey, Erithales; Erithales guatemalteca Felder, Mesenochroa; Mesenochroa guineensis Fabricius, Zygaena; Euchromia gundlachiana Neumoegen, Ira; Cubira guttata Walker, Artaxa; Artaxa guttipalpis Walker, Leucopsumis; Glaucostola guttivaga Walker, Soloe; Soloella guttivena Walker, Claphe; Claphe guttivitta Walker, Agrisius; Agrisius guttivitta Walker, Cecrita; Cecrita guttulosa Walker, Syntomis; Agaphthora gyge Druce, Hiera; Hiera gymna Collenette, Amphekes; Amphekes gyon Fabricius, Zygaena; Phaeochlaena

habanera Schaus, Soritena; Soritena haematica Holland, Metarctia; Thyretarctia haematobasis Hampson, Phoeniostacta; Phoeniostacta haematochrysia Kiriakoff, Bergeria; Bergeria haematoessa Holland, Metarctia; Mecistorhabdia haemon Druce, Tithraustes; Tithraustes haemorrhoidalis Stoll, Sphinx; Aethria hagnoleuca Dyar, Mannina; Mannina haigi Collenette, Thambeta; Thambeta

halmaphylla Dyar, Calomathetes; Calomathetes hammamelis Mell, Stigmatophorina; Stigmatophorina hampsoni Dognin, Fasslia; Fasslia hampsoni Swinhoe, Migoplastis; Paraplastis hampsonia van Eecke, Katmeteugoa; Katmeteugoa hanga Herrich-Schäffer, Laemocharis; Aristodaema haplodes Meyrick, Teratopora; Teratopora hapygia Felder, Lobogona; Lobogona harpalyce Schaus, Cosmosoma; Mirandisca harpotoma Turner, Mesodrepta; Mesodrepta harrisii Boisduval, Ctenucha; Pygoctenucha hazara Butler, Pericopis; Neolaurona hearseyana Moore, Digama; Digama hebe Linnaeus, Phalaena; Eucharia heber Cramer, Sphinx; Aclytia hebes Walker, Delphyre; Delphyre heliconia Linnaeus, Phalaena; Hypsa helops Cramer, Phalaena; Ammalo helvola Hübner, Bombyx; Katha hemicalla Lower, Orgyia; Oligeria hemiluta Collenette, Eala; Eala heminephes Meyrick, Tigrioides; hera Linnaeus, Phalaena; Euplagia herbida Walker, Luca; Luca herbuloti Toulgoët, Philenora; Novosia herona Druce, Anace; Heronina hesperia Cramer, Phalaena; Otroeda heterochroa Mabille, Deiopeia; Callicereon heterogena Oberthür, Ecpantheria; Catenina heterogyna Hampson, Ilemodes; Ilemodes heteroides Collenette, Lymantralex; Lymantralex hexamitobalia Collenette, Dasychira; Lanitra hierax Distant, Desmeocraera; Pararethona hilaris Walker, Doliche; Adites hipparchus Cramer, Sphinx; Pompostola hippia Stoll, Phalaena; Evius hippona Druce, Odozana; Prepiella hippotes Cramer, Sphinx; Thyretes hirsuta Walker, Euchromia; Enope hirta Walker, Lithosia; Calamidia histrionica Herrich-Schäffer, Notodonta; Minara histrionica Herrich-Schäffer, Phaegoptera; Phaegoptera hoemorrhoa Walker, Uraga; Uraga hoenei Daniel, Hypasura; Hypasura hoenei Daniel, Lithosarctia; Lithosarctia hoenei Daniel, Paralithosia; Paralithosia hoenei Gaede, Melagonina; Melagonina hoenei Kiriakoff, Paranerice; Paranerice holdingii Felder, Cataphractes; Cataphractes holoclera Karsch, Caryatis; Karschiola homopteridea Butler, Cercopimorpha; Cercopimorpha horae Druce, Sagaris; Sagaropsis horsfieldi Moore, Eumeta; Brykia huebneri Boisduval, Syntomis; Buthysia humeralis Walker, Ityca; Ityca humilis Herrich-Schäffer, Chrysostola; Chrysostola humilis Walker, Cyllene; Lyclene hunyada Swinhoe, Pydna; Hunyada hyalina Janse, Lepidopalpus; Lepidopalpus hyalozona Felder, Sphecopsis; Sphecopsis hydromeli Harvey, Litodonta; Litodonta hyhochloe Kiriakoff, Sidisca; Sidisca hypoleucus Herrich-Schäffer, Ecdemus; Ecdemus hypophaeola Hampson, Neasura; Neasura hypoxantha Hübner, Dysschema; Dysschema hypoxantha Wallengren, Ornithopsyche; Ornithopsyche hypparchus Cramer, Sphinx; Marmax

icasia Cramer, Phalaena; Hypercompe ichneumonea Felder, Rhynchopyga; Rhynchopyga icilia Stoll, Phalaena; Nygmia idiopis Collenette, Sychnacedes; Sychnacedes ignea Druce, Mesothen; Mesothen ignepicta Hampson, Stenaroa; Stenaroa igninix Walker, Mieza; Mieza ignita Walker, Euchromia; Chrysocale illuminata Maassen, Pais; Protracta ima Boisduval, Tipulodes; Tipulodes imaon Cramer, Sphinx; Syntomoides imbuta Walker, Arctia; Nayaca imbutana Walker, Deua; Deua imitans Butler, Zelomera; Zelomera imitans Draudt, Loxomima; Loxomima immaculata Grote, Phryne; Calybia imparepunctata Oberthür, Lithosia; Calpoparia impeditus Walker, Cossus; Stenostaura imperialis Herrich-Schäffer, Eupyra; Eupyra imperialis Heylaerts, Romanoffia; Romanoffia imperialis Kollar, Euprepia; Melanareas imperialis Walker, Euchromia; Epanycles impervia Walker, Lithosia; Petalopleura implens Walker, Manoba; Manoba impolita Schaus, Anaulosia; Anaulosia impomaeae Doubleday, Schizura; Schizura impunctibasis Kiriakoff, Mismia; Mismia inachia Schaus, Charidea; Callopepla incarnata Freyer, Heliothis; Chazaria incarnata Walker, Lerina; Lerina incendiaria Hübner, Glaucopis; Isanthrene incertus Walker, Malabus; Malabus inclusa Walker, Addua; Addua incompleta Gaede, Antheusina; Antheusina incomptaria Walker, Utidava; Utidava inconcisa Walker, Sciatta; Sciatta inconspicua Draudt, Paramevania; Paramevania inconspicua Kiriakoff, Parafentonia; Hemifentonia inconspicuana Strand, Desmeocraerula; Desmeocraerula inconstans Butler, Dictenus; Dictenus incurvata Kiriakoff, Leptonadatoides; Leptonadatoides indeterminata Walker, Casama; Casama indiana Grote, Nystalea; Eunystalea indica Moore, Dysauxes; Gurna indica Moore, Pydna; Rebita indica Walker, Gluphisia; Varmina indistincta Gaede, Pydna; Afropydna indistincta Hampson, Costarcha; Costarcha inexpectata Kiriakoff, Rhynchophalerina; Rhynchophalerina inexpectata Rothschild, Phalera; Aprosdocetos inexpectata Toulgoët, Pseudogaltara; Pseudogaltara infans Kiriakoff, Helga; Helga infausta Hering, Pseudostracilla; Pseudostracilla infernalis Butler, Thanatarctia; Thanatarctia infumata Felder, Setinochroa; Setinochroa infumata Hampson, Diplonyx; Didymonyx infusca Matsumura, Horishachia; Horishachia infuscata Matsumura, Kikuchiana; Kikuchiana ingens Karsch, Amyops; Amyops ingens Walker, Porsica; Porsica innocens Hering, Croremopsis; Croremopsis innotata Walker, Spilosoma; Casama inops Walker, Hypsa; Philona inordinata Walker, Sybrida; Sybrida inornata Walker, Amatissa; Amatissa inscripta Walker, Marcipa; Marcipa inscripta Walker, Sesapa; Sesapa insignior Kiriakoff, Neoclostera; Neoclostera insignis Aurivillius, Euproctilla; Euproctilla instructalis Walker, Mamala; Mamala insueta Kiriakoff, Diodorina; Diodorina

insularis Kiriakoff, Romaleostaura; Romaleostaura

insulata Walker, Halysidota; Pareuchaetes intacta Walker, Eloria; Eloria integra Felder, Clastognatha; Clastognatha integra Walker, Josia; Phalcidon integra Walker, Mennis; Antiotricha intercissa Freyer, Eyprepia; Lacydes interfixa Walker, Phaos; Phaos interiorata Walker, Numenes; Camptoloma interlecta Walker, Nyctemera; Nyctemera intermixta Walker, Felia; Felia intermixta Warren, Phytometra; Thysanoplusia interpellatrix Wallengren, Stauropus; Desmeocraera interrupta Linnaeus, Phalaena; Creatonotos intersecta Hampson, Paramya; Metamya interstitialis Herrich-Schäffer, Macrobrochis; Macrobrochis intervenata Schaus, Pernambis; Pernambis intricata Walker, Euchromia; Echoneura intricata Walker, Isia; Isia invaria Walker, Anace; Elsita invaria Walker, Repena; Thelde invasa Walker, Turriga; Turriga inversa Schaus, Paratalara; Paratalara involuta Sepp, Phalaena; Eupseudosoma io Viette, Fentonina; Acrasiodes iocephala Turner, Maurilia; Austrocarea iodea Herrich-Schäffer, Phaegoptera; Pachydota ipomoeae Harris, Glaucopis; Syntomeida iridescens Kiriakoff, Iridoplitis; Iridoplitis iridescens Schaus, Munona; Munona irregularis Toulgoët, Alepista; Alepista irrorata Butler, Olceclostera; Olceclostera irrorata Hampson, Pasteosia; Pasteosia irrorata Moore, Daplasa; Daplasa irrorata Rothschild, Calamidia; Striosia irrorata Rothschild, Diacrisia; Chlorocrisia irrorella Linnaeus, Phalaena; Setina isabella Smith, Phalaena; Pyrrharctia isabella White, Trichetra; Sthenadelpha isophragma Meyrick, Comarchis; Comarchis isthmia Felder, Euagra; Mydromera ivyalba Van Eecke, Sicciaemorpha; Sicciaemorpha

jacobaeae Linnaeus, Phalaena; Tyria jaliscana Schaus, Notela; Notela jamaicensis Schaus, Prumala; Prumala jansei Kiriakoff, Iostaura; Iostaura jansei Kiriakoff, Metascrancia; Metascrancia japonica Marumo, Yazawaia; Yazawaia japonica Walker, Atolmis; Epatolmis japonica Wileman & South, Tarsolepis; Tarsolepisoides jason Dognin, Lithoprocris; Lithoprocris javana Cramer, Phalaena; Asota javanica Roepke, Darantasiella; Darantasiella javensis Kiriakoff, Chadisrella; Chadisrella jessica Butler, Plusia; Sclerogenia jezoensis Matsumura, Ptilophoroides; Ptilophoroides jezoensis Wileman & South, Spatalia; Stenospatalia joiceyi Talbot, Heptaptosis; Heptaptosis jonasii Butler, Aroa; Topomesoides jordani Hering, Pirgulina; Pirgulina jucunda Walker, Pallene; Heliosia judas Hübner, Apistosia; Apistosia jugifera Felder, Micrommia; Micrommia julumito Dognin, Sphecopsyche; Sphecopsyche junctura Walker, Melanochroia; Hyperphara justina Stoll, Phalaena; Ernassa juturna Cramer, Phalaena; Antaea juvenis Walker, Ptilomacra; Hondella

kalisi Roepke, Macrohoplitis; Macrohoplitis kanshireiensis Wileman & South, Neeugoa; Neeugoa kawachiensis Matsumura, Nagandopsis; Nagandopsis kefersteinii Butler, Hydrusa; Tritonaclia kitchingi Bethune-Baker, Laelia; Sphragista kivensis Dufrane, Apisa; Diakonoffia korbi Staudinger, Bombyx; Albarracina kosemponica Strand, Hyperaeschra; Hyperaeschrella

l-nigrum Müller, Phalaena; Arctornis laboulbeni Bar, Palustra; Palustra labuana Swinhoe, Ctenane; Ctenane lacrumans Schaus, Agunaix; Agunaix lactea Bethune-Baker, Pellucens; Pellucens lactea Talbot, Monarctia; Monarctia lacteata Holland, Nudaria; Lymantriopsis lacteolaria Hulst, Leucula; Leuculodes lacticinia Cramer, Phalaena; Nyctemera lactinea Cramer, Phalaena; Aloa ladakensis Moore, Lachana; Lachana ladislai Oberthür, Lophopteryx; Fusapteryx laennus Walker, Glaucopis; Gymnelia laeta Walker, Termessa; Termessa laeticolor Felder, Paratype; Paratype laevis Butler, Fentonia; Fentonia lajonquierei Viette, Mafana; Mafana lamprolepis Hampson, Megaceramis; Megaceramis lanaria Holland, Terphothrix; Terphothrix lanassa Druce, Edema; Poresta lanceolata Roepke, Stenopydna; Stenopydna lanceolata Walker, Collyta; Collyta lanceolata Walker, Josia; Loxozona lanceolata Walker, Rilia; Rilia lanceolatus Butler, Chloropsinus; Chloropsinus lanifera Walker, Eragisa; Eragisa lapidaria Walker, Mulona; Mulona lapponica Thunberg, Bombyx; Pararctia lasea Druce, Xenosoma; Somatoxena lassa Schaus, Anita; Proanita lata Holland, Aroa; Usimbara latana Walker, Tospitis; Garudinia lateralis Walker, Motada; Motada lateritia Herrich-Schäffer, Automolis; Hebena laticilia Walker, Areva; Areva laticostalis Hampson, Spatalia; Pseudallata latifascia Walker, Cleapa; Cleapa latifascia Walker, Eucereon; Acridopsis latilinea Walker, Glaucopis; Leucotmemis latipennis Butler, Gonoclostera; Gonoclostera latipennis Butler, Phalera; Leucophalera latipennis Strand, Alatanadata; Alatanadata lativitta Moore, Rajendra; Rajendra lativitta Walker, Armiana; Armiana latreillana Kirby, Ctenucha; Ctenucha latreillii Godart, Chelonia; Artimelia latus Grote, Cytorus; Cytorus latus Rothschild & Jordan, Eriomastyx; Eriomastyx laudabilis Druce, Aclytia; Epectaptera laura Schaus, Eumaschane; Eumaschane lauta Scott, Aglaosoma; Aglaosoma lavinia Druce, Idalus; Hyponerita lavinia Fawcett, Phalera; Eulavinia laymerisa Grandidier, Lithosia; Axiopoeniella lectrix Linnaeus, Phalaena; Episteme lenistriata Dognin, Microdota; Trocodima lentiginosa Collenette, Adetoneura; Adetoneura leonensis Hampson, Cerura; Afrocerura leonina Kiriakoff, Jozinia; Jozinia leopoldi Hering, Leopoldina; Leopoldina

lephina Field, Epeiromulona; Epeiromulona lepidocaena Dyar, Dixanaene; Dixanaene lepidoides Dognin, Pseudhemiceras; Pseudhemiceras leptalina Druce, Areva; Disaulota leptopepla Hampson, Cymaroa; Cymaroa leucanina Felder & Rogenhofer, Halysidota; Leucanopsis leucanioides Herrich-Schäffer, Idioctetus; Idioctetus leucocloea Collenette, Perinetia; Perinetia leucomela Herrich-Schäffer, Phaegorista; Stemonoceras leuconephra Hampson, Amphiphalera; Amphiphalera leuconoe Hopffer, Nyctemera; Chiromachla leucophaea Walker, Ela; Urocoma leucophaea Walker, Theages; Theages leucoptera Hampson, Automolis; Nyearctia leucospila Walker, Cozola; Cozola leucospilota Moore, Pitasila; Pitasila leucostigma Smith, Phalaena; Hemerocampa leucota Hampson, Odozana; Saozana levis Druce, Phaegoptera; Xanthophaeina lianga Semper, Satara; Heliozona liboria Stoll, Phalaena; Exotrocha lichenicolor Oberthür, Semidonta; Rachiades lichenigera Felder, Carcinopyga; Carcinopyga lichenina Butler, Somera; Parasinga licormas Cramer, Phalaena; Caroia lignaria Walker, Listoca; Listoca lignea Butler, Nioda; Ankova lignea Schaus, Naduna; Naduna ligneata Walker, Etobesa; Etobesa ligniclusa Walker, Braura; Braura lignicolor Walker, Ianassa; Ianassa lignifera Walker, Sarbena; Sarbena lignilutea Walker, Biula; Biula lignitecta Walker, Lirimiris; Lirimiris lignosa Felder, Leucaniochroa; Leucaniochroa lignosa Möschler, Elasmia; Elasmia lignosa Walker, Petrinia; Petrinia lignosa Walker, Xulora; Xulora lignosa Walker, Zana; Zana ligula Hübner, Hypocrita; Josia lilacea Kiriakoff, Janthinisca; Janthinisca limacodiformis Strand, Dactylorhynchides;

Dactylorhynchides limacodoides Walker, Alimala; Alimala limbalis Herrich-Schäffer, Porthesia; Urocoma limbaria Warren, Hadesina; Hadesina limnatis Schaus, Navarcostes; Navarcostes limona Schaus, Pseudalus; Pseudalus limonensis Thiaucourt, Lemairegisa; Lemairegisa limpida Swinhoe, Sapelia; Sapelia lineata Matsumura, Macroshachia; Macroshachia lineata Oberthür, Notodonta; Epodonta lineata Walker, Destolmia; Destolmia lineata Walker, Lymantria; Psilochira lineola van Eecke, Plumareola; Plumareola lineolata Kiriakoff, Gastrochrysia; Gastrochrysia lineolata Walker, Dinara; Dinara linta Moore, Artaxa; Micronygmia lirioleuca Turner, Panachranta; Panachranta lithosiana Hampson, Acanthofrontia; Acanthofrontia lithosioides Moore, Baradesa; Baradesa lithosioides Snellen, Tampea; Tampea litura Walker, Charnidas; Charnidas liturata Guérin-Méneville, Orgyia; Creagra liturata Walker, Cerura; Neocerura lixivia Dognin, Orgya; Blazia lobata Strand, Melisoides; Melisoides lobifer Herrich-Schäffer, Creatonotos; Sutonocrea lobula Hübner, Eubaphe; Eubaphe locuples Walker, Cifuna; Cifuna lodra Moore, Euproctis; Bradytera longa Holland, Dinotodonta; Dinotodonta

longicauda Holland, Macrosenta; Macrosenta longicornis Herrich-Schäffer, Agaraea; Agaraea longinasus Joannis, Bifurca; Bifurca longinervis Felder, Gnatholophia; Gnatholophia longipennis Moore, Norraca; Norraca longipennis Neumoegen, Phaio; Phaio longipennis Walker, Damata; Damata longipennis Walker, Hypercompa; Nikaea longipennis Walker, Lithosia; Pteroodes longivitta Felder, Josiomorpha; Josiomorpha longivitta Gaede, Allodonta; Kiriakoffia longivitta Kiriakoff, Psalisodes; Gargettina longivitta Walker, Bireta; Bireta lophocampoides Felder, Symphlebia; Symphlebia lophura Aurivillius, Paramelisa; Paramelisa lorquinii Felder, Arctioneura; Arctioneura lowii Butler, Kettelia; Kettelia lubricipeda Linnaeus, Phalaena; Spilosoma lucasii Rosenstock, Discophlebia; Themerastis lucilinea Walker, Bardaxima; Bardaxima lucinia Druce, Automolis; Diaxanthia luctifera Walker, Anaphela; Anaphela luctosa Oberthür, Naclia; Melanonaclia luctuosa Kiriakoff, Microbergeria; Microbergeria lugubris Gaede, Pygaerina; Pygaerina lunifer Herrich-Schäffer, Ochrogaster; Ochrogaster lunifera Walker, Adlullia; Adlullia lunifera Walker, Zarfa; Zarfa luridipennis Burmeister, Cydosia; Progona lutea Druce, Lithosia; Phaeosia lutea Hufnagel, Phalaena; Spilarctia lutea Kiriakoff, Sericanaphe; Sericanaphe luteata Holloway, Cofimpacia; Cofimpacia luteicincta Holland, Hemonia; Paremonia luteilinea Druce, Heterocampa; Azaxia luteovirens Felder, Psilacron; Psilacron luxa Grofe, Euhalisidota; Euhalisidota luzonica Felder & Rogenhofer, Cyme; Xanthetis lycoides Oberthür, Mimica; Mimica lycoides Walker, Euchromia; Pionia lycone Druce, Ichoria; Holophaea lygropis Collenette, Dasychira; Sirana lymantrioides Felder, Caenina; Caenina lymantrioides Kiriakoff, Pinheyia; Pinheyia

macrodonta Hampson, Ramesa; Rasemia maculapex Hampson, Bracharthron; Bracharthron maculata Harris, Lophocampa; Phoegoptera maculata Moore, Heterocampa; Polystictina maculatrix Walker, Godasa; Godasa maculifascia Moore, Aemene; Parasiccia maculifer Matsumura, Hexafrenum; Hexafrenum maculosa [Denis & Schiffermüller], Phalaena; Chelis maculosa Matsumura, Takashachia; Takashachia maculosa Stoll, Phalaena; Alpenus madagascariensis Boisduval, Chelonia; Phryganeomorpha madagascariensis Draeseke, Desmeocraera; Vietteella magnaria Felder, Ophitis; Ophitis magnifica Walker, Chrysocale; Gardinia magnifica Walker, Lithosia; Chrysaeglia magniplaga Schaus, Pronola; Pronola maguila Schaus, Fentonia; Maguila maja Fabricius, Zygaena; Macrocneme majas Jörgensen, Hassleria; Hassleria makala Bethune-Baker, Olapa; Pseudogenusa malangae Bethune-Baker, Leptolepida; Leptolepida malgassica Kenrick, Stauropus; Griphocerura malgassica Kiriakoff, Nesoptilura; Nesoptilura malgassica Toulgoët, Spatulosia; Spatulosia

maligna Butler, Parorgyia; Gallienica mangholda Schaus, Fentonia; Erconholda manleyi Leech, Drymonia; Naganoea manteo Doubleday, Lochmaeus; Lochmaeus marcescens Felder, Agaposoma; Agaposoma margarethae Kiriakoff, Deriddera; Deriddera margarethae Kiriakoff, Margaropsecas; Margaropsecas margaritaceus Walker, Aphyle; Aphyle marginalis Matsumura, Formotensha; Formotensha marginalis Walker, Amsacta; Amsacta marginata Drury, Phalaena; Are marginata Moore, Cobanilla; Cobanilla marginata Walker, Nudaria; Xanthocraspeda marginepuncta Griveaud, Vadonaclia; Vadonaclia marginepunctata Saalmüller, Cypra; Scaphocera mariae Dufrane, Pseudothyretes; Pseudothyretes mariamne Geyer, Eucharia; Daritis marita Schaus, Pamcoloma; Pamcoloma marmorata Hampson, Nolosia; Nolosia marmorata Kiriakoff, Blakeia; Blakeia marmorea Butler, Calledema; Calledema marmorea Schaus, Graphea; Graphea marojejia Kiriakoff, Desmeocraera; Ochrosomera marpissa Wallengren, Eutimia; Eutimia marshalli Hampson, Cerura; Cerurina marthesia Cramer, Phalaena; Macrurocampa martynovi Kusnezov, Oligamatites; Oligamatites mascarena Butler, Dasychira; Lemuriana mathani Rothschild, Ceramidia; Ceramidiodes matronula Linnaeus, Phalaena; Pericallia maura Eichwald, Bombyx; Axiopoena meeki Rothschild, Chadisra; Timoraca meeki Rothschild & Jordan, Eucocytia; Eucocytia megaera Fabricius, Bombyx; Phalcidon megalopia Schaus, Arpema; Arpema megaspila Walker, Talara; Talara melaleuca Walker, Lambula; Lambula melanchroia Boisduval, Cocastra; Acribia melanistis Bethune-Baker, Parathemerastis; Parathemerastis melanitis Hübner, Centronia; Centronia melanocephala Walker, Lymire; Lymire melanochlora Hampson, Malachitis; Malachitis melanochloros Sepp, Sesia; Illipula melanogramma Janse, Hypophiala; Hypophiala melanolepia Dudgeon, Miltochrista; Cyclomilta melanopasta Dognin, Idalus; Arctiarpia melanopsis Walker, Arctia; Rhodareas melanoptera Turner, Pseudophanes; Pseudophanes melanora Meyrick, Agaphthora; Agaphthora melanosticta Joicey & Talbot, Gargetta; Gargettiana melas Dyar, Ardonipsa; Ardonipsa melaxantha Hübner, Hipocrita; Ephestris melissograpta Collenette, Dasychira; Vohitra membliaria Cramer, Phalaena; Euplocia menas Doubleday, Stauropus; Balia mendica Clerck, Phalaena; Diaphora mendosa Hübner, Olene; Olene menea Drury, Phalaena; Ruscino menthastri [Denis & Schiffermüller], Bombyx; Spilosoma mentiens Walker, Crorema; Crorema mentiens Walker, Virbia; Virbia meona Stoll, Phalaena; Eulophopteryx meones Stoll, Sphinx; Euclera merra Wallengren, Lasioprocta; Lasioprocta mesembrinum Collenette, Aphomoeoma; Aphomoeoma mesogona Godart, Callimorpha; Apaidia mesomelaena Holland, Artaxa; Euproctillina mesomella Linnaeus, Phalaena; Ecteina mesopotamia Hampson, Chrysocyma; Chrysocyma metabus Cramer, Phalaena; Antarctia metachrysea Druce, Phoenicoprocta; Phoenicoprocta

metaleuca Hampson, Lamoriodes; Lamoriodes

metalica Dognin, Odozana; Procridia metallica Dyar, Serincia; Serincia metallica Gaede, Neardonaea; Neardonaea metallica Kiriakoff, Zadadrina; Zadadrina metamelaena Hampson, Carcinarctia; Carcinarctia metamelaena Hampson, Neotalara; Neotalara metamelas Walker, Glaucopis; Mallostethus metaphaea Walker, Ceira; Ceira metarhoda Walker, Spilosoma; Hyarias metaxanthus Walker, Glaucopis; Herea meteus Stoll, Sphinx; Automolis methyalea Hampson, Lithoprocris; Lithoprocris meticulodina Oberthür, Notodonta; Uropyia mexicana Druce, Lithosia; Haematomis mhondana Bartel, Spilosoma; Eminaria miara Hering, Theriophila; Theriophila micans Bremer & Grey, Setina; Stigmatophora miccyla Collenette, Papuaroa; Papuaroa michaellae Laporte, Koffoleania; Koffoleania micilia Cramer, Phalaena; Agyrta micrommata Turner, Eurodes; Eurodes microsippia Strand, Meganaclia; Nacliodes microsticta Hampson, Damata; Damatoides mida Dyar, Palaeozana; Palaeozana miles Butler, Sphragidium; Sphragidium milhauseri Fabricius, Bombyx; Harpyia militaris Butler, Dycladia; Hypocladia militta Stoll, Phalaena; Thyrgis milupa Nye, Noliproctis; Noliproctis mimetica van Eecke, Snellenopsis; Snellenopsis mimetica Rothschild, Caprimimodes; Caprimimodes mimica Butler, Schasiura; Schasiura mimica Druce, Pseudopompilia; Pseudopompilia minceus Stoll, Sphinx; Eutomis mingens Herrich-Schäffer, Anurocampa; Anurocampa miniata Bethune-Baker, Euproctoides; Euproctoides miniata Forster, Phalaena; Miltochrista minima Butler, Eustixis; Diarhabdosia ministra Drury, Phalaena; Datana minor Hampson, Meganaclia; Mesonaclia minuta Druce, Actea; Leptactea minuta Druce, Crinodes; Rincodes minuta Druce, Nola; Neoselca minuta Hampson, Paurophleps; Paurophleps minuta Hulstaert, Deloplotela; Deloplotela minuta Rothschild, Halysidota; Nannodota minx Dyar, Geridixis; Geridixis mioides Hampson, Stauropus; Miostauropus mira Butler, Agylla; Hyperthagylla mirabilis Aurivillius, Pirga; Pirga mirabilis Butler, Barsinella; Barsinella mirabilis Draudt, Chrysocneme; Chrysocneme mirabilis Viette, Anaphe; Nesanaphe miraculosa Staudinger, Himeropteryx; Himeropteryx mirax Prout, Dolophrosyne; Dolophrosyne mirifica Butler, Choerotriche; Chrysopsyche mirifica Oberthür, Chelonia; Preparctia miroides Hampson, Metagylla; Metagylla miselioides Kenrick, Dasychira; Radamaria mixtipennis Walker, Gopha; Gopha modesta Butler, Atyphopsis; Atyphopsis modesta Holland, Scrancia; Scrancia modesta Schaus, Homoneuronia; Homoneuronia modesta Schaus, Kurtia; Kurtia modesta Walker, Morasa; Morasa modestus Distant, Polienus; Polienus modicus Lucas, Diphtheraspis; Diphtheraspis molliculana Walker, Cabarda; Cabarda mollis Lucas, Nudaria; Psilopepla moloneyi Druce, Anaphe; Epanaphe moluccana Felder, Ortholomia; Ortholomia

molybdipera Schaus, Lamprostola; Lamprostola

moma Schaus, Automolis; Viviennea monacha Linnaeus, Phalaena; Lymantria monacheicauda Dyar, Grucia; Grucia monetaria Oberthür, Notodonta; Peridea monilis Hübner, Hipocrita; Mitradaemon moninna Schaus, Carilephia; Carilephia monochroa Turner, Aedoea; Aedoea monogrammaria Walker, Chiriphe; Chiriphe monotica Holland, Apisa; Anapisa monteiroi Butler, Thyretes; Eressades monticola Roepke, Pydnella; Pydnella morio Linnaeus, Phalaena; Penthophera morio Walker, Ardonea; Ardonea morona Druce, Edema; Drugera morosa Schaus, Hyaleucerea; Prosopidia moruma Schaus, Salluca; Salluca mucescens Felder, Dasysphinx; Dasysphinx multifaria Walker, Apistosia; Euctenucha multilinea Schaus, Antiopha; Antiopha multilinea Walker, Magava; Magava multilineata Gaede, Solwezia; Solwezia multipunctata Hampson, Aeolosia; Aeolosia multistrigata Hampson, Eugoa; Pareugoa mumetes Cramer, Phalaena; Hippia munda Kiriakoff, Antoroka; Antoroka munda Walker, Lymantria; Imaus mundana Linnaeus, Phalaena; Nudaria muricolor Walker, Lithosia; Poliosia murina Heylaerts, Lithosia; Bitecta murina Hübner, Phalaena; Paidia murina Kiriakoff, Plastystaura; Plastystaura murina Walker, Comacla; Comacla muscella Mabille, Syntomis; Maculonaclia muscerda Hufnagel, Phalaena; Pelosia muscosa Walker, Cascera; Cascera muscula Staudinger, Nudaria; Nudaridia musculus Kiriakoff, Antsalova; Antsalova musiforme Kaye, Pseudodiptera; Pseudodiptera myiella van Eecke, Eressa; Nannoceryx myopsychoides Strand, Epitoxis; Syntomerea mystacina Kiriakoff, Nyssiopsis; Nyssiopsis mystica Walker, Euchromia; Pampa

naclioides Felder, Cacosoma; Cacosoma naclioides Felder, Ctenandra; Ctenandra naenia Druce, Pseudocharis; Pseudocharis nana Bryk, Podocryptula; Podocryptula nana Druce, Calonotos; Metaloba nana Hampson, Paratuerta; Hemituerta nana Swinhoe, Bireta; Dragonera nana Walker, Agyrta; Arctagyrta nantana Walker, Ammalo: Hoplarctia natalensis Kiriakoff, Cerurella; Cerurella nebulosa Butler, Rhyparioides; Rhyparioides nebulosa Toulgoët, Agaltara; Agaltara nebulosa Wileman, Pydna; Kuohsingia necyria Felder & Rogenhofer, Sangala; Cyanotricha negrottoi Berio, Croperoides; Croperoides nephocrossa Bethune-Baker, Elaphrodes; Elaphrodes nerias Felder, Adelphoneura; Adelphoneura nero Weymer, Amastus; Mellamastus nesiotica Collenette, Carpenterella; Carpenterella neurabrunnea Bethune-Baker, Parapirga; Parapirga neurias Schaus, Hypidota; Hypidota neuroptera Sepp, Phalaena; Sontia nevosa Dognin, Areva; Neagylla nexa Boisduval, Lithosia; Byssophaga niepelti Gaede, Agyrtiola; Agyrtiola nieuwenhuisi Roepke, Notodontella; Notodontella

nigeriensis Laporte, Medlerana; Medlerana nigra Aurivillius, Scrancia; Subscrancia nigra Laporte, Deinhugia; Deinhugia nigrapex Strand, Phryganopsis; Tesma nigrescens Kiriakoff, Induba; Induba nigribursa Chou & Lu, Shensiplusia; Shensiplusia nigricosta Felder, Xenosoma; Xenosoma nigricosta Hampson, Oedaleosia; Oedaleosia nigricosta Leech, Oenistis; Conilepia nigricostata Butler, Ghoria; Onychipodia nigrifera Walker, Cothocida; Cothocida nigrifrons Walker, Icambosida; Icambosida nigripalpis Walker, Inous; Inous nigripennis Aurivillius, Megapisa; Megapisa nigripes Heylaerts, Laemocharis; Neotrichura nigripuncta Hampson, Mulona; Lomuna nigripuncta Rothschild, Turnaca; Trabanta nigripunctella Viette, Dinotodonta; Malgadonta nigrisquamata Swinhoe, Pseudoblabes; Melastrota nigristriata Holland, Tegulata; Pliniola nigrivena Schaus, Ischnocampa; Pseudischnocampa nigrocincta Aurivillius, Paradiastema: Paradiastema nigrolinea Lucas, Notodonta; Paradestolmia nigrolineata Aurivillius, Haplozana; Haplozana nigrolineata Joicey, Noakes & Talbot, Epicoma; Tearina nigronervata Gaede, Odontosina; Odontosina nigropunctata Bethune-Baker, Eminaria; Eminaria nigropunctata Wallengren, Lithosia; Siccia nigropunctum Hampson, Holocraspedon; Holocraspedon nigrorosea Walker, Boenasa; Boenasa nigroruba Rego Barros, Ypiranga; Lepypiranga nigrostriata Möschler, Lomatosticha; Lomatosticha nihonica Wileman, Fentonia; Eufentonia niobe Weymer, Porthetria; Argyrostagma nitens Gaede, Lamorina; Lamorina nitens Walker, Setora; Setora nitida Cramer, Phalaena; Zatrephes nivea Aurivillius, Acantharctia; Acantharctia nivea Bethune-Baker, Pseudarctia; Pseudarctia nivea Herrich-Schäffer, Charidea; Eupseudosoma nivea Kiriakoff, Menegites; Menegites nivea Matsumura, Shironia; Shironia niveiceps Walker, Trabala; Euhampsonia niveifascia Walker, Getta; Getta niveimaculata Hampson, Lobobasis; Lobobasis niveipicta Bryk, Pheosiopsis; Pheosiopsis niveipicta Kiriakoff, Bireta; Norracana nivosa Walker, Lerna; Lerna nobilis Walker, Dudusa; Dudusa nobilitella Cramer, Phalaena; Cydosia noctella Dyar, Zamolis; Zamolis noctuiformis Barnes & McDunnough, Ursia; Ursia noctuiformis Hampson, Euplexidia; Euplexidia noctuiformis Janse, Pectinophora; Eujansea nodicornis Guenée, Hapigia; Hapigia nodyna Swinhoe, Notodonta; Macroshachia noliformis Strand, Chrysaegliodes; Chrysaegliodes nomo Kiriakoff, Parosica; Parosica nonagrioides Walker, Saliocleta; Saliocleta norema Schaus, Nesebra; Nesebra notabilis Walker, Glaucopis; Methysia notata Butler, Laelapia; Laelapia notodontina Rothschild, Stauropus; Hoplitina notodontoides Walker, Elymiotis; Elymiotis novicia Draudt, Cercocladia: Cercocladia nubica Strand, Cinciana; Cinciana nubifera Hampson, Chadisra; Nepheliphora nubifuga Holland, Notohyba; Notohyba nubila Butler, Birnara; Birnara nubila Druce, Rifargia; Euphastia nubila Schaus, Pauluma; Pauluma nubila Walker, Sorema; Sorema

nubilosa Hampson, Steneugoa; Steneugoa nubilosa Staudinger, Nudaria; Nudina nuda Fabricius, Bombyx; Perina nuda Hampson, Achroosia; Achroosia nudarioides Butler, Homopsyche; Homopsyche nudiuscula Aurivillius, Hyaloperina; Hyaloperina nugatrix Felder, Desmeocraera; Desmeocraera nulliferana Walker, Tospitis; Tospitis nyctemeroides Walker, Olina; Aphra nyikae Kiriakoff, Umncumbata; Umncumbata nyseus Cramer, Phalaena; Cyrrhesta

oberthueri Rothschild, Micronaclia; Tenuinaclia oberthuri Hampson, Correbia; Mimica obliqua Walker, Turuptiana; Turuptiana obliqualinea Bethune-Baker, Dasychirana; Dasychirana obliqualinea Bethune-Baker, Lymantriades; Lymantriades obliquaria Warren, Authyala; Authyala obliquilinea Walker, Comidava; Comidava obliquipuncta Rothschild, Thyrosticta; Toulgoetinaclia obliquistria Hampson, Streptoplebia; Streptophlebia obliterata Wileman & South, Stauropus; Palaeostauropus oblonga Toulgoët, Philenora; Astacosia obryzos Mabille, Eusemia; Schausilla obscura Birket-Smith, Yelva; Yelva obscura Butler, Ceramidia; Ceramidiodes obscura Wileman, Phalera; Horishachia obscurata Butler, Neritos; Phaeomolis obsoleta Bethune-Baker, Dasychiroides; Dasychiroides obsoleta Kiriakoff, Periergos; Periergos obtusa Walker, Arina; Polypoetes obtusa Walker, Cassidia; Cassidia occidentalis Strand, Pusiola; Pusiolania ocellata Kenrick, Dasychira; Kintana ocellatula Hering, Dasychira; Kintana ocellifera Walker, Alope; Olepa ochracea Bethune-Baker, Blaviodes; Blaviodes ochracea Bethune-Baker, Neorgyia; Neorgyia ochracea Bethune-Baker, Polienus; Lopiena ochracea Felder, Gymnopoda; Apeplopoda ochraceorufa Rothschild, Melanaema; Diaconisia ochribasis Hampson, Chadisra; Achaera ochricostata Gaede, Metarctina; Metarctina ochrivena Hampson, Eugoa; Meteugoa ochroderoea Mabille, Dasychira; Alina ochrogutta Herrich-Schäffer, Cnethocampa; Tanystola ochropasa Dyar, Exemasia; Exemasia ochropis Hampson, Phalera; Epiphalerina ochrosterna Felder, Sallaea; Sallaea ochsenheimeri Boisduval, Syntomis; Myopsyche ochthoeba Hampson, Turuptiana; Tamsita ockendeni Rothschild, Automolis; Melanarctia ocularia Fabricius, Bombyx; Ecpantheria oculata Kiriakoff, Spodiosomera; Spodiosomera oculata Kiriakoff, Ulina; Ulina oculata Toxopeus, Xanthonygmia; Xanthonygmia ocypete Bremer, Harpyia; Urocampa oedippus Boisduval, Mastigocera; Drucea ogovensis Holland, Ctenogyna; Paratrotonotus olivacea Gaede, Eurystaura; Eurystauridia olivacea Möschler, Pseudodryas; Pseudodryas olivacea Schaus, Odozana; Lamprostola olivina Kiriakoff, Scarnica; Scarnica ombrophanes Meyrick, Anestia; Anestia omophanes Meyrick, Scaeodora; Scaeodora omphale Hübner, Cosmosoma; Cosmosoma oneili Janse, Prionocentrum; Prionocentrum onytes Cramer, Phalaena; Trichromia oophora Zeller, Pseudoblabes; Pseudoblabes

opaca Walker, Myrmecopis; Phaeosphecia opalescens Rebel, Pseudogenusa; Pseudogenusa opalina Walker, Cimola; Cimola opalocincta Druce, Urolosia; Urolosia opharina Schaus, Amastus; Romualdia ophidera Dognin, Euphastia; Euphastia ophion Walker, Glaucopis; Argyroeides opponens Walker, Arunda; Arunda opponens Walker, Pampa; Pseudaclytia opposita Walker, Euchromia; Apiconoma opulentana Walker, Maepha; Maepha orbicularis Walker, Hypsa; Peridrome orbifer Hampson, Stauropus; Formofentonia orbiferana Walker, Hemonia; Hemonia orbigera Herrich-Schäffer, Coryphyala; Coryphyala orcus Walker, Sanega; Sanega orientalis Walker, Areas; Areas orion Esper, Noctua; Diphteramoma orislinea Walker, Mennis: Mennis ornata Oberthür, Ptilodontis; Eguria ornata Walker, Cynosarga; Cynosarga ornatissima Walker, Plusia; Antoculeora ornatrix Linnaeus, Phalaena; Utetheisa osiris Bethune-Baker, Paraproctis; Paraproctis osmophora Hampson, Disoidemata; Disoidemata otanes Druce, Dioptis; Paradioptis ovale Hampson, Idopterum; Idopterum ovalis Eecke, Stauropus; Vaneeckeia owamboensis Kiriakoff, Owambarctia; Owambarctia oweni Schaus, Castrica; Castrica oxiana Djakonov, Paragluphisia; Paragluphisia

pacifera Schaus, Clemendana; Clemendana paetula Hering, Laeliolina; Laeliolina paidicus Dyar, Gaudeator; Gaudeator pallida Butler, Bireta; Mimopydna pallida Butler, Cincia; Amplicincia pallida Butler, Hyperaeschra; Hyperaeschra pallida Hampson, Charnides; Dactylorhyncha pallida Moore, Korawa; Korawa pallida Packard, Crambidia; Crambidia pallida Schaus, Eunotela; Eunotela pallida Schaus, Kaseria; Kaseria pallidicosta Walker, Onythes; Onythes pallidifascia Hampson, Stauropus; Vaneeckeia pallidior Dyar, Palpidia; Palpidia pallidistriata Hampson, Dialeucias; Dialeucias pallidula Grünberg, Meneclia; Meneclia palliola [Denis & Schiffermüller], Noctua; Nola palmeri Druce, Urgedra; Herbulotedra palpina Clerck, Phalaena; Euchila pampa Druce, Metastatia; Mimagyrta panda Boisduval, Bombyx; Zastonia pandula Boisduval, Lithosia; Lepista pannosa Moore, Zaranga; Zaranga panochra Turner, Ameleta; Ameleta pantheraria Felder, Lomaspilis; Argina parabolica Matsumura, Neoshachia; Neoshachia paradoxa Butler, Chionophasma; Chionophasma paradoxa Hampson, Acidon; Acidon paradoxa Hering, Balacra; Neobalacra paradoxa Hering, Marbloides; Marbloides paradoxa Schultze, Neoliparina; Neoliparina parallela Felder, Cratosia; Cratosia pardalina Püngeler, Tancrea; Tancrea pardalina Walker, Minara; Minara pardalis Walker, Amaxia; Amaxia parnassia Möschler, Marissa; Parastatia parthenos Harris, Arctia; Platarctia

parthia Druce, Ichoria: Cloesia parva Barnes & McDunnough, Aresia; Aresia parva Hampson, Notata; Notata passalis Fabricius, Zygaena; Amata pastinacella Duponchel, Haemilis; Piesta patagiata Berg, Chlanidophora; Chlanidophora patula Walker, Thelde; Thelde pectinicornis Smith, Eucalyptera; Aretypa peculiaris Butler, Mardara; Masoandro peculiaris Butler, Orthoclostera; Orthoclostera pedrana Schaus, Corania; Corania pega Dognin, Agkonia; Agkonia pelecoides Tams, Boadicea; Boadicea pelias Kiriakoff, Peltostaura; Peltostaura pellex Linnaeus, Phalaena; Atasca pellopis Bethune-Baker, Scoliacma; Hyposhada pellucens Herrich-Schäffer, Psycharium; Psycharium penates Druce, Neotrichura; Neotrichura penetrata Walker, Josia; Josiomorpha penicillata Walker, Binna; Binna pennatula Fabricius, Bombyx; Arestha pennipes Grote, Callicarus; Callicarus pentacyma Meyrick, Hectobrocha; Hectobrocha pentelia Druce, Eusemia; Sergiusia perbella Schaus, Thyone; Thyonaea perculta Distant, Aroa; Pseudobazisa pergrisea Hampson, Stauropus; Epiphalera perinetensis Kiriakoff, Scalmicauda; Analama perisema Turner, Axiocleta; Axiocleta permagna Butler, Drymonia; Epizaranga perornata Moore, Gonerda; Gonerda perornata Walker, Bizone; Bizone perpallida Hampson, Miltochrista; Korawa perplexa Schaus, Talmeca; Talmeca perpusilla Mabille, Aglaope; Tsarafidynia perrottetii Guérin-Méneville, Chelonia; Rajendra persimilis Hampson, Chadisra; Disracha pertyi Herrich-Schäffer, Laemocharis; Laemocharis pexicera Schaus, Prepiella; Prepiella phaeocephala Meyrick, Petalopleura; Petalopleura phaeocraspis Hampson, Crocosia; Crocosia phaeocyma Meyrick, Neobrocha; Neobrocha phaeogramma Turner, Oxymetopa; Oxymetopa phaeoplaga Schaus, Rhodographa; Rhodographa phalaenoides Drury, Sphinx; Castrica phalaenoides Toulgoët, Turlinia; Turlinia phalarota Meyrick, Thallarcha; Thallarcha phantoma Dalman, Bombyx; Pania phasma Harvey, Euerythra; Enerythra phegea Linnaeus, Sphinx; Syntomis pheloides Felder, Monocreagra; Monocreagra phemonoe Hübner, Antichloris; Antichloris phemonoides Möschler, Antichloris; Passineura phileta Drury, Phalaena; Caryatis philocosma Meyrick, Atopomima; Atopomima phlegmon Cramer, Sphinx; Calonotos phoebe Siebert, Phalaena; Tritophia pholus Drury, Sphinx; Lycomorpha phryganea Kiriakoff, Scranciella; Scranciella phryganeoides Felder, Tmetoptera; Tmetoptera phyllodes Meyrick, Palaexera; Palaexera pica Butler, Drymonia; Eucerura pica Walker, Halysidota; Galethalea picaria Walker, Balaca; Balaca picta Boisduval, Liparis; Lacipa picta Butler, Pachycispia; Pachycispia picta Druce, Paramya; Metamya picta Felder, Trichotis; Trichotis picta Hampson, Notodonta; Khasidonta picta Herrich-Schäffer, Eurata; Eurata picta Walker, Deiopeia; Tatargina pictibasis Hampson, Pheosia; Chokaia

pictula Walker, Glaucopis; Eurota pictus Kiriakoff, Subwilemanus; Subwilemanus pierreti Rambur, Nototrachus; Nototrachus pilicornis Herrich-Schäffer, Trichaea; Trichaea pilosa Warren, Uliolepis; Uliolepis pilumnia Stoll, Phalaena; Purius pinheyi Kiriakoff, Pseudagoma; Pseudagoma pinivora Treitschke, Gastropacha; Traumatocampa piperita Herrich-Schäffer, Phaegoptera; Bernathonomus pirgula Hering, Marblepsis; Marblepsis pitthea Druce, Enioche; Enioche plagiata Walker, Anaxila; Anaxila plagiata Walker, Castulo; Castulo plagiata Walker, Hypsa; Asota plagiata Walker, Teulisna; Teulisna plagidisca Hampson, Eucyclopera; Eucyclopera plagidotata Walker, Cyclidia; Mahoba plagifera Matsumura, Metasomera; Metasomera plagifera Walker, Lyces; Cacolyces plagifera Walker, Nyctemera; Trypheromera plagipennis Dyar, Brecontia; Brecontia plagiviridis Moore, Heterocampa; Viridifentonia plagosus Gaede, Pseudostauropus; Pseudostauropus plana Kiriakoff, Batempa; Batempa plantaginis Linnaeus, Phalaena; Parasemia platyzona Felder, Scytale; Clystea plebeja Oberthür, Notodonta; Allodonta plexa Möschler, Lysana; Lysana plicata Butler, Maepha; Callisthenia plumbea Hampson, Poliopastea; Poliopastea plumicornis Butler, Coracia; Viettesia plumigera [Denis & Schiffermüller], Bombyx; Ptilophora plumigera Matsumura, Higena; Higena plumipes Drury, Sphinx; Phylloecia plumitarsus Janse, Polelassothys; Polelassothys plumosa Hampson, Pteredoa; Pteredoa plumosa Kiriakoff, Pararheneades; Pararheneades plumosa Leech, Bireta; Torigea plumosa Moore, Rachia; Rachia plusiata Walker, Celeia; Celeia plusioides Bryk, Pydna; Ogulina plusioides Walker, Eleale; Tamseale plutonica Felder, Meringocera; Meringocera podarce Walker, Josia; Phintia poecila Felder, Nycterotis; Nycterotis pogonoda Hampson, Hypareva; Hypareva polia Druce, Psilopleura; Psilopleura polidamon Cramer, Sphinx; Cacoethes polioplaga Hampson, Macrosia; Lophilema polioplaga Hampson, Stenostaura; Boscawenia poliotis Hampson, Gerarctia; Gerarctia polioxutha Turner, Pheraspis; Pheraspis polistes Hübner, Pseudosphex; Myrmecopsis politia Stoll, Phalaena; Sericochroa pollinia Boisduval, Ctenucha; Stenophaea polygrapha Felder, Cyme; Graptasura polymena Linnaeus, Sphinx; Phalanna polyxenus Druce, Evius; Paranerita porpara Schaus, Noctulodes; Noctulodes postalba Wileman, Pida; Dokuga postalbida Hampson, Liparopsis; Liparopsis postflava Schaus, Metacrocea; Metacrocea postica Moore, Ceira; Biraia postica Walker, Atrasana; Atrasana postica Walker, Melese; Melese praestans Saalmüller, Numenes; Peloroses praeusta Felder, Anchyneura; Anchyneura prasena Moore, Tripura; Tripura prasina Butler, Calliteara; Labordea prasina Kiriakoff, Iphigeniella; Iphigeniella pratti Bethune-Baker, Parabasis; Parabasis pretus Cramer, Sphinx; Horama

preussi Aurivillius, Metarctia; Epibalacra preussi Mabille & Vuillot, Liparis; Abynotha priscilla Schaus, Automolis; Amphelarctia prittwitzi Möschler, Desmidocnemis; Desmidocnemis processionea Linnaeus, Phalaena; Thaumetopoea procne Schaus, Symmerista; Gisara procridiformis Hampson, Leptopepla; Leptopepla procris Felder & Rogenhofer, Lithosia; Procrimima procroides Walker, Opharus; Opharus producta Kiriakoff, Kumataia; Kumataia prolifera Walker, Josia; Glissa promathides Druce, Eucereon; Protomolis pronapides Druce, Zatrephes; Ochrodota propylea Druce, Scena; Scena proserpina Hübner, Euchromia; Histioea proserpina Staudinger, Euarctia; Euarctia protea Kiriakoff, Neochadisra; Neochadisra protea Kiriakoff, Neochadisra; Neochadisra prouti Hering, Xenormicola; Xenormicola psecas Druce, Cleapa; Paracleapa psectriphora Distant, Pusiola; Ctenosia pseudameoides Rothschild, Automolis; Xanthoarctia pseudotortrix Toulgoët, Philenora; Mimulosia pteridaula Turner, Eurypepla; Eurypeplella pterostomina Kiriakoff, Epiptilodontis; Epiptilodontis ptochias Meyrick, Tylanthes; Tylanthes ptychoglene Hampson, Propyria; Propyria pua Dyar, Metacrisiodes; Metacrisiodes pudibunda Linnaeus, Phalaena; Elkneria pudica Esper, Phalaena; Cymbalophora pudica Kiriakoff, Lomela; Lomela puella Boisduval, Naclia; Pseudonaclia puella Drury, Phalaena; Chionaema pugione Linnaeus, Sphinx; Empyreuma pulchella Linnaeus, Phalaena; Deiopeia pulcherrima Moore, Anodonta; Hupodonta pulchra Aurivillius, Balacra; Lamprobalacra pulchripicta Walker, Mirobriga; Birgorima pulverina Neumoegen, Bruceia; Bruceia pulverosa Schaus, Automolis; Astralarctia pumila Butler, Dasychira; Noliproctis pumila Staudinger, Dasychira; Dasorgyia punctalis Brandt, Mekrania; Mekrania punctana Walker, Tospitis; Narosodes punctata Griveaud, Turlina; Turlina punctata Guérin-Méneville, Phylloecia; Phylloecia punctata Hampson, Palaeosiccia; Palaeosiccia punctata Hampson, Philenora; Exilisia punctata Moore, Ranghana; Ranghana punctata Stoll, Phalaena; Malocampa punctatus Druce, Purius; Hypocrisias puncticilia Moore, Naxa; Axana punctifascia Walker, Cispia; Cispia punctifera Walker, Caschara; Caschara punctilineata Moore, Kerala; Kerala punctivaga Walker, Cycnia; Baroa punctum Gaede, Fentonina; Fentonina pura Hulstaert, Paramsacta; Paramsacta pura Lucas, Teara; Axiologa purata Walker, Galtara; Galtara purissima Butler, Plusia; Puriplusia purpurascens Hampson, Palaeomolis; Palaeomolis purpurascens Schaus, Xantholopha; Xantholopha purpurascens Walker, Sarsina; Sarsina purpurata Linnaeus, Phalaena; Rhyparia purpurea Linnaeus, Phalaena; Rhyparia purpurescens Matsumura, Dokuga; Dokuga pusilla Kiriakoff, Leptoceryx; Leptoceryx pygmaeodes Turner, Nesotropha; Nesotropha pylotis Drury, Phalaena; Calepidos pyralidina Kiriakoff & Viette, Fanambana; Fanambana pyraliformis Moore, Danaka; Danaka

pyraloides Walker, Isionda; Isionda pyraloides Walker, Scotura; Scotura pyrgion Druce, Amaxia; Antaxia pyrrha Druce, Heterocampa; Cargida pyrrha Schaus, Dycladia; Mesothen pyrrhias Kiriakoff, Pyrsopsyche; Pyrsopsyche pyrrhocrocis Felder & Rogenhofer, Colletria; Colletria pyrrhopa Meyrick, Hestiarcha; Hestiarcha pyrrhopsamma Hampson, Prinasura; Prinasura pyrrhopyga Walker, Eucereon; Thysanoprymna pyrrhorhoea Hübner, Hyelosia; Metastatia pythia Druce, Idalus; Thyromolis

quadra Linnaeus, Phalaena; Lithosia quadrata Walker, Marthula; Marthula quadrata Walker, Procodeca; Procodeca quadrata Walker, Zigira; Zigira quadricincta Kaye, Tigridania; Tigridania quadricolor Walker, Mevania; Mevania quadrifasciata Walker, Gerba; Gerba quadrifasciatus Cockerell & Le Veque, Chionaemopsis;

Chionaemopsis
quadrigutta Walker, Euchromia; Ichoria
quadrimacula Mabille, Naclia; Thyrosticta
quadriplaga Moore, Mithuna; Mithuna
quadriplagiata Walker, Pseudomesa; Pseudomesa
quadripunctaria Poda, Phalaena; Euplagia
quadripunctata Aurivillius, Stenarctia; Stenarctia
quadripunctata Gaede, Scranciola; Scranciola
quadripunctata Wallengren, Orgyia; Bracharoa
quadristrigata Felder, Isoctenia; Isoctenia
quadristrigata Walker, Pseudomya; Dixophlebia
quenseli Paykull, Bombyx; Grammia
quincuncialis Holloway, Codalithia; Codalithia
quinquemaculata Kiriakoff, Mesonadata; Mesonadata
quinquepunctata Möschler, Eudasychira; Eudasychira
quinta Kiriakoff, Automolis; Pinheyata

rabdina Butler, Leptosphetta; Leptosphetta radama Mabille, Amblythyris; Amblythyris radians Walker, Apantesis; Apantesis radiata Bethune-Baker, Parazana; Parazana radiosa Pagenstecher, Amsacta; Amsactarctia rafflesi Moore, Eumeta; Ambadra ragonotalis Snellen, Sybrida; Kawiella ragua Druce, Lithosia; Leucorhodia ramburi Zerny, Episema; Episema ramosula Staudinger, Lithosia; Paidina rangei Gaede, Phlyctaenogastra; Phlyctaenogastra ranruna Matsumura, Asuropsis; Asuropsis ratovosoni Viette, Euproctis; Viettema rattrayi Rothschild, Metapiconoma; Metapiconoma rava Lucas, Scaeodora; Pelobrochis rectangulata Kenrick, Phryganopteryx; Paraphrygia recurviloba Rothschild, Garudinodes; Megalobosia recussa Kiriakoff, Otpada; Otpada regina Cramer, Phalaena; Idalia rejecta Geyer, Centronia; Phavaraea relictus Kozhanchikov, Ocneropsis; Gissarus rema Druce, Tuerta; Tuertella remelana Moore, Lithosia; Vamuna remicauda Butler, Tarsolepis; Tarsolepis remota Walker, Penora; Penora rendalli Distant, Dasychira; Homochira repanda Walker, Neritos; Neritos resumens Walker, Psaphida; Psaphida

reticulata Felder, Cyme; Cyme reticulata Moore, Lithosia; Dolgoma reticulata Walker, Anaphe; Anaphe reussi Gaede, Pleuromnema; Pleuromnema reversa Schaus, Epitalara; Epitalara reversa Walker, Lithosia; Ocrosia rhabdophora Hampson, Pydna; Roppa rhabdophora Turner, Thallarcha; Phenacomorpha rhebus Cramer, Sphinx; Diospage rhoda Druce, Phaegoptera; Pseudohemihyalea rhodapicata Holland, Heteronygmia; Heteronygmia rhodesi Bethune-Baker, Paradrallia; Paradrallia rhodobroncha Felder, Isostola; Isostola rhodographa Hampson, Araeomolis; Araeomolis rhodopepla Felder, Sarothropyga; Sarothropyga rhodophaea Walker, Aloa; Teracotona rhodophila Walker, Eucereon; Heliura rica Hübner, Dioptis; Lauron ricini Fabricius, Bombyx; Olepa rinaria Moore, Redoa; Pendria rivata Hampson, Stauropus; Chlorocalliope roberti Schaus, Hardingia; Hardingia roboris Fabricius, Noctua; Chaonia rogenhoferi Schaus, Blera; Sericochroa rosacea Hampson, Pydna; Pydnella roscida [Denis & Schiffermüller], Noctua; Endrosa rosea Beutenmüller, Euhyparpax; Euhyparpax rosea Fabricius, Bombyx; Calligenia rosea Schaus, Lampruna; Lampruna rosea Walker, Sangata; Sangata roseata Walker, Castabala; Castabala roseireta Hampson, Eriostepta; Eriostepta roseivenata Röber, Protolithosia; Protolithosia roseiventris Kiriakoff, Arciera; Arciera rosenbergi Rothschild, Poliopasta; Pseudophaio roseoalba Walker, Heorta; Heorta roseothorax Rothschild, Diarhabdosia; Schalotomis rosinaria Hampson, Chadisra; Rosinella rossii Curtis, Laria; Byrdia rostrata Dognin, Caelicola; Caelicola rostrata Wallengren, Euchelia; Diota rotundata Hulstaert, Metamicroptera; Metamicroptera rotundata Matsumura, Formofentonia; Formfentonia rotundata Matsumura, Marushachia; Marushachia rotundata Walker, Teara; Iropoca rougeoti Laporte, Batuana; Batuana ruatana Druce, Antichloris; Bombopsyche rubea [Denis & Schiffermüller], Bombyx; Oeneria; rubella Grote, Burtia; Burtia ruber Hampson, Arboricornus; Arboricornus rubicunda [Denis & Schiffermüller], Noctua; Miltochrista rubida Walker, Anigraea; Anigraea rubiginea Walker, Besaia; Besaia rubra Neumoegen, Antarctia; Diacrisia rubra Walker, Anace; Anace rubrana Walker, Pontala; Pontala rubriceps Hampson, Haemanota; Haemanota rubriceps Mabille, Chelonia; Parachelonia rubricollis Linnaeus, Phalaena; Atolmis rubricosta Walker, Cluaca; Cluaca rubrifera Walker, Caranusca; Caranusca rubrilineata Bethune-Baker, Metarctia; Rhabdomarctia rubripes Blanchard, Mallocephala; Mallocephala rubriplaga Walker, Ameles; Aemilia rubripuncta Hampson, Metarctia; Metarhodia rubripuncta Joannis, Billetia; Billetia rubrizonea Hampson, Eugraphosia; Eugraphosia rubroflava Hampson, Darantoides; Darantoides rubrostriata Aurivillius, Metarctia; Callobalacra rubrosuffusa Kiriakoff, Elsa; Rhipidarctia rufeola Hampson, Hypagoptera; Hypagoptera rufeola Hampson, Stenosia; Stenosia

rufescens Hampson, Micraroa: Micraroa rufescens Walker, Gadiana; Gadiana rufescens Walker, Metarctia; Metarctia rufibarbis Kiriakoff, Barbarossula; Barbarossula ruficeps Kiriakoff, Diplonychodes; Diplonychodes ruficollis Donovan, Eudesmia; Eudesmia ruficollis Schaus, Talara; Paraprepia ruficornis Hufnagel, Bombyx; Chaonia rufifemur Walker, Polymona; Polymona rufifrons Moore, Churinga; Churinga rufina Kiriakoff, Paulisana; Paulisana rufipex Hampson, Paradoxosia; Paradoxosia rufiventris Schaus, Aclytia; Neacerea rufizonata Hampson, Thermograpta; Thermograpta rufodiscalis Rothschild, Godasa; Antigodasa rufotincta Hampson, Omichlis; Omichlis rufotincta Rothschild, Manoba; Celamodes rufoviridis Walker, Idalus; Chlorhoda rufovittata Aurivillius, Anticyra; Parazana russula Grote, Pseudorgyia; Aretypa russula Linnaeus, Phalaena; Diacrisia rutilus Stoll, Sphinx; Ordishia

sabulosa Felder, Agarista; Neotuerta sachalinensis Matsumura, Kara; Parascotia sacrifica Hübner, Eucharia; Dysschema sadima Schaus, Lithosia; Parablavia sagada Semper, Trichaeta; Neeressa sagana Druce, Heterocampa; Farigia sagenaria Wallengren, Setina; Tumicla saginaea Turner, Calligenia; Gymnasura sagittaria Lucas, Chlenias; Ecnomodes saitonis Matsumura, Quadricalcarifera; Hypostauropus sakania Berio, Poneopirga; Poneopirga saleuta Meyrick, Parelictis; Parelictis salicis Boisduval, Phoegoptera; Phoegoptera salicis Germar, Pterostoma; Pterostoma salicis Lamarck, Furcula; Furcula salicis Linnaeus, Phalaena; Leucoma salmoni Druce, Coreura; Anticoreura salona Druce, Phya; Skaphita salsoma Swinhoe, Zethes; Focillistis salvini Felder, Dialephtis; Dialephtis samarinda Kiriakoff, Pseudoturnaca; Pseudoturnaca sanctaejohannis Walker, Eudryas; Epithisanotia sangala Druce, Tuina; Meterythrosia sangarida Stoll, Phalaena; Curoba sangaris Wallengren, Buthysia; Buthysia sanguiceps Druce, Chrysostola; Sphecopsyche sanguiceps Hampson, Micragra; Micragrella sanguinea Butler, Cisthene; Hypermaepha sanguinea Inoue, Nipponasura; Nipponasura sanguinea Walker, Glaucopis; Phoenicoprocta sanguineata Walker, Halysidota; Hyperthaema sanguinolenta Cramer, Phalaena; Ernassa sanguipuncta Schaus, Marecidia; Marecidia sannio Linnaeus, Phalaena; Diacrisia saphira Staudinger, Agyrta; Cacostatia saskai Berio, Bifrontipta; Bifrontipta saturata Felder, Tineopsis; Tineopsis saturata Walker, Gangamela; Gangamela saturata Walker, Hypsa; Methypsa satyroides Felder, Stenoplastis; Stenoplastis saucia Walker, Phaloesia; Phaloesia saundersi Moore, Calpenia; Calpenia scapularis Herrich-Schäffer, Copaena; Copaena scapulosa Wallengren, Eutaenia; Paralacydes sceletaria Dyar, Bahaia; Bahaia schistacea Kiriakoff, Odontosiana; Odontosiana

schistacea Rothschild, Procrimima; Prolobosia schistaceoplagiata Rothschild, Talara; Neothyone schwarziorum Dyar, Ozodania; Ozodania scieropis Meyrick, Strepsimanes; Strepsimanes scintillans Walker, Somena; Somena sciocosma Meyrick, Symmetrodes; Symmetrodes scissa Warren, Oricia; Cleptophasia scoparioides Strand, Pseudoedaleosia; Pseudoedaleosia scoteola Hampson, Neoblavia; Neoblavia scurrilis Wallengren, Psilura; Oreinobia scutellata Wallengren, Lithosia; Sozusa secunda Ershov, Phothedes; Iranada securis Hübner, Psalis; Psalis selene Felder, Microgiton; Microgiton selenitica Esper, Phalaena; Gynaephora semiaurata Walker, Neola; Neola semicincta Walker, Alope; Nagunda semiclusa Walker, Coutha; Coutha semifalcata Dognin, Dusponera; Dusponera semifascia Hampson, Trichogyia; Trichogyia semiferrea Hampson, Chadisra; Trincomala semiflava Hampson, Chadisra; Stemmatophalera semifusca Turner, Arrhythmica; Arrhythmica semifusca Walker, Orgyia; Ocybola semihyalina Hampson, Dendrophleps; Dendrophleps semiluna Walker, Euchromia; Fregella semilutea Walker, Oxytenis; Staetherinia semimarginata Dognin, Dialephtis; Dialephtis semiochrea Butler, Xanthodule; Xanthodule semiopalina Felder, Ischnognatha; Ischnognatha semiramis Staudinger, Arctia; Nebrarctia semirufa Walker, Halvsidota: Premolis semirufescens Walker, Hatima; Hatima semistria Walker, Dioptis; Erchia semivaria Walker, Halysidota; Tessellarctia senescens Kiriakoff, Mesaeschra; Mesaeschra senex Walker, Ptilomacra; Ptilomacra separata Walker, Apyre; Apyre serena Karsch, Galona; Galona seriatopunctata Matsumura, Disparia; Disparia sericea Felder, Hypolochma; Hypolochma sericea Moore, Stilpnotia; Charala sericea Rothschild, Lambula; Lambulodes sericea Schaus, Hyalarctia; Hyalarctia sericea Walker, Cicynna; Cicynna serrata Thunberg, Bombyx; Peridea sertata Berg, Halysidota; Tessella seruba Herrich-Schäffer, Charidea; Thalesa serva Walker, Lithosia; Gandhara sesamiodes Hampson, Notoxantha; Notoxantha sesiiformis Walker, Glaucopis; Sarosa sexguttata Tessmann, Opoboa; Opoboa seychellarum Fryer, Mahensia; Mahensia seydeli Hering, Laeliophila; Laeliophila shaowuica Daniel, Microlithosia; Microlithosia shepherdi Newman, Termessa; Termessa sibyllae Druce, Josiodes; Pseudoricia sicciaecolor Hulstaert, Nolinophanes; Nolinophanes sicciana Hampson, Nilgiricola; Nilgiricola sidae Fabricius, Noctua; Godasa sideridis Kiriakoff, Bireta; Ottachana sigmata Butler, Phalera; Mesophalera signata Aurivillius, Rhynchophalera; Rhynchophalera signata Dognin, Astapa; Astapa signata Kiriakoff, Eutricholoba; Eutricholoba signata Walker, Ernolatia; Ernolatia signatoria Christoph, Ocneria; Uliolepis signatura Walker, Percote; Percote signifera Walker, Naroma; Naroma sikkima Moore, Celeia; Neophyta sikkimensis Moore, Stauropus; Mesostauropus siletti Walker, Numenes; Numenes

silvandra Stoll, Phalaena; Hypsa silvicola Holloway, Pectinidia; Pectinidia similis Walker, Chilena; Chilena simplex Aurivillius, Synogdoa; Synogdoa simplex Butler, Pseudonaclia; Micronaclia simplex Gaede, Eurystaura; Romalina simplex Rothschild, Idalus; Haplonerita simplex Walker, Antheua; Antheua simplex Walker, Corymbia; Lobogona simplex Walker, Ogoa; Ogoa simplex Walker, Pagara; Pagara simplex Walker, Savara; Bucaea simplex Walker, Thyone; Euthyone simplicior Matsumura, Neasuroides; Neasuroides simulans Butler, Epilacydes; Epilacydes sinensis Walker, Anzabe; Anzabe singularis Aurivillius, Paraphlebs; Paraphlebs singularis Roepke, Xanthomaenas; Xanthomaenas singularis Schaus, Goaxis; Goaxis sinistra Schaus, Coxeya; Coxeya sinuata Toxopeus, Imaida; Imaida sinuata Wallengren, Setina; Mosoda sippia Plötz, Naclia; Meganaclia sisenna Druce, Evius; Parevia sjostedti Aurivillius, Leptonadata; Leptonadata sjostedti Aurivillius, Stemmatophalera; Stemmatophalera slossonii Dyar, Ceruridia; Ceruridia smaragdina Walker, Eunomia; Sesiura smaragdiplena Walker, Exaereta; Oxoia smerinthoides Walker, Corymbia; Corymbia smithii Holland, Parauchmis; Parauchmis snelleni Ritsema, Pseudoblabes; Blabioides sobria Schaus, Apocerea; Apocerea sobria Walker, Arcilasisa; Arcilasisa sobria Walker, Dioptis; Erbessa sobria Walker, Halone; Halone sobria Walker, Misogada; Misogada sobria Walker, Xeniara; Xeniara socorra Schaus, Vulsinia; Vulsinia solaris [Denis & Schiffermüller], Noctua; Acontia solicauda Butler, Heliura; Heliura solox Dognin, Parascepsis; Parascepsis sommeri Hübner, Crino; Tarsolepis sordida Aurivillius, Lophilema; Lophilema sordida Aurivillius, Rhypopteryx; Rhypopteryx sordida Birket-Smith, Archithosia; Architesmia sordida Butler, Eudoliche; Phaulosia sordida Felder, Ischnocampa; Ischnocampa sordida Felder, Phryganopsis; Phryganopsis sordida Walker, Eunomia; Pezaptera sordida Wileman, Fentonia; Disparia sordidus Walker, Purius; Purius sparsata Gaede, Antizana; Antizana sparshalli Curtis, Arcturus; Trichiocercus sparsipennis Walker, Hemiceras; Epicoria speciosa Drury, Phalaena; Pseudhypsa speciosa Schaus, Euzeugapteryx; Euzeugapteryx spectabilis Tauscher, Noctua; Acymba specularis Herrich-Schäffer, Phaegoptera; Bertholdia speculigera Kiriakoff, Hypambadra; Hypambadra sperchius Cramer, Sphinx; Euchromia sphingidea Perty, Glaucopis; Euplesia sphingiformis Clemens, Ormetica; Ormetica sphingiformis Moore, Dudusa; Dudusoides spilosomata Walker, Thyrgorina; Thyrgorina spilosomoides Walker, Bithra; Bithra spiritalis Distant, Cerura; Notocerura spissicornis Mabille, Ichthyura; Ambina splendens Butler, Crambomorpha; Paraona splendens Möschler, Eulophopteryx; Eulophopteryx splendida Rambur, Orgya; Clethrogyna spurca Dyar, Anaene; Anaene

spurca Rambur, Omochroa; Omochroa squalida Dognin, Proanita; Proanita squamicornis Felder, Zygaenopsis; Zygaenopsis squamipunctum Felder, Pterygosoma; Oenosandra squamosa Walker, Lopera; Lopera statices Linnaeus, Sphinx; Adscita staudingeri Saalmüller, Fodinoidea; Fodinoidea stefanellii Giacomelli, Cyanohypsa; Cyanohypsa steinbachi Rothschild, Rifargia; Grafiria stellans Draudt, Xantharete; Xantharete stellata Dognin, Pachasura; Pachasura stellata Guérin-Méneville, Agarista; Arnophila stelligera Schaus, Marthantia; Marthantia steniptera Hampson, Stauropus; Epicerura stenopa Meyrick, Chiriphe; Stenarcha stenoptera Turner, Scythrophanes; Scythrophanes stigmatica Janse, Phycitimorpha; Phycitimorpha stilbioides Moore, Adrepsa; Adrepsa straeleni Debauche, Prytania; Prytania straminea Herrich-Schäffer, Diastema; Antheua straminea Moore, Ceira; Cutuza stretchii Edwards, Scotogramma; Scotogramma striata Aurivillius, Grammoa; Grammoa striata Collenette, Jacksonia; Jacksoniana striata Druce, Heterocampa; Urgedra striata Linnaeus, Phalaena; Spiris striatella Matsumura, Tensha; Tensha strigata Aurivillius, Dasylaelia; Dasylaelia strigata Aurivillius, Ramesa; Bisolita strigata Walker, Secusio; Secusio strigicosta Walker, Darabitta; Darabitta strigifascia Moore, Niganda; Niganda strigifera Druce, Stidzaeras; Stidzaeras strigilata Saalmüller, Macrobrochis; Phryganopteryx strigillata Rothschild, Elysius; Graphelysia strigipennis Moore, Locharna; Locharna strigosa Butler, Trichromia; Lycomorphodes strigosa Walker, Arctia; Calidota strigosa Walker, Euchromia; Rhipha strigosa Walker, Rethona; Rethona strigosa Walker, Rosama; Rosama strigosus Grünberg, Paramaenas; Paramaenas strigularis Walker, Belciana; Dialephtis structa Walker, Pallene; Prinasura stygne Walker, Josia; Phavaraea styphlopis Turner, Polychoa; Polychoa styracis Felder, Desmoloma; Desmoloma suava Staudinger, Pygopteryx; Pygopteryx subaequa Kiriakoff, Eurystaurella; Eurystaurella subapicalis Walker, Glaucopis; Lasioprocta subapicalis Walker, Pericopis; Addua subargentea Felder, Leucoma; Candidata subargentea Walker, Marane; Marane subaurata Walker, Glaucopis; Trianeura subcarnea Warren, Turnaca; Rhabdoturnaca subcrinita Dognin, Pilodonta; Pilodonta subcyanea Walker, Euchromia; Procalypta subfascia Walker, Hypsa; Anagnia subformicina Bethune-Baker, Ceryx; Paralaethia subgeneris Strand, Stauropus; Quadricalcarifera subgriseola Strand, Eilema; Ovenna subinanis Walker, Topomesa; Topomesa subjecta Walker, Cisthene; Cisthene subjecta Walker, Echlida; Echlida sublactigera Walker, Euchontha; Euchontha subluna Walker, Antona; Antona submacula Walker, Virbia; Microgiton submarginata Rothschild, Trischalis; Palaeotype submarginata Schaus, Ascaptesyle; Ascaptesyle submarginata Walker, Redoa; Redoa subnigra Bethune-Baker, Porthmeia; Porthmeia subnotata Walker, Orvasca; Orvasca

subovalis Turner, Nycteropa; Nycteropa subquadrata Herrich-Schäffer, Aganopis; Aganopis subrosea Matsumura, Shachia; Shachia subrosea Walker, Anthora; Laelia subserena Kiriakoff, Bireta; Odnarda substrigosa Walker, Caviria; Caviria substyxana Kiriakoff, Apistaeschra; Apistaeschra subterminalis Kiriakoff, Mainiella; Mainiella subterminalis Kiriakoff, Medanella: Medanella subtilis Felder, Anatolis; Anatolis subumbrata Holland, Lepista; Archilema subvarda Schaus, Arhabdosia; Arhabdosia subvelutina Kiriakoff, Oligaeschra; Oligaeschra subviridis Walker, Boreconia; Boreconia subvitrea Walker, Leucoma; Kanchia suffusa Hampson, Lyclene; Nephelomilta suffusa Walker, Ricine; Ricine sumatrana Kiriakoff, Grangulina; Grangulina sumptuosa Dognin, Thyana; Thyana sundana Roepke, Parafentonia; Parafentonia superba Janse, Arattatha; Arattatha superpulverea Dyar, Spodarctia; Spodarctia suprema Schaus, Dukinfieldia; Dukinfieldia surgens Walker, Borseba; Borseba suriga Schaus, Turnaca; Suriga surinamensis Rothschild, Automolis; Aphyarctia suzukii Matsumura, Suzukia; Suzukiana sylvestris Schaus, Peroara; Peroara synemonistis Strand, Pemphigostola; Pemphigostola syntomina Butler, Caryatis; Euthoracia syntomoides Butler, Callitomis; Callitomis sypnoides Butler, Bocana; Palaeocoleus

tabernaria Kiriakoff, Andocidia; Andocidia taeniata Guérin-Méneville, Chelonia; Ormetica taeniosoma Hering, Laelioproctis; Laelioproctis taiwana Wileman, Chrysorabdia; Heliorabdia taiwana Wileman, Tarsolepis; Tarsolepis takamukuana Matsumura, Megashachia; Megashachia takamukui Matsumura, Acmeshachia; Acmeshachia takamukui Matsumura, Takadonta; Takadonta takaonis Matsumura, Hiradonta; Hiradonta tamahonis Matsumura, Euproctis; Epeuproctis tamarix Kiriakoff, Pycnographa; Pycnographa tamsi Collenette, Euproctis; Miccotrogia tanda Bethune-Baker, Epicerura; Epicerura tapina Dyar, Altimaenas; Altimaenas taprobanis Walker, Aemene; Aemene tarsalis Walker, Glaucopis; Pompilopsis tasmanica Hampson, Thermeola; Thermeola taurica Herrich-Schäffer, Heliothis; Philareta telamonophorum Prittwitz, Diptilon; Diptilon telesilla Druce, Neozatrephes; Neozatrephes temenus Stoll, Sphinx; Thrinacia temperata Walker, Olapa; Olapa tendinosa Hübner, Phaeochlaena; Phaeochlaena tenebrifera Hampson, Pseudoscolia; Parascolia tenebriplaga Walker, Clostra; Clostra tenebrosa Dognin, Acyclania; Acyclania tenebrosa Moore, Phalera; Scotodonta tenebrosa Walker, Apistosia; Nyctosia tenebrosa Walker, Glaucopis; Phacusa teneiformis Walker, Syntomis; Trichaeta tenera Hübner, Cycnia; Cycnia tenuis Butler, Hylemera; Hylemera tephra Hübner, Dasychira; Dasychira terminalis Kiriakoff, Phalerina; Phalerina terminalis Walker, Apistosia; Protosia terminalis Walker, Euchromia; Napata

terminalis Walker, Eutane; Eutane terminalis Walker, Pionia; Correbidia terminata Walker, Clamara; Clamara terminicana Kiriakoff, Bireta: Liccana teroensis Collenette, Paramarbla; Paramarbla terrifica [Denis & Schiffermüller], Bombyx; Harpyia tessellaris J.E. Smith, Phalaena; Halysidota testacea Aurivillius, Pseudapiconoma; Pseudapiconoma testacea Moore, Harapa; Harapa testacea Roepke, Kawiella; Kawiella testacea Walker, Cropera; Cropera testacea Walker, Pydna; Eupydna testacea Walker, Spilosoma; Eyralpenus tetragona Walker, Lithosia; Thysanoptyx tetrasticta Kiriakoff, Parasoloe; Parasoloe tettigonioides Heylaerts, Leptothrix; Chionaema textor Harris, Hyphantria; Hyphantria thalassina Hampson, Stauropus; Heraia tharis Stoll, Phalaena; Disphragis thaumasta Franclemont, Apocrisias; Apocrisias thaumatopoeides Schultze, Dasychira; Dyasma theon Druce, Amaxia; Neaxia thermidoides Talbot, Thermidarctia; Thermidarctia thetis Klug, Eyprepia; Daritis thoracica Edwards, Eulithosia; Chichimeca thoracica Walker, Euchromia; Psoloptera thricophora Hampson, Nodozana; Nodozana thyretiformis Wallengren, Naclia; Ceryx thyria Druce, Desmidocnemis; Hyalomis tibetensis Daniel, Lacydoides; Lacydoides tigniferum Felder, Ichthyosoma; Trichomaplata tigrina van Eecke, Sylescaptia; Sylescaptia tigrinus Hampson, Leucopardus; Leucopardus timon Hübner, Bombyx; Pygaera timoniorum Bremer, Pygaera; Gonoclostera tina Walker, Euchromia; Bodosa tinctaria Hübner, Geometra; Crocota tineoides Walker, Mantala; Mantala tipulina Hübner, Glaucopis; Pseudomya tiresias Cramer, Phalaena; Notophyson tolumnensis Herrich-Schäffer, Trichela; Trichela tomponis Matsumura, Meteuproctis; Meteuproctis torniplaga Jones, Clemensia; Neomulona torta Walker, Corcura; Corcura tortricella Walker, Macaduma; Macaduma tortriciformis Möschler, Afrida; Afrida tortriciformis Zeller, Rhagophanes; Rhagophanes tortricina Felder, Dylomia; Dylomia tortricoides Walker, Chatracharta; Chatracharta tortuosa Moore, Neochera; Antichera tosta Kiriakoff, Aoba; Aoba tosta Walker, Ramesa; Ramesa toxopei Rocpke, Allochrista; Allochrista toxophora Turner, Psapharacis; Psapharacis tranducta Walker, Ecregma; Ecregma transcisa Walker, Phaeochlaena; Euchlaenidia transiens Hampson, Tuerta; Hypotuerta transiens Walker, Spilosoma; Phissama translucida Butler, Pachylaelia; Pachylaelia translucida Griveaud, Cadorela; Cadorela translucida Oberthür, Leucoma; Stracilla transversa Gaede, Ogovodonta; Ogovodonta transversa Walker, Cyllene; Padenia transversa Walker, Sermyla; Sermyla treitschkii Frivaldszky, Heliothis; Philareta triangularis Kiriakoff, Bireta; 'Dypna triangulifera Felder, Sphenoptera; Asphenoptera triangulum Draudt, Phyllopalpia; Phyllopalpia trichopteroides Walker, Chamaita; Chamaita trichosa Hampson, Stauropus; Daulopaectes trichosticha Hampson, Hyperaeschra; Anthyperaeschra tricolor Butler, Isorropus; Isorropus

tricolor Felder, Heterodontia; Heterodontia tricolor Herrich-Schäffer, Torycus; Torycus tricolor Sulzer, Phalaena; Aphisaon tricolor Walker, Scaptesyle; Scaptesyle tricolora Sulzer, Phalaena; Thebrone tridesma Turner, Atelophleps; Atelophleps trifascia Bethune-Baker, Paradohertya; Paradohertya trifasciata Burmeister, Halysidota; Sellota trigona Leech, Nyctemera; Micrarctia trigutta Walker, Isine; Isine trigutta Walker, Soloe; Soloe trilineata Walker, Agabra; Agabra trimacula Hampson, Setinochroa; Eurosia trimaculata Hampson, Monotaxis; Monosyntaxis trimenii Felder, Aegocera; Agoma trimenii Felder, Dyphlebia; Dyphlebia trinitas Strand, Eilema; Zobida tripars Walker, Sureta; Sureta triplaga Hampson, Cisthene; Costalimaida triplaga Walker, Docyna; Docyna tripunctaria Linnaeus, Phalaena; Orphanos tripunctata Gaede, Psegmaphora; Psegmaphora tripunctata Griveaud, Tsirananaclia; Tsirananaclia triseriata Herrich-Schäffer, Gnophaela; Paraethria tristicolor Gaede, Antheua; Castastygne tristis Lewin, Bombyx; Epicoma tristis Schaus, Nodoza; Nodoza tristis Staudinger, Notodonta; Allodonta tristriata Bryk, Disconeura; Disconeura tritophus [Denis & Schiffermüller], Bombyx; Tritophia trivittata Walker, Dioptis; Aphra trocta Karsch, Barobata: Barobata troglodyta Graeser, Pygaera; Micromelalopha trogonoides Walker, Aclytia; Ixylasia tropica Walker, Euchromia; Sciopsyche trotschi Druce, Calonotos; Xanthopleura trucidata Butler, Xanthodura; Xanthodura truncata Hampson, Parashada; Parashada truncata Walker, Oricia; Oricia tumida Walker, Tegulata; Tegulata turbata Butler, Gogane; Raphipeza turbida Möschler, Phedosia; Phedosia turcosa Retzius, Adscita; Adscita turlini Toulgoët, Seydelia; Seydelia typica Linnaeus, Phalaena; Naenia typica Strand, Syntomerea; Syntomerea

uelleburgensis Strand, Eilema; Palilema ulmi [Denis & Schiffermüller], Noctua; Dicranura umber Cramer, Phalaena; Pseudapistosia umbra Kiriakoff, Pheosilla; Pheosilla umbrata Dognin, Euclemensia; Euclemensoides umbrifera Felder, Crambomorpha; Crambomorpha umbrifera Walker, Dioptis; Pseuderbessa umbrina Kiriakoff, Sagamora; Sagamora umbrosa Kiriakoff, Astaura; Astaura umbrosa Leech, Lophopteryx; Inouella unca Dognin, Hamidonta; Hamidonta uncina Hübner, Gonodonta; Gonodonta undulata Kiriakoff, Sawia; Sawia undulifera Hampson, Leucolopha; Leucolopha undulosa Walker, Acontia; Philenora unicolor Hopffer, Crocota; Ochrota unicolor Packard, Lochmaeus; Stematocampa unicornis Smith, Phalaena; Coelodasys unidentata Bryk, Pseudonerice; Pseudonerice unifascia Rothschild, Padenodes; Padenodes unifascia Walker, Andala; Andala unifascia Walker, Thosea; Thosea

uniformis Gaede, Epiphalera; Epiphalera uniformis Hampson, Lophoneura; Lophoneura uniformis Möschler, Thagona; Thagona uniformis Schaus, Haematomis; Haematomis uniformis Swinhoe, Lophopteryx; Drapetides unilinea Walker, Lymphorta; Lymphorta unimacula Warren, Tanaostyla; Euforbesia unipectinata Aurivillius, Lepidilema; Lepidilema unipunctata Möschler, Laelia; Conigephyra unita [Denis & Schiffermüller], Noctua; Manulea univitta Hampson, Euryptidia; Euryptidia urania Püngeler, Phragmatobia; Oroncus uranophila Walker, Eucyane; Agyrtidia urapterides Walker, Ciaca; Ciaca urophora Herrich-Schäffer, Cercophora; Cercophora ursina Schaus, Phaegoptera; Pseudotessellarctia ursipes Walker, Zama; Zamana ursula Stoll, Phalaena; Epidesma ussuriensis Püngler, Ochrostigma; Chadisroides ustulata Felder, Eriphia; Eriphioides

v-nigrum Fabricius, Bombyx; Arctornis vacillans Walker, Amphissa; Phissama vacillans Walker, Eunomia; Phoenicoprocta vadoni Toulgoët, Asura; Proxhyle vagans Boisduval, Arctia; Rhagonis vagesa Moore, Lithosia; Capissa valdiviesoi Dognin, Orthosoma; Meragisa varana Moore, Lithosia; Tarika variabilis Schaus, Goacampa; Goacampa varians Bethune-Baker, Damata; Parachadisra variegata Aurivillius, Phalera; Alenophalera variegata Moore, Heterocampa; Chadisrina variegata Rothschild, Nudaria; Crocodeta variegata Villers, Phalaena; Ptilophora variegata Walker, Eudule; Syntomidopsis variegata Walker, Teara; Aglaosoma variegata Wileman, Fentonia; Epifentonia varii Kiriakoff, Anticleapa; Anticleapa varipes Walker, Euchromia; Mystrocneme varipes Walker, Glaucopis; Leucotmemis varipes Walker, Hysibada; Hysibada velitaris Hufnagel, Phalaena; Drynobia veltini Dognin, Lirimiris; Dognina velutina Birket-Smith, Asythosia; Asythosia velutina Oberthür, Drymonia; Togepteryx velutinum Butler, Mitradaemon; Mitradaemon venata Aurivillius, Dionychoscelis; Dionychoscelis venata Aurivillius, Poecilarctia; Poecilarctia venata Butler, Episcepsis; Episcepsis venata Butler, Melanaema; Melanaema venata Rothschild, Baritius; Haemaphlebiella venata Swinhoe, Cadurca; Cadurca veninotata Warren, Xenomigia; Xenomigia venosa Hering, Stilpnaroma; Stilpnaroma venosa Kiriakoff, Euanthia; Euanthia venosa Moore, Carbisa; Carbisa venosa Moore, Prabhasa; Prabhasa venosa Walker, Ctenucha; Philoros venosa Walker, Hebena; Hebena venosata Hering, Lepidolacipa; Lepidolacipa venosata Walker, Gazalina; Gazalina ventralis Schaus, Machaeraptenus; Machaeraptenus venusta Lucas, Mosoda; Stenoscaptia verania Druce, Phaloe; Pseudophaloe verbasci Fabricius, Cossus; Neoharpyia vermiculata Grote, Omoiala; Omoiala vernalis Gaede, Pseudohoplitis; Pseudohoplitis veronica Oberthür, Naclia; Vitronaclia

vesparis Butler, Poecilosoma; Loxophlebia vesperalis Nakamura, Nepalia; Nepalia vespiformis Druce, Metaxanthia; Metaxanthia vespina Harris, Mastigocera; Horama vespoides Strand, Pseudosphecosoma; Pseudosphecosoma vexata Felder, Antiotricha; Antiotricha viettei Collenette, Crorema; Madema viettei Kiriakoff, Artanasa; Artanasa viettei Kiriakoff, Ochrocalliope; Ochrocalliope viettei Kiriakoff, Omocerina; Omocerina viettei Laporte, Syrrusoides; Syrrusoides vieui Viette, Rothia; Pararothia vilis Felder, Dicranuropsis; Dicranuropsis vilis Holland, Ctenogyna; Epitrotonotus vilis Walker, Euproctis; Casama villacresi Dognin, Idalus; Neidalia villica Linnaeus, Phalaena; Epicallia vinosa Drury, Phalaena; Hyalurga vinula Linnaeus, Phalaena; Cerura viola Butler, Mardara; Mpanjaka violacea Pelletier, Marcipa; Marcipalina violascens Gaede, Sentana; Sentana violascens Herrich-Schäffer, Notodonta; Oligocentria virens Butler, Syntrichura; Syntrichura virescens Matsumura, Aoshachia; Aoshachia virescens Möschler, Pseudonotodonta; Pseudonotodonta virescens Moore, Cadrusia; Cadrusia virescens Moore, Stauropus; Benbowia virgata Packard, Xylinodes; Xylinodes virginalis Boisduval, Chelonia; Platyprepia virginica Charpentier, Sphinx; Ctenucha virgo Butler, Platychasma; Platychasma virgo Herrich-Schäffer, Ctenucha; Ctenuchidia viridata Walker, Lithosia; Chrysorabdia viridescens Kiriakoff, Pararhenea; Pararhenea viridescens Walker, Netria; Netria viridescens Walker, Ripisa; Ripisa viridifusca Walker, Somera; Somera viridinota Hampson, Fentonia; Chloroceramis viridipennis Strand, Stauropussa; Stauropussa viridipicta Kenrick, Dasychira; Rivotra viridipicta Kiriakoff, Panteleclita; Panteleclita viridipulverea Gaede, Chadisra; Chlorochadisra viridirosea Dognin, Libyssa; Eudognina viridis Plötz, Caryatis; Carpostalagma viridisparsa Griveaud, Soganaclia; Soganaclia viridissima Kiriakoff, Eramos; Eramos viridissimus Bethune-Baker, Stauropus; Chlorostauropus viriditincta Strand, Someropsis; Someropsis vitrina Oberthür, Pseudapiconoma; Neophemula vittata Gaede, Allodonta; Corinella vittata Hulstaert, Stenocyttara; Onymapata vittata Möschler, Eudoliche; Eudoliche vittata Staudinger, Notodonta; Paradrymonia vittata Walker, Pheia; Psilopleura vittata Walker, Rigema; Rigema vittata Wing, Trichomaplata; Trichomaplata vittigera Felder, Euplesia; Euplesia vittigerum Blanchard, Compsoprium; Compsoprium vittipalpis Walker, Harma; Hippia vittipes Walker, Arara; viviana Schaus, Dyasia; vocula Stoll, Phalaena; Paralacydes volcancita Dognin, Marcella; Marcella voluptuosa Bryk, Libido; Libido vulcanus Walker, Euchromia; Endera vulgaris Druce, Nola; Eutelesia vulnerata Butler, Hyaleucerea; Pseudohyaleucerea

walkeri Hemiceras, Schaus; Salamboria wallengreni Aurivillius, Caligula; Paralacydes wanderbilti Rego Barros, Wanderbiltia; Wanderbiltia warionis Oberthür, Bombyx; Albarracina wernickei Draudt, Eucereum; Euceriodes weyenberghi Snellen, Woerdenia; Woerdenia whiteleyi Druce, Belemnia; Belemniastis wittei Debauche, Paramyopsyche; Paramyopsyche woodfordii Butler, Hyalaethea; Hyalaethea wukongensis Chou & Lu, Cerviplusia; Cerviplusia

xenopis Lower, Oenone; Diceratucha xenops Kiriakoff, Takwa; Takwa xolotl Schaus, Hapigia; Hapigiodes xylinata Walker, Besida; Besida xylinoides Walker, Edema; Pentobesa xylinoides Walker, Rifargia; Rifargia xylocampoides Walker, Nagidusa; Nagidusa xylogramma Turner, Scaphidriotis; Scaphidriotis

ydatodes Dyar, Afrida; Aresia

xanthodelta Lower, Scoliacma; Poliodule xanthogastra Perty, Glaucopis; Andrenimorpha xanthogramma Hampson, Holochrea; Holochrea xanthomela Druce, Nodozana; Nodozana xanthomelas Walker, Mapeta; Pyralopsis xanthopa Holland, Lepista; Muxta xanthopera Hampson, Euproctis; Pronygmia xanthoperas Hampson, Sphingomorpha; Rougeotiana xanthophila Walker, Menapia; Menapia xanthosoma Dognin, Syntomostola; Syntomostola xanthosoma Hampson, Phacusosia; Phacusosia xanthosoma Wallengren, Phiala; Phiala xanthospila Plötz, Aroa; Mylantria xanthosticta Hampson, Hemihyalea; Machadoia xanthura Schaus, Parablavia; Parablavia xenodora Meyrick, Peronetis; Peronetis

zaba Dyar, Rhabdatomis; Rhabdatomis zapata Schaus, Kalkoma; Praeschausia zatesima Hampson, Poliosia; Ateucheta zatrephica Dyar, Epimolis; Epimolis zavattarii Berio, Paraonagylla; Paraonagylla zelica Stoll, Phalaena; Zelica zelotes Talbot, Barlowia; Barlowia zerbina Stoll, Phalaena; Pericopis zethus Druce, Dasylophia; Theroa zethus Hübner, Pseudosphex; Pseudosphex zeus Schaus, Ptychotricos; Ptychotricos ziczac Linnaeus, Phalaena; Eligmodonta zonata Joannis, Sterrhosia; Sterrhosia

